Efficiency of the EmERGE Pathway to provide continuity of care for Spanish people living with medically stable HIV

Article (Accepted Version)

Beck, E J, Mandalia, S, Yfantopoulos, P, Jones, C I, Bremner, S, Whetham, J, Etcheverry, F, Moreno, L, Gonzalez, E, Merino, M J, Leon, A and Garcia, F (2021) Efficiency of the EmERGE Pathway to provide continuity of care for Spanish people living with medically stable HIV. *Enfermedades Infecciosas y Microbiología Clínica*. ISSN 0213-005X

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/102949/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Efficiency of the EmERGE Pathway to provide Continuity of Care for Spanish People Living with Medically Stable HIV

Background:

The 2020 Spanish Covid19 epidemic reiterated the need for people living with chronic diseases to remain in touch with care-givers, especially during emergencies [1]. The global HIV community increasingly wants to track, monitor and evaluate the use, cost, outcome and impact of national health services for people living with HIV (PLHIV) [2]. Mobile Health (mHealth), employs wireless technology to deliver health information [3,4] and plays an increasingly important role in linking care and integrating health services [5]. mHealth has become an important tool to respond to the Covid-19 Pandemic [6,7]

Increased global access to anti-retroviral drugs (ARVs) [8], resulted in increased life-expectancy of PLHIV, similar to that of people not living with HIV [9]. This will increase the number of PLHIV, including those aged 50 years or older [10]. Non-HIV cancers, cardiovascular disease and other non-communicable diseases (NCDs) are now the commonest cause of death for PLHIV in high income countries [11], while the incidence and prevalence of NCDs are increasing among PLHIV in low- and middle-income countries [12]. PLHIV will increasingly need to use HIV and non-HIV health and social services and integrated health information systems can assist service integration and make them more cost-effective [5].

The Evaluating mHealth technology in HIV to improve Empowerment and healthcare utilisation: Research and innovation to Generate Evidence for personalised care (EmERGE) Project developed and implemented a new mHealth Pathway in five HIV centres in five European countries: Belgium, Croatia, England, Portugal and Spain. The Pathway includes a mobile health application (App)], which enables PLHIV access to communicate with care-givers and personal health information [13]. Recruitment criteria are described in Table 1.

Table 1 near here

A recent review identified nine communication functions for a mHealth App [4], but missed two important criteria: firstly, the confidentiality and security of the data need to be protected. Secondly, the technology needs to be affordable and efficient (Table 2; [15]).

Table 2 near here

Most mHealth studies have not included the cost of developing and implementing these mHealth Apps, nor their efficiency. The need for efficiency studies was recognized long ago [16,17], but few efficiency studies have been performed since [18].

This before-and-after study calculated the use, cost and cost-effectiveness of HIV services for EMERGE participants one-year pre- and one-year post-implementation of the EmERGE Pathway at the Hospital Clinic-IDIBAPS (HC- IDIBAPS), University of Barcelona, Barcelona, Spain. The specific objectives were: 1) calculate the use of HIV outpatient services at HC- IDIBAPS pre- and post-EmERGE; 2) calculate the unit costs of departments supporting HIV outpatient services; 3) calculate the annual costs of HIV outpatient services pre- and post-EmERGE; 4) calculate the efficiency of implementation of the EmERGE Pathway at HC- IDIBAPS.

Methods

Context

546 participants were followed up between 1st July 2016-30th October 2019 at the Instituto Clinic de Medicina y Dermatología (ICMiD). The ICMiD is a research centre of the HC-IDIBAPS, and provides the infectious diseases service for this Hospital.

At the start of the study, there were three separate outpatient clinics where PLHIV were followed up: General Outpatient Clinic (Dispensario), the HIV Daycare Hospital and the ICMiD Daycare. While the former two conducted face-to-face appointments the latter clinic offered PLHIV virtual appointments. This video-consultation appointment system – Hospital Virtual – had been in operation since 2005 [19]. During the second year of the study the virtual clinic was closed for internal technical reasons. The videoconferencing and sound system in use was old and inadequate. As this could not be upgraded in a timely manner to continue services, this service was therefore closed during the second year of the study. As a consequence, those participants that were recruited to attend the virtual clinic during the first year of the study, were transferred to the two face-to-face clinics during the second year. The virtual clinic was closed in October 2019.

At the time of recruitment EmERGE participants signed an informed consent form, and this visit was the base-line for the analyses. Participants were followed up one year retrospectively pre-EmERGE and prospectively post-EmERGE implementation. All investigative procedures and minor surgical procedures were performed in the ICMiD Daycare Office or HIV Daycare Hospital.

Tests results were transmitted electronically, reviewed and stored in electronic medical records. ARVs were provided free and most participants collected them from the Hospital . HIV outpatient services were supported by the following departments: laboratories, pharmacy, surgical and radiology. The HC-IDIBAPS developed its own method to cover overheads. It provides departments with a fixed additional percentage of their budget to cover the following overheads: cleaning & waste, laundry & clothes, materials, messaging & communication, security, information technology services, legal, administrative, human resources, financial, and managerial services. None of EmERGE participants had inpatient stays during the study period, hence the inpatient ward was not costed. As per EmERGE protocol, one visit per year was an electronic ‘visit’ when the physician reviewed the results of blood tests and sent their recommendation for future treatment via the EmERGE App; the other annual visit, part of routine management, was a face-to-face visit, or virtual visit for the ICMiD Daycare Outpatient participants during their first year of the study.

Data collected

Information on the use, cost and outcome of HIV services by EmERGE participants pre- and post-EmERGE were required to estimate efficiency of the Pathway. This required process information on use of services, the unit costs of the services used, average annual cost of services pre- and post-EmERGE and changes in primary or secondary outcome measures.

Costing Health Facilities: General Principles

The “top-down” or “bottom-up or ingredient-based” methods are the two main costing methods [20,21]. The more detailed “bottom-up” method defines the type and quantity of input or ingredients used to produce the service output. The ‘top-down’ method is easier to perform and determines the total cost of providing services on accounts of past expenditure. This expenditure is then divided by the number of ‘products’ provided during the period. Where possible, the preferred ingredient-based approach was used, but this was dependent on the availability of detailed data.

Process data

The use of services by EmERGE participants in the year pre- and post-EmERGE provided the individual process data, which were obtained from the electronic medical records. They predominantly used outpatient services and therefore the focus of the costing exercise was on the three HIV outpatient services at HC- IDIBAPS.

Unit costs data

A micro-costing study was performed between October-November 2018. The focus was on outpatients' services used by EmERGE participants for the 2018 financial year. The costing study was based on the UNAIDS Costing Manual [20] and UNAIDS Costing Workbook [21]. To estimate the unit costs, workload and financial data were collected from departments supporting EmERGE outpatient services. The individual process data of EmERGE participants identified which services were used and these were costed. The financial data included *staff*, *consumables*, *overhead*, *procedures* and *equipment* costs (SCOPE) [22]. Unit costs of outpatient services were combined with process data to calculate average annual costs of outpatient services for participants pre- and post- EmERGE [23].

Statistical Methods

All participants with a baseline visit, contributed to the data and analyses presented. Summary statistics are presented with point estimates and indication of variability and missing data. Linear mixed models were used to calculate difference in three-monthly averages in CD4 counts and viral load measurements (DAVG; 24). Time weighted changes of CD4 count and viral load were analysed over a two-year period and measured by time point changes during one year before recruitment and one year after EmERGE recruitment [25]. MIXED procedure in SAS was used by fitting routine values of CD4 counts and viral loads results as dependent variables [25]. Independent variables included the fixed effects of study visit time points and time in months. A covariance matrix was used to model the within patient errors. Estimates of effects are based on MIXED models and assume any missing data were missing at random. Trends over time are presented as point estimates derived from the models. Viral load data were transformed logarithmically to stabilise their variance.

The mean number of services used per patient year (MPPY) was calculated using methods employed previously [26-28], based on the following formula:

Equation 1.1 Mean Per Patient Year (MPPY) Calculation Formula

$$M = \frac{\sum_{i=1}^n \sum_{j=1}^k S_{ij}}{\sum_{i=1}^n \sum_{j=1}^k (t_{ij} - t_{i(j-1)})} \times 365$$

Where:

- n = total number of individuals;
- k = day of censoring;
- S_{ij} = use of service by individual i on jth day;
- t_{ij} = number of days of follow up for individual i;
- M = mean of services S per patient-year.

The denominator comprised the total duration of follow up for all patients during a calendar year, one year from baseline and one-year prior to study entry. The data were left censored at

the one-year pre-baseline visit. Post-EmERGE data were right censored at either one year since baseline visit if patients were still under follow-up at one year, or if they had died during follow up then their date of death, or if they were lost to follow up, which ever came first. Numerators were calculated by summing the use of outpatient services (Equation 1.1). Exact Poisson 95% confidence intervals (CIs) were estimated for MPPY based on the distribution of the observed number of outpatient visits divided by the total duration of follow up for all patients during a calendar year. All statistical analyses were performed using SAS version 9.4 software [25].

Average annual costs per patient-year (PPY) of EmERGE outpatient service provision were produced by multiplying the MPPY by their respective unit costs. The total average annual costs for providing services was obtained by adding the annual costs for outpatient visits, tests, procedures and drugs at the HC- IDIBAPS [22,23]. Costings were performed from a societal perspective.

The original focus of the analysis was the use and costs for different HIV outpatient clinics, but as the Virtual Clinic was closed during the second year, an additional analysis was performed to compare the use and cost of participants followed up only in face-to-face clinics (F2F) with those participants that were transferred from the virtual to the face-to-face clinics during the second year of the study (V2F).

Primary and secondary outcome measures

Changes in CD4 count and viral load measurements were primary outcome measures pre- and post-EmERGE. The secondary outcome measures included changes in patient activation, measured by PAM13 [29], and PROQOL-HIV quality of life measures [30], from month 0 (baseline) to month 12 post-EmERGE.

Cost-effectiveness analyses

Incremental cost-effectiveness ratios (ICERs) were calculated based on the average annual costs, primary and secondary outcome measures before and after the implementation of EmERGE using the following formula [31]:

$$ICER = \frac{[Costs A - Costs B]}{[Outcome A - Outcome B]}$$

As the primary and secondary outcome measures did not change substantially between periods, the study ended up being a cost-minimization study comparing annual costs between periods.

Out of pocket expenditure for EmERGE Participants

Information was also collected from EmERGE participants' socio-economic background, time off work for clinic appointments, return travel time and costs for clinic appointments.

Results:

546 participants were followed up between July 2016-October 2019, of whom 21% (n=113) were seen in the Virtual Clinic pre- EmERGE; 92% of all participants were men and mean age at study entry was 43.3 years (95%CI 42.0-44.6). Median age was 43.0 years (IQR 36.9-49.4) and age at time of baseline ranged from 23.7 to 70.1 years.

Of those participants who self-identified their ethnicity, 77% were Caucasian, 20% Hispanic, 1% Black and 3% 'other'. 76% described themselves as 'men who have sex with men', 14% heterosexuals and two percent as bisexual.

Of participants with known employment status, 84% had fulltime employment of a median 40 hours working-week (IQR 34-40) and a median monthly income of €1450 (IQR €1050-€2050). Five percent received additional social services benefits; 52% received income support, 49% housing benefits and 24% received pension credits. Median monthly additional income was €465 (IQR €400-€738).

Median number of sick days three month prior to enrolment was 0 days (IQR 0-0). Around half of the participants took a day off for professional consultations, blood tests or pick up medication. The median return travel time to the HC- IDIBAPS was 1.5 hours (IQR: 0.5-2.) and median cost was €3 (IQR: €0-8).

Estimated Unit Costs

The unit costs estimated through the micro-costing exercise, are displayed in Table 3 [22]: the unit cost for a virtual clinic visit was less than that for face-to-face visits.

Table 3 near here

Annual Use and Cost of Services pre- and post-mHealth

Across all clinics, outpatient visits increased between periods from 5.2 MPPY (95%CI: 5.0-5.4) to 5.6 MPPY (95%CI: 5.4-5.8) an increase of 8% (Table 4).

Table 4 near here

The annual cost PPY of outpatient visits increased by 12% from €696 (95%CI €654-€741) pre-EmERGE to €778 (95%CI €738-€820) post-EmERGE. The mean number of tests decreased between the two periods (Table 4), as did average costs by 29% (Table 5). The mean number of anal canal procedures increased resulting in a 33% increase of costs for

these procedures, while costs for dexascans decreased and that for fibroscans remained the same between periods (Table 4).

The average annual costs of outpatient visits, tests and procedures increased by 8% across periods from €1,073 (95%CI €999- €1,157) to €1,158 (95%CI €1,084-€1,238) PPY. The average annual cost of ARVs was €7,557 and total average annual costs increased by 1% from €8,430 (95%CI €8,356– 8,514) to €8,515 (95%CI €8,441– 8,595) PPY (Table 4).

When comparing the 433 participants only attended F2F clinics, with those who started with the virtual service and then transferred to a face-to-face (V2F), the annual use and cost for participants that only used F2F services decreased by 5% from €958 (95%CI 905 -1018) to €904 (95%CI 863-945; Table 5). Conversely, post-EmERGE 113 V2F participants were transferred to the F2F outpatient services (Table 3); cost of services for V2F participants increased by a factor of 2.2 from €115 (95%CI 94 -139) pre- to €251 (95%CI 219-290) post-EmERGE.

Table 5 near here

Primary and secondary outcomes pre- and post- mHealth

Median CD4 count at baseline was 759 cells/mm³ (IQR 578-976 cells/mm³) and CD4 counts did not differ significantly between periods. Viral load at baseline was less than 20 copies/ml for all but four participants; the viral load of all participants was undetectable shortly after study entry and remained so during the study (Figure 1).

Figure 1 near here

EmERGE participants' secondary outcome measure PAM-13, measure of patient activation, and their quality of life indicators based on PROQOL-HIV, did not change substantially over time. (Table 6).

Table 6 near here

Discussion

The 8% increase in use and associated costs of HIV outpatient services at HC-IDIBAPS was related to the transfer of EmERGE participants from the virtual ICMID Daycare Office to the two F2F clinics. The participants who only used F2F clinics recorded a 5% reduction in use and cost of services. Similar reductions to those observed in the F2F population were, mutatis mutandis, to those observed in the other EmERGE clinics [32]. The increased cost generated by V2F participants occurred because they were transferred during the second year to the more 'expensive' F2F.

The observed increase in some of the procedures also contributed to the rise in cost. During the first year of EmERGE, anal canal procedures were performed by staff who also had other duties. During the second year, dedicated staff were appointed, which increased the number of referrals for anal scans and the increased number of scans performed, increased expenditure. These changes, however, only minimally affect the conclusion of this analysis, that the implementation of the EmERGE Pathway was an efficient intervention in the HC-IDIBAPS.

Participants, including women and migrants, enjoyed having the EmERGE mHealth App and following the new Care Pathway. The App provided the participants with autonomy, as they became less dependent on going to the clinic for routine consultations. The test results function was the most important for participants [33].

The majority of participants expressed trust in the privacy and security of the EmERGE platform and this was growing over time. Some privacy concerns remained especially among black and migrant women; they were anxious that the App could be seen on the phone by friends or family members, who then would ask questions [33].

Participants remained medically stable and did not have substantive changes in their patient activation and quality of life measures [34]. The before-and-after study provides a robust method for assessing the effect of the intervention and is a favoured method for health-technology assessments [35]. The study is limited by the fact that a control-group was not constituted for the secondary outcome measures – PAM-13 and PROQOL-HIV.

The costs of ARVs used by EmERGE participants was €7,557, within the 2017 price range for ARVs in Spain [36]. ARVs were the main cost-driver of outpatient services for EmERGE participants and a switch to quality-assured and affordable generic ARVs could reduce these costs, although the use of generic drugs can raise other issues [37]. Health care costs can be further reduced when using a single daily pill regimen [38], especially when using generic versions.

Most departments could be micro-costed, apart from the radiology and overhead departments. The HC-IDIBAPS overhead costing system was adopted but the accuracy of the fixed percentages and the extent they covered all overhead costs could not be established [22].

This is the most recent Spanish cost-of-illness study for medically stable PLHIV; no other recent studies could be identified. A study from the Canary Islands reported an annual cost for asymptomatic PLHIV services of €10,351 in 2003 [39]; a second study, reported the cost for asymptomatic PLHIV to be €8,548 at 2010 prices [40].

The expansion of electronic health information systems highlights the need to protect the confidentiality and security of personal health information: at rest - on a phone or server(s) - and in-transit; such protection is of paramount importance [41]. Guidance for protecting personal health information have been published [15] and paper-based and electronic tools developed to investigate the existence and implementation of these guidelines at facility, data warehouse and national levels [42].

The EmERGE study was successfully implemented in five different European countries, which demonstrated its applicability in different cultural settings [32]. It also points to the benefits of including virtual appointments as part of outpatient managements for PLHIV.

Many published systematic reviews high-light the variable effectiveness of mHealth tools. [16]. Some reviews were HIV-specific [3,4,43] whereas others reviewed mHealth tools for other chronic diseases [44,45]. Most studies were performed in high-income countries, but the use of mHealth is increasingly promoted in low- and middle-income countries [46-51]. However, some observations remain relevant today: *“Most of the studies analysed were small scale, short term, pragmatic evaluations that added little to our knowledge of the costs and benefits that would be expected to result from the introduction of telemedicine services into routine clinical practice* [16]. EmERGE involved more participants across different health care systems and countries, but only involved medically stable PLHIV and follow-up was still only one year post-EmERGE implementation.

While efficiencies were achieved at HC- IDIBAPS, broader implementation of the EmERGE Pathway to all PLHIV are likely to lead to greater efficiencies; these savings can be used to address other needs within the clinic. Extending the use of the EmERGE Pathway should be systematically and closely monitored and implementation evaluated and costed [3,4,18,23]. As this study focused on medically stable PLHIV, the services for managing PLHIV with more complex HIV disease or comorbidities need to be costed. Funding should be sought from relevant agencies to monitor and evaluate changes in service provision.

The use of mHealth tools has expanded during the Covid19 Pandemic for both acute and chronic diseases [52], including cancer and mental health services [7,53]. This expansion is expected to continue; in 2019, the global mHealth market was valued at US\$46 billion and is expected to reach \$230 billion by 2027 [54]. The expansion of these tools should make tracking the use, cost, outcomes and impact of health services across facilities easier. Linking individual health information over time and place is part of the aim of *Universal Health Coverage* [55].

References

- 1) Blanco JL, Ambrosioni J, Garcia F, Martínez E, Soriano A, Mallolas J et al on behalf the COVID-19 in HIV Investigators. COVID-19 in patients with HIV: clinical case series. *Lancet HIV* April 15, 2020
[https://www.thelancet.com/journals/lanhiv/article/PIIS2352-3018\(20\)30111-9/fulltext](https://www.thelancet.com/journals/lanhiv/article/PIIS2352-3018(20)30111-9/fulltext)
(Access checked 10-11-2020)
- 2) Beck EJ, Shields JM, Tanna G, Henning G, de Vega I, Andrews G et al. Developing and Implementing National Health Identifiers in Resource Limited Countries: why, what, who, when and how? *Global Health Action*, 2018, volume 11 – issue 1
<https://www.tandfonline.com/doi/full/10.1080/16549716.2018.1440782>
(Access checked 10-11-2020)
- 3) Taylor D, Lunny C, Loli P, Warje O, Geldman J, Wong T, et al. Effectiveness of text messaging interventions on prevention, detection, treatment, and knowledge outcomes for sexually transmitted infections (STIs)/HIV: a systematic review and meta-analysis *Systematic Reviews* (2019) 8:12
<https://doi.org/10.1186/s13643-018-0921-4>
(Access checked 10-11-2020)
- 4) Cooper V, Clatworthy J, Whetham J and EmERGE Consortium. *mHealth Interventions to Support Self-Management In HIV: A Systematic Review. The Open AIDS Journal*, 2017, 11, 119-132
<https://openaidsjournal.com/VOLUME/11/PAGE/119/> (Access checked 10-11-2020)
- 5) Hyle EP, Naidoo K, Su AE, El-Sadr WM, Freedberg KA. HIV, Tuberculosis, and Non-Communicable Diseases: What is known about the costs, effects, and cost-effectiveness of integrated care? *J Acquir Immune Defic Syndr*. 2014; 67(0 1): S87–S95.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4147396/pdf/nihms604877.pdf>
(Access checked 10-11-2020)
- 6) IAM Newswire, *TeleMedicine – one of the first COVID-19 By-Products*, May 19, 2020
https://iamnewswire.com/2020/05/19/telemedicine-one-of-the-first-covid-19-by-products/?mod=article_inline
(Access checked 19-8-2020)

- 7) Cheney C. How to use telehealth in Cancer Care during the Coronavirus Pandemic and beyond. *HealthLeaders*, May 19, 2020

<https://www.healthleadersmedia.com/clinical-care/how-use-telehealth-cancer-care-during-coronavirus-pandemic-and-beyond>

(Access checked 10-11-2020)

- 8) World Health Organization. *Guideline on when to start antiretroviral therapy and on pre-exposure prophylaxis for HIV*. Geneva, World Health Organization 2015

http://apps.who.int/iris/bitstream/10665/186275/1/9789241509565_eng.pdf?ua=1

(Access checked 10-11-2020)

- 9) Marcus JL, Chao CR, Leyden WA, Xu LF, Quesenberry C, Klein D. et al. Narrowing the Gap in Life Expectancy Between HIV-Infected and HIV-Uninfected Individuals With Access to Care. *J Acquir Immune Defic Syndr*. 2016;73(1):39-46.

<https://insights.ovid.com/jaids-acquired-immune-deficiency-syndromes/jaids/2016/09/010/narrowing-gap-life-expectancy-hiv-infected/6/00126334>

(Access checked 10-11-2020)

- 10) Autenrieth CS, Beck EJ, Mallouris C, Stelzle D, Mahy M, Ghys P *Global and regional trends of people living with HIV aged 50 and over: estimates and projections for 2000–2020* PLOS ONE 2018, 13(11): e0207005.

<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0207005>

(Access checked 10-11-2020)

- 11) Althoff KN, Smit M, Reiss P, Justice AC. HIV and ageing: improving quantity and quality of life *Curr Opin HIV AIDS* 2016, 11:527–536

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5084838/>

(Access checked 10-11-2020)

- 12) Shah ASV, Stelzle D, Lee KK, Beck EJ, Alam S, Clifford S et al. Global Burden of Atherosclerotic Cardiovascular Disease in People Living With HIV: Systematic Review and Meta-Analysis. *Circulation*. 2018; 138:1100–1112

<https://www.ahajournals.org/doi/abs/10.1161/CIRCULATIONAHA.117.033369>

(Access checked 10-11-2020)

- 13) Chausa P, Gárate FJ, Cáceres C, Wallitt E, Whetham J and Gómez EJ and the EmERGE Consortium. EmERGE Platform: A New mHealth Solution for People Living with HIV. In:

https://link.springer.com/chapter/10.1007%2F978-3-030-31635-8_99

(Access checked 10-11-2020)

- 14) Waldrop G, Doherty M, Vitoria M and Ford N. Stable patients and patients with advanced disease: consensus definitions to support sustained scale up of antiretroviral therapy. *Trop Med Int Health*, 2016. 21(9): p. 1124-30.

<https://onlinelibrary.wiley.com/doi/full/10.1111/tmi.12746>

(Access checked 10-11-2020)

- 15) Beck EJ, Gill W and De Lay PR. Protecting the Confidentiality and Security of Personal Health Information in Low- and Middle-Income Countries in the Era of the SDGs and Big-Data, *Global Health Action*, 2016.

<https://www.tandfonline.com/doi/full/10.3402/gha.v9.32089>

(Access checked 10-11-2020)

- 16) Whitten PS, Mair FS, Haycox A, May CR, Williams TL, Hellmich S. Systematic review of cost effectiveness studies of telemedicine interventions *BMJ* 2002;324:1434–7

<https://www.bmj.com/content/324/7351/1434>

(Access checked 10-11-2020)

- 17) Hailey D. The need for cost-effectiveness studies in telemedicine. *Journal of Telemedicine and Telecare*, December 1, 2005

<https://doi.org/10.1177/1357633X0501100801>

(Access checked 10-11-2020)

- 18) de Jongh T, Gurol-Urganci I, Vodopivec-Jamsek V, Car J, Atun R. Mobile phone messaging for facilitating self-management of long-term illnesses. *Cochrane Database of Systematic Reviews* 2012, Issue 12. Art. No.: CD007459.

<https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD007459.pub2/abstract>

(Access checked 10-11-2020)

- 19) León A, Cáceres C, Fernández E, Chausa P, Martín M, Codina C, A et al. New Multidisciplinary Home Care Telemedicine System to Monitor Stable Chronic Human

(Access checked 10-11-2020)

- 20) UNAIDS. *Manual for costing HIV facilities and services*. Geneva, Switzerland, 23rd May 2011.

http://www.unaids.org/en/media/unaids/contentassets/documents/document/2011/20110523_manual_costing_HIV_facilities_en.pdf

(Access checked 10-11-2020)

- 21) UNAIDS. *Workbook for the collection of cost information on HIV facilities and services*. Geneva, Switzerland, 23rd May 2011.

http://www.unaids.org/en/media/unaids/contentassets/documents/document/2011/20110523_workbook_cost_info_en.pdf

(Access checked 10-11-2020)

- 22) Beck EJ, Yfantopoulos P, Mandalia S. EU Deliverable 3.3 Costs and Unit Costs for EmERGE clinical sites, 29th April 2020. EU Grant Agreement No: 643736

- 23) Beck EJ, Mandalia S, Yfantopoulos P. EU Deliverable 3.4 *Cost-efficiency of the mHealth Pathway at the Five EmERGE Sites*. 29th April 2020. EU Grant Agreement No: 643736

- 24) West BT, Welch KB, Galecki AT. *Linear Mixed Models. A Practical Guide Using Statistical Software*. Chapman & Hall/CRC, Boca Raton, FL, 2007

- 25) SAS Version 9.4. Cary, NC: SAS Institute Inc; 2014

- 26) Beck EJ, Mandalia S, Gaudreault M, Brewer C, Zowall H, Gilmore N et al. The Cost-effectiveness of HAART, Canada 1991-2001. *AIDS*, 2004; 18: 2411-9.

<https://journals.lww.com/aidsonline/pages/articleviewer.aspx?year=2004&issue=12030&article=00008&type=Fulltext>

(Access checked 10-11-2020)

- 27) Badri M, Maartens G, Mandalia S, Bekker L-G, Penrod JR, Platt RW, Wood R, Beck EJ. Cost-effectiveness of Highly Active Antiretroviral Therapy in South Africa. *Plos Medicine* January 2006; 3: e4

<http://www.plosmedicine.org/article/info:doi/10.1371/journal.pmed.0030004>

(Access checked 10-11-2020)

28) Mandalia S, Mandalia R, Lo G, Chadborn T, Sharott P, Youle M, et al. Rising Population Cost for Treating People Living with HIV in the UK, 1997-2013, *PLoS ONE* 2010 5(12): e15677.

<http://www.plosone.org/article/info:doi/10.1371/journal.pone.0015677>

(Access checked 10-11-2020)

29) Hibbard JH, Mahoney ER, Stockard J, Tusler M (2005) Development and testing of a short form of the patient activation measure. *Health Serv Res* 40: 1918–1930.

<https://europepmc.org/article/PMC/1361231>

(Access checked 10-11-2020)

30) Duracinsky M, Herrmann SE, Berzins B, Armstrong AR, Kohli R, Le Coeur S et al.. The Development of PROQOL-HIV: An International Instrument to Assess the Health-Related Quality of Life of Persons Living With HIV/AIDS. *J Acquir Immune Defic Syndr* 2012; 59:498–505.

https://www.researchgate.net/publication/51977525_The_development_of_PROQOL-HIV_An_international_instrument_to_assess_the_health-related_quality_of_life_of_persons_living_with_HIVAIDS

(Access checked 10-11-2020)

31) Wikipedia. *Incremental cost-effectiveness ratio*

https://en.wikipedia.org/wiki/Incremental_cost-effectiveness_ratio

(Access checked 10-11-2020)

32) Beck EJ, Mandalia S, Yfantopoulos P, Lean A, Merino MJ, Garcia F et al on behalf of the EmERGE Consortium *Ensuring Continuity of Care for People living with HIV in Five European Countries: the efficacy of the EmERGE Platform P117*, HIV Glasgow 2020, Virtual Meeting 5-8 October 2020, Glasgow, Scotland

33) Marent B, Henwood F, Darking M EU Deliverable 2.3 Final Evaluation Report: Best practice in co-design and user (clinician and patient) involvement across study sites. 28th April 2020. EU Grant Agreement No: 643736

34) Jones CI, Bremner S, Whetham J. EU D6.2 *Report on clinical outcomes; patient empowerment and quality of life analyses; patient experience & usability*. April 2020. EU Grant Agreement No: 643736

35) Public Health England: *Guidance, Before-and-after study: comparative studies*. Gov.UK, 30 January 2020

<https://www.gov.uk/guidance/before-and-after-study-comparative-studies>

(Access checked 10-11-2020)

36) Rivero A, Pérez-Molina JA, Blasco AJ, Arribas JR, Asensio V, Crespo M et al. Costs and cost-efficacy analysis of the 2017 GESIDA/Spanish National AIDS Plan recommended guidelines for initial antiretroviral therapy in HIV-infected adults. *Enferm Infecc Microbiol Clin*. 2018;36(5):268–276

<https://doi.org/10.1016/j.eimc.2017.04.002>

(Access checked 10-11-2020)

37) Beck EJ, Mandalia S, Dongmo Nguimfack B, Pinhero E, 't Hoen E, Boulet P et al Does the Political Will exist to bring Quality-Assured and Affordable Drugs to Low- and Middle-Income Countries? *Global Health Action* 2019, 12:1, 1586317

<https://doi.org/10.1080/16549716.2019.1586317>

38) Beck EJ, Mandalia S, Sangha R, Youle M, Brettell R, Gompels M et al. Lower Healthcare Costs Associated with the Use of a Single-Pill ARV Regimen in the UK, 2004-2008. *PLoS ONE*, 2012,7(10): e47376.

<http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0047376>

(Access checked 10-11-2020)

39) Lopez-Bastida J, Oliva-Moreno J, Perestelo-Perez L & Serrano-Aguilar P, The economic costs and health-related quality of life of people with HIV/AIDS in the Canary Islands, Spain. *BMC Health Services Research* 2009, 9:55 doi:10.1186/1472-6963-9-55

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2670289/>

(Access checked 10-11-2020)

40) Trapero-Bertran, M, Oliva-Moreno, J. Economic impact of HIV/AIDS: a systematic review in five European countries. *Health Econ Rev* 4, 15 (2014).

<https://doi.org/10.1186/s13561-014-0015-5>

(Access checked 10-11-2020)

41) Hall JL and McGraw D. For Telehealth to Succeed, Privacy and Security Risks Must Be Identified and Addressed, *Health Affairs* 33, NO. 2 (2014): 216–221

<https://www.healthaffairs.org/doi/10.1377/hlthaff.2013.0997>

(Access checked 10-11-2020)

42) UNAIDS: The Privacy, Confidentiality and Security Assessment Tool: Protecting personal health information. Geneva 14th February 2019.

http://www.unaids.org/en/resources/documents/2019/confidentiality_security_assessment_tool

(Access checked 10-11-2020)

43) Catalani C, Philbrick W, Fraser H, Mechael P and Israelski DM mHealth for HIV Treatment & Prevention: A Systematic Review of the Literature, *The Open AIDS Journal*, 2013, 7, 17-41

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3795408/pdf/TOAIDJ-7-17.pdf>

(Access checked 10-11-2020)

44) de la Torre-Díez I, Lo´pez-Coronado M, Vaca C, Saez Aguado J and Carlos de Castro C, Cost-Utility and Cost-Effectiveness Studies of Telemedicine, Electronic, and Mobile Health Systems in the Literature: A Systematic Review *Telemedicine and e-Health*, 2015, 21 (2); 81-5

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4312789/>

(Access checked 10-11-2020)

45) Sarabi RE, Sadoughi F, Orak RJ and Bahaadinbeigy K. *The Effectiveness of Mobile Phone Text Messaging in Improving Medication Adherence for Patients with Chronic Diseases: A Systematic Review*. *Iran Red Crescent Med J*. 2016 May; 18(5): e25183.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4939231/pdf/ircmj-18-05-25183.pdf>

(Access checked 10-11-2020)

46) Kaplan WA. Can the ubiquitous power of mobile phones be used to improve health outcomes in developing countries? *Globalization and Health* 2006, 2:9 doi:10.1186/1744-8603-2-9

<https://doi.org/10.1186/1744-8603-2-9>

(Access checked 10-11-2020)

47) Govindasamy D, Meghij J, Negussi EK, Baggaley RC, Ford N and Kranzer K. Interventions to improve or facilitate linkage to or retention in pre-ART (HIV) care and initiation of ART in low- and middle-income settings _ a systematic review. *JIAS*, August 2014

<http://dx.doi.org/10.7448/IAS.17.1.19032>

(Access checked 10-11-2020)

48) Thirumurthy H & Lester RT. M-health for health behaviour change in resource-limited settings: applications to HIV care and beyond. *Bull World Health Organ* 2012; 90:390–392

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3341690/pdf/BLT.11.099317.pdf>

(Access checked 10-11-2020)

49) Iribarren SJ, Cato K, Falzon L, Stone PW. What is the economic evidence for mHealth? A systematic review of economic evaluations of mHealth solutions. *PLoS ONE* 2017, 12(2): e0170581.

<https://journals.plos.org/plosone/article/file?id=10.1371/journal.pone.0170581&type=printable>

(Access checked 10-11-2020)

- 50) Rout SK, Gabhale YR, Dutta A, Balakrishnan S, Lala MM, Setia MS, et al. Can telemedicine initiative be an effective intervention strategy for improving treatment compliance for pediatric HIV patients: Evidences on costs and improvement in treatment compliance from Maharashtra, India. *PLoS ONE*, 2019, 14(10): e0223303.

<https://journals.plos.org/plosone/article?id=10.1371%2Fjournal.pone.0223303>

(Access checked 10-11-2020)

- 51) Demena BA, Artavia-Mora L, Ouedraogo D, Thiombiano BA and Wagner N. A Systematic Review of Mobile Phone Interventions (SMS/IVR/Calls) to Improve Adherence and Retention to Antiretroviral Treatment in Low and Middle-Income Countries. *AIDS Patient Care and STDs*, 2020,34, (2)

https://www.liebertpub.com/doi/10.1089/apc.2019.0181?url_ver=Z39.88-2003&rfr_id=ori:rid:crossref.org&rfr_dat=cr_pub%20%200pubmed

(Access checked 10-11-2020)

- 52) Wright JH and Caudill R Remote Treatment Delivery in Response to the COVID-19 Pandemic *Psychother Psychosom* 2020; 89:130–132,

<https://www.karger.com/Article/FullText/507376> (Access checked 10-11-2020)

- 53) Humer E, Pieh C, Kuska M, Barke A, Doering BK, Gossmann K et al Provision of Psychotherapy during the COVID-19 Pandemic among Czech, German and Slovak Psychotherapists, *Int. J. Environ. Res. Public Health* **2020**, 17(13), 4811.

<https://www.mdpi.com/1660-4601/17/13/4811> (Access checked 10-11-2020)

- 54) Allied Market Research. *COVID-19 Impact on mHealth Market Size to hit \$230 billion by 2027, Top Prominent Players: Bayer Healthcare., Boston Scientific Corporation*. Press Release 27-7-2020

<https://www.openpr.com/news/2098085/covid-19-impact-on-mhealth-market-size-to-hit-230-billion>

(Access checked 10-11-2020)

- 55) WHO / International Bank for Reconstruction and Development / The World Bank. *Tracking universal health coverage: 2017 global monitoring report*, 2017

<http://pubdocs.worldbank.org/en/193371513169798347/2017-global-monitoring-report.pdf>

(Access checked 10-11-2020)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1 CD4 count and Viral Load Changes during pre- and post-EmERGE periods at HC- IDIBAPS

Table 1. Inclusion and Exclusion criteria for EmERGE Participants [14]*

<i>Inclusion criteria:</i> Patients who meet all of the following criteria were eligible for this study:	• Documented HIV infection
	• Aged at least 18 years old
	• Able to give informed consent
	• In possession of a smartphone, tablet, or similar technology supporting the mHealth platform
	• Clinically stable on anti-retroviral therapy (ART). This was defined as receiving ART for at least 1 year and unchanged regimen for at least 3 months, 2 consecutive undetectable viral load measures (<50 copies/ml), no current pregnancy and without any new WHO clinical stage 2, 3 or 4 events within the previous 12 months*.
<i>Exclusion criteria:</i> Patients who met one or more of the following criteria were excluded from the study	• Aged less than 18 years
	• Pregnant
	• Participating in a clinical trial or receiving an investigational medication
	• Unable to comprehend the patient information sheet
	• Unable to comprehend the instructions for using the mHealth platform
	• Considered for any other reason by their regular physician to be unsuitable for study participation

Table 2 Eleven Requirements of a Tele-Medicine System [4]*, [15]**

1. Patient-provider and peer communication*;
2. Medication and appointment reminders*;
3. A medication checklist, pill identification function and list of current and discontinued medicines*;
4. Laboratory reports (CD4 count, viral load, sexually transmitted infections, glucose and complete blood count)*;
5. Pharmacy information*;
6. Nutrition and fitness trackers;
7. Resources, links to social services, substance abuse support, video testimonials, case-management*;
8. Settings (profile picture, password and alerts)*;
9. A search function*.
10. Protecting the confidentiality and security of personal information at rest and in-transit**
11. Affordability and efficiency of the technology**

Table 3 Unit Costs of HIV Outpatient Services for HC- IDIBAPS EmERGE Participants	
HC- IDIBAPS	Cost / Unit cost
Unit cost per EmERGE Participant Dispensario clinic Outpatient visit	€173
Unit cost per EmERGE Participant ICMiD Office Virtual Outpatient visit	€51
Unit cost per EmERGE Participant Day-care Hospital Outpatient visit	€118
Unit cost per EmERGE Participant Hospital Outpatient visits across all Clinics	€114
Unit cost per EmERGE Participant ICMiD Office Anal Canal Day ward Procedure	€398
Unit cost per EmERGE Participant Day-care Hospital Fibroscan Day ward Procedure	€64
Unit cost per EmERGE Participant Day-care Hospital Dexascan Day ward Procedure	€64
Unit cost per EmERGE Participant Pharmacy patient excluding ARVs costs	€45
Annual cost of ARVs for EmERGE Participant	€7,357
Unit cost per EmERGE Participant Biochemistry Laboratory test	€0.7
Unit cost per EmERGE Participant Haematology Laboratory test	€2.0
Unit cost per EmERGE Participant Microbiology Laboratory test	€13
Unit cost per EmERGE Participant Immunology Laboratory test	€20
Unit cost per EmERGE Participant Radiology investigation	€39

Table 4 Annual Mean Use and Cost PPY HIV Outpatient Visits, Tests and Procedures, Pre- and Post-EmERGE implementation at the HC- IDIBAPS.				
N=546	Pre-EmERGE		Post-EmERGE	
	Mean/average	95% CI interval	Mean/average	95% CI interval
Mean Dispensario Outpatient visits PPY and average costs PPY	2.1	2.0 to 2.3	2.3	2.2 to 2.5
	€ 368	€ 346 to € 391	€ 401	€ 379 to € 424
Mean ICMiD Office virtual visit and average costs PPY	0.6	0.5 to 0.6	0.05	0.04 to 0.08
	€ 28	€ 24 to € 32	€ 3	€ 2 to € 4
Mean Daycare Outpatient visits and average costs PPY	2.6	2.4 to 2.7	3.2	3.0 to 3.3
	€ 300	€ 284 to € 318	€ 374	€ 357 to € 392
Outpatient visit MPPY	5.2	5.0 to 5.4	5.6	5.4 to 5.8
TOTAL Average Cost visits PPY	€ 696	€ 654 to € 741	€ 778	€ 738 to € 820
Mean Biochemistry tests and average costs PPY	51.0	50.4 to 51.7	36.7	36.1 to 37.2
	€ 36	€ 35 to € 36	€ 26	€ 25 to € 26
Mean Haematology tests and average costs PPY	11.3	11.0 to 11.6	8.4	8.1 to 8.6
	€ 23	€ 22 to € 23	€ 17	€ 16 to € 17
Mean Microbiology tests and average costs PPY	3.4	3.3 to 3.6	2.4	2.2 to 2.5
	€ 44	€ 42 to € 47	€ 31	€ 29 to € 32
Mean Immunology tests and average costs PPY	4.6	4.4 to 4.8	3.2	3.0 to 3.4
	€ 92	€ 88 to € 96	€ 64	€ 61 to € 67
Mean tests PPY	70.3	69.2 to 71.7	50.7	49.4 to 51.7
Total average tests PPY	€ 195	€ 187 to € 202	€ 138	€ 131 to € 142
Mean Anal Canal Procedures and average costs PPY	0.39	0.34 to 0.45	0.56	0.50 to 0.63
	€ 155	€ 135 to € 179	€ 223	€ 199 to € 251
Mean Day care Hospital Dexascans and average costs PPY	0.21	0.17 to 0.26	0.08	0.06 to 0.11
	€ 13	€ 11 to € 17	€ 5	€ 4 to € 7
Mean Day care Hospital Fibroscans and average costs PPY	0.05	0.03 to 0.08	0.05	0.03 to 0.07
	€ 3	€ 2 to € 5	€ 3	€ 2 to € 5
Mean Imaging Procedures and average costs PPY	0.28 (95%CI 0.24 - 0.33) - € 11 (95%CI € 10 - € 13)			
Mean procedures per PPY	0.93	0.78 to 1.12	0.97	0.83 to 1,14
Total annual cost procedures	€182	€158 to €214	€242	€215 to €276
Total annual cost of outpatient services	€1,073	€999- €1,157	€1,158	€1,084-€1,238
Annual ARV cost	€7,557			

Total annual outpatient costs including ARVs	€8,430	€8,356 – 8,514	€8,515	€8,441 – 8,595
---	---------------	-----------------------	---------------	-----------------------

Table 5 Annual Mean Use and Cost PPY HIV Outpatient Visits, Tests and Procedures, Pre- and Post-EmERGE implementation at the HC- IDIBAPS for participants used face-to-face outpatients only (F2F) and participants who use virtual and face-to-face clinics (V2F).

F2F participants Average annual cost clinic visits	€668	95%CI €630-709	€594	95%CI €573-614
F2F participants average costs tests & procedures	€290	95%CI €275 - 309	€310	95%CI €290 – 331
F2F participants total average annual outpatient cost	€958	95%CI €905 -1018	€904	95%CI €863 -945
V2F participants average annual outpatients visit costs	€ 28	95%CI €24- 32	€181	95%CI €163-203
V2F participants average costs tests & Procedures	€87	95%CI €70-107	€70	95%CI €56-87
Total Average annual cost V2F participants	€115	95%CI €94-139	€251	95%CI €219-290
Total annual cost of outpatient services	€1,073	95%CI €999- €1,157	€1,158	95%CI €1,084-€1,238

Table 6. Median and IQR for PAM13and PROQOL-HIV Months 0 and 12, baseline - and post-EmERGE at the HC-IDIBAPS						
PAM 13	Month 0- recruitment baseline			Month 12 Post- EmERGE		
	Median	IQR		Median	IQR	
	70.2	(58.1-80.9)		67.8	(58.1-77.7)	
PROQOL-HIV	Month 0 (Baseline)			Month 12 (Post- EmERGE)		
	n	Median	IQR	n	Median	IQR
Physical Health and Symptoms	535	88.9	77.8 to 97.2	281	83.3	69.4 to 94.4
Body Change	535	93.8	81.3 to 100.0	280	87.5	68.8 to 100.0
Social Relationships	540	100.0	100.0 to 100.0	283	100.0	100.0 to 100.0
Intimate Relationships	534	91.7	66.7 to 100.0	282	83.3	50.0 to 100.0
Stigma	540	50.0	25.0 to 87.5	282	50.0	25.0 to 87.5
Emotional Distress	537	93.8	75.0 to 100.0	280	87.5	68.8 to 100.0
Health Concerns	537	81.3	62.5 to 100.0	279	81.3	56.3 to 93.8
Treatment Impact	533	87.5	80.0 to 95.0	277	85.0	77.5 to 92.5