First do no harm: reconsidering our approach to weight in primary care

Article (Accepted Version)

Shaw, Sebastian C K and Meadows, Angela (2022) First do no harm: reconsidering our approach to weight in primary care. *British Journal of General Practice*, 72 (716). pp. 102-103. ISSN 0960-1643

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/104170/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

First do no harm: re-considering our approach to weight in primary care

¹Sebastian C. K. Shaw & ²Angela Meadows

¹Honorary Clinical Lecturer, Department of Medical Education, Brighton and Sussex Medical School, Brighton, UK; ²Lecturer, Department of Psychology, University of Essex, Colchester, UK.

SCK Shaw (corresponding author)

Email: S.Shaw2@bsms.ac.uk

A Meadows

Email: drameadows@gmail.com

Word count: 940

Acknowledgements: None.

Conflict of interest: We declare no competing interests.

Weight stigma may be defined as “the social devaluation, denigration, and marginalization of” higher-weight individuals (1). Weight stigma is well documented in the healthcare context, including in general practice, with implicit and explicit anti-fat biases consistently found in primary care physicians and serious implications for quality of care and patient outcomes (2). Weight stigma is not only directed at patients. As a higher-weight doctor in the UK, SS has experienced numerous incidents of being publicly shamed or pointed to as a cautionary tale by colleagues in both primary and secondary care settings. Such experiences are validated by the recent findings from a North American study, in which 87% of doctors were found to harbour implicit anti-fat bias where higher-weight colleagues were concerned (3). Interestingly, when asked explicitly, most participants reported low levels of bias toward higher-weight doctors, suggesting that many may be unaware of their biases.

The situation seems unlikely to improve, as weight stigma is not only entrenched in our everyday lives, but also further fostered during medical education. Healthcare education and discourses surrounding obesity promote a view of fatter as lesser and often seem to reduce the complexity of body weight to a

simplistic, individual-blaming rhetoric (4). Additionally, it is not unusual for medical students to observe denigration and discriminatory treatment of higher-weight patients by faculty and staff, contributing to the (not so) “hidden curriculum” that promotes implicit, and in some cases explicit, anti-fat bias in our medical workforce (5). In one UK study, exploring beliefs in negative stereotypes about obese people, only 2.1% of medical students expressed neutral or better attitudes towards this group (6).

Weight stigmatisation should have no place in a profession whose ethos is to first do no harm. Experiences of weight stigma are associated with physical and psychological morbidity and mortality, independent of body mass index (BMI) (1). In young people, experienced stigma is associated with increased suicidality – even labelling young people as “overweight” is associated with disordered eating, unhealthy weight-control behaviours, and long-term weight gain, independent of baseline BMI (7).

Unsurprisingly, weight stigma among healthcare providers is linked to healthcare avoidance in patients, including under-utilisation of preventive healthcare and screening (2). And lest one should believe that stigma serves the patients’ best interests, motivating them to “do something” about their weight, evidence consistently demonstrates that stigma is more likely to lead to increased allostatic load, higher prevalence of chronic diseases, and further weight gain, again, independent of BMI (1). Even encouraging higher-weight patients to view their weight as a problem could have paradoxical effects. Evidence suggests that chronic weight dissatisfaction over time predicts increased risk of developing type 2 diabetes mellitus, even when controlling for BMI – the effect being greater than having a family history of diabetes (8).

Our training and praxis seem to encourage a culture of ‘othering’ of higher-weight individuals, that seems to give us licence to treat higher-weight individuals as a problem or a burden. However, the foundations on which our attitudes towards higher-weight individuals apparently rest may not be as solid as generally thought. Research has demonstrated that high-weight status may not be the strong determinant of health it is generally considered to be, and that weight loss is not required to improve metabolic health (1,7). It is also unlikely to happen. Diets are ineffective in the long-term and frequently produce paradoxical results, worsening health and driving further weight gain (9).

While body weight naturally varies across a continuum, as with other physical characteristics, the intersectional nature of obesity prevalence speaks to the social determinants of health (10) – an issue that cannot be solved by individual behaviour change. That many clinical guidelines continue to treat obesity as a disease, the solution to which is located at the level of the individual patient, can

be at least in part attributed to powerful vested interests and a narrow focus on an evidence base that is limited in its scope and that takes pathology as its starting point and continues to publish in an academic echo chamber (11,12). This situation is not dissimilar to the field of autism research, which has been revolutionised in the last 20 years through the introduction of the neurodiversity movement, reframing research discourses away from the pathologizing lens of deviance and toward one of difference.

Thus, the dogmatic promotion of individual weight loss is likely to be ineffective, damage patient–physician relationships, promote overmedicalisation, propagate weight stigma, and is at odds with a holistic approach to healthcare and the ethical principles of beneficence and social justice. People have a “right to exist in their bodies without prejudice, stigmatization, marginalization, or oppression” (1). As advocates for our patients, including their holistic wellbeing and autonomy, our training may place us in a difficult position. As physicians, our aim is to centre the needs of our patients, yet we carry a deeply ingrained instinct to shield the healthcare system from the long-term economic burden that we have been taught obesity represents. Yet this role conflict arises from false foundations.

We call for a better understanding and approach to education of the issues and discourse surrounding weight and health. We need to build a healthcare system that acknowledges and aims to eliminate the impact of social inequality on health outcomes, and to recognise that weight stigma perpetuates and magnifies health disparities in already marginalised populations. While diversity-related legislation does not protect higher-weight individuals from discrimination, we as healthcare professionals have an ethically bound duty to do so. In primary care, we are the gatekeepers to specialist services and are typically patients’ first port of call for medical concerns. Let us lead the way in challenging tradition and fostering positive change.

References:

1. Calogero RM, Tylka TL, Mensinger JL, Meadows A, Danielsdottir S. Recognising the fundamental right to be fat: a weight-inclusive approach to size acceptance and healing from sizeism. *Women & Therapy* 2019;**42(1-2)**:22–44. DOI: 10.1080/02703149.2018.1524067
2. Phelan SM, Burgess DJ, Yeazel MW, Hellerstedt WL, Griffin JM, Van Ryn M. Impact of weight bias and stigma on quality of care and outcomes for patients with obesity. *Obesity Reviews* 2015;**16(4)**:319–326. DOI: 10.1111/obr.12266

3. McLean ME, McLean LE, McLean-Holden AC, Campbell LF, Horner AM, Kulkarni ML, et al. Interphysician weight bias: A cross-sectional observational survey study to guide implicit weight bias training in the medical workplace. *Academic Emergency Medicine* 2021;**28(9)**:1024–1034. DOI: 10.1111/acem.14269
4. Lomax-Sawyers I. On being a fat medical student, at the start of our metabolism module. *Journal of Paediatrics and Child Health* 2017;**53**:733–734. DOI: 10.1111/jpc.13648
5. Meadows A, Barreto M, Dovidio JF, Burke SE, Wittlin NM, Herrin J, van Ryn M, Phelan SM. Signalling hostility: The relationship between witnessing weight-based discrimination in medical school and medical student wellbeing. *Journal of Applied Social Psychology* 2021;In press.
6. Swift JA, Hanlon S, El-Redy L, Puhl RM, Glazebrook C. Weight bias among UK trainee dietitians, doctors, nurses and nutritionists. *Journal of Human Nutrition and Dietetics* 2013;**26**:395–402. DOI: 10.1111/jhn.12019
7. Hunger JM, Smith JP, Tomiyama AJ. An evidence-based rationale for adopting weight-inclusive health policy. *Social Issues and Policy Review* 2020;**14**:73–107.
8. Wirth M, Blake CE, Hébert JR, Sui X, Blair SN. Chronic weight dissatisfaction predicts type 2 diabetes risk: *Aerobic Center Longitudinal Study*. *Health Psychology* 2014;**33(8)**:912–919.
9. Rothblum ED. Slim chance for permanent weight loss. *Archives of Scientific Psychology* 2018;**6**:63–69.
10. Ailshire JA, House JS. The unequal burden of weight gain: an intersectional approach to understanding social disparities in BMI trajectories from 1986 to 2001/2002. *Social Forces* 2011;**90(2)**:397–423.
11. O'Hara L, Thille P, Adams L, Meadows A. Rhetoric versus substance: How new are Canada's 'new' obesity guidelines? 2021;Manuscript submitted for publication.
12. O'Hara L, Taylor J. What's wrong with the 'war on obesity'? A narrative review of the weight-centered health paradigm and development of the 3C framework to build critical competency for a paradigm shift. *SAGE Open* 2018;**8(2)**.

Dr Sebastian Shaw is a General Practice Speciality Trainee in the Southeast of England. His academic interests stem from the social sciences and his PhD focused on medical education. He is an Honorary Clinical Lecturer at Brighton

and Sussex Medical School, where he lectures in research methods and medical education, with interests in diversity, inclusivity, and support.

Dr. Angela Meadows is a Lecturer in Psychology at the University of Essex. She is a social psychologist specializing in prejudice relating to weight and body size. Her work focuses on how higher-weight people respond to stigma in their daily lives, and the implications for individuals and for social change.