

Cloning and characterisation of the rad9 DNA repair gene from *Schizosaccharomyces pombe*

Article (Published Version)

Murray, Johanne M, Carr, Antony M, Lehmann, Alan R and Watts, Felicity Z (1991) Cloning and characterisation of the rad9 DNA repair gene from *Schizosaccharomyces pombe*. *Nucleic Acids Research*, 19 (13). pp. 3525-3531. ISSN 0305-1048

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/26347/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Cloning and characterisation of the *rad9* DNA repair gene from *Schizosaccharomyces pombe*

Johanne M. Murray, Antony M. Carr¹, Alan R. Lehmann¹ and Felicity Z. Watts*

School of Biological Sciences, University of Sussex, Falmer, Brighton, East Sussex, BN1 9QG and

¹MRC Cell Mutation Unit, University of Sussex, Falmer, Brighton, East Sussex, BN1 9RR, UK

Received April 23, 1991; Revised and Accepted June 7, 1991

EMBL accession no. X58231

ABSTRACT

The *rad9.192* DNA repair mutant from the fission yeast, *Schizosaccharomyces pombe*, is sensitive to both UV and ionising radiation. The *rad9* gene has been cloned by complementation of the gamma-ray sensitivity of the mutant cell line. A 4.3kb *HindIII* fragment was found to confer resistance to both types of radiation. The region of complementation was further localised to a 2.6kb *HindIII-EcoRV* fragment, which, by DNA sequence analysis, was found to contain sequences capable of coding for a 427 amino acid protein, if three introns were postulated to remove stop codons. The introns were confirmed by sequence analysis of cDNA clones and PCR products derived from cDNA. The product of transcription is a 1.6kb mRNA of low abundance. The putative *rad9* protein shows no homology to any published sequence. A truncated protein is capable of complementing the radiation sensitivity of the *rad9.192* mutant. Deletion of the gene is not lethal and the null allele has a similar phenotype to the *rad9.192* mutant.

INTRODUCTION

In the fission yeast, *Schizosaccharomyces pombe*, mutants that are associated with defects in DNA repair have been assigned to 23 different complementation groups (1, 2). These *rad* mutants, unlike those identified in *Saccharomyces cerevisiae*, have not been extensively characterised. Most mutants are sensitive to both UV and gamma radiation. In this respect, they differ from the majority of *S. cerevisiae* *RAD* mutants, which are mostly sensitive to either UV- (excision repair) or gamma-irradiation (recombination repair) but not to both.

DNA repair is of fundamental importance in all living organisms. This is illustrated by the existence in man of a number of genetic disorders due to defects in DNA repair. The cloning and characterisation of *S. pombe* DNA repair genes should provide information relevant to DNA repair in other eukaryotes. This may be particularly relevant to the study of DNA repair in humans as the conservation of basic cellular mechanisms between *S. pombe* and higher eukaryotes has been well documented: as demonstrated, for example, in the cloning of a human cell cycle gene by functional complementation of an *S.*

pombe mutant (3). *S. pombe* is also more like higher eukaryotes than *S. cerevisiae* in its prevalence of introns (4) and its ability to splice a mammalian intron correctly (5).

We report here the cloning of the *S. pombe rad9* gene. The *rad9.192* mutant is highly sensitive to both UV and gamma radiation. It also has a greater percentage of cell death during mitotic growth than wild type strains. It has been reported that, unlike *rad*⁺ cells, the survival of *rad9.192* cells after irradiation is not decreased in the presence of caffeine, suggesting that *rad9* acts in a caffeine-sensitive repair pathway (1). It has been suggested that caffeine inhibits intergenic recombination in *S. pombe* (6) and that this implies that *rad9.192* may be defective in a recombination repair pathway. However, spontaneous intragenic recombination in *rad9.192* cells is approximately double that in wild type cells (1), indicating that cells containing the *rad9.192* mutation are still capable of undergoing some types of recombination. The UV-induced forward mutation rate has been reported to be slightly reduced (1).

MATERIALS AND METHODS

Strains and Plasmids

S. pombe strains and plasmids used are shown in tables 1 and 2 respectively. The *S. pombe* gene bank, constructed from partially *HindIII* digested genomic DNA, in the yeast shuttle vector pDB262, containing *S. cerevisiae* *LEU2* and 2 μ sequences has been described previously (7). The *S. pombe* cDNA library was a gift from L. Guarente (8). Rescued plasmids were subcloned into the yeast vectors pSTA12 (7), pUR19 (9) and M13mp19.

Plasmids were grown in the *E. coli* strain DH5 α , genotype: F-*endA1*, *hsdR17*, (*rk*-, *mk*-), *supE44*, *thi-1*, *l*-, *recA1*, *gyrA96*, *relA1*, *lacIqZ-M15(lacproAB)* (10) and rescued from *S. pombe* into the *recBC* strain JA226 (11). M13 derivatives were grown in DH5 α F' (genotype as for DH5 α but containing an integrated F').

Growth Conditions

S. pombe strains were grown and routinely maintained on complete medium (12). Selection for plasmid-containing cells utilised minimal medium or agar plates, as appropriate. Minimal

* To whom correspondence should be addressed

medium agar plus sorbitol was used to plate cells after transformation (11). Minimal medium containing 0.05g/l NH₄Cl (low nitrogen) was used for mating and sporulation. Bacteria were grown in LB or 2YT (13) containing appropriate antibiotics.

Transformation

Yeast transformation was performed according to the spheroplast method of Beach *et al.*, (11). Bacterial transformation was by the calcium chloride method described by Maniatis *et al.* (13).

DNA Extraction and Purification

Plasmid and chromosomal DNA from yeast were purified according to the method of Aves *et al.*, (14). Plasmids from *E. coli* were prepared according to the alkali lysis method of Burke and Ish-Horowitz (13). Large scale preparations of plasmids were purified by caesium chloride-ethidium bromide centrifugation (13). DNA fragments were purified by extraction from low gelling temperature agarose using 'GeneClean' (Stratatech Scientific Ltd.) according to the manufacturer's recommended methods.

RNA Extraction and Purification

Total RNA was extracted from *S. pombe* by vortexing the cells in the presence of glass beads and phenol (15). Poly A⁺ RNA was made by passing total RNA over oligo dT cellulose (13). For Northern analysis, polyA⁺ RNA was run on 1% agarose gels after glyoxal treatment and transferred to Genescreen Plus (New England Nuclear Research Products) and probed by the manufacturer's recommended method.

DNA Sequence Analysis

Nucleotide sequencing was performed using Sequenase (US Biochemical Corp.) on DNA fragments subcloned into M13mp19. For the sequence of both strands of the complementing 4.3kb genomic fragment, nested series of deletions, in both directions, were made using exonuclease III according to the protocol of Henikoff (16). The sequence was analysed using the DNASTAR computer programs.

The sequence of the cDNA clones across the introns was determined in M13mp19 using the universal primer and

oligonucleotides made to specific regions of the genomic sequence as alternative primers. These were:

O.3 (5'-ATGACTCCTACAATACGTATTCGTAC-3'),
O.11 (5'-ATGTTAATCTTCGGGACCTCGCAAG-3'),
O.13 (5'-AGTCTTCCTGAGAGAAAATGCCATGA-3'),
see figure 3.

Polymerase Chain Reaction

DNA sequences were amplified from cDNA made from *S. pombe* polyA⁺ RNA, and from clones isolated from the cDNA library, using the polymerase chain reaction. Oligonucleotide sequences used as primers were as follows:

O.11 5'-ATGTTAATCTTCGGGACCTCGCAAG-3' upstream of intron I,
O.15 5'-GATGAGTCACTTTGATCAGTTTGCCA-3' spanning intron III,
O.17 5'-ATTTTATATGTTTTAATCACTCCGTGC-3' spanning intron II,

cDNA was synthesised with murine Moloney leukemia virus reverse transcriptase (Pharmacia) using a modification of the procedure of Yang *et al.* (17). To prime the reaction 1.5μg polyA⁺ RNA were annealed with 50ng oligo dT by heating to 80°C for 3 min and incubating at 37°C for 5 min. After cDNA synthesis for 1 hr at 37°C, 10% of the cDNA was used directly for PCR analysis.

PCR (18) was carried out for 30 cycles with 1 min at 94°C, 2 min at 58°C and 2.5 min at 70°C. 100ng plasmid DNA was used to amplify sequences from the cDNA clones, 1ng and 100ng genomic DNA used as control.

Pulse Field Gel Electrophoresis and General Molecular Methods

Molecular biology methods were as described in Maniatis *et al.* (13). Preparation of high molecular weight DNA, *NotI* digestion and transagonal pulse field gel electrophoresis of *S. pombe* chromosomal DNA was performed according to the Beckman protocol using the Beckman TAFE system. *NotI* digested *S. pombe* chromosomal DNA was separated in two phases. The first stage consisted of 4 sec pulses at 170mA for 30 min and the second a 60 sec pulse at 150mA for 18 h. The gel was stained with ethidium bromide to visualise the DNA bands. DNA was denatured and transferred to a Hybond-N filter (Amersham, UK).

Irradiation of *S. pombe* Cells

Gamma irradiation was routinely carried out using a ⁶⁰Co gamma source, with a dose rate of 5 Gy/min, on 1 × 10⁷ cells/ml in minimal medium with appropriate supplements. The cells were then plated on minimal medium plates and colonies counted after 3 days at 30°C. Patch tests were carried out on

Table 1. *S. pombe* strains

sp.037 *rad9.192*, *h*⁻: from the Canadian National Cell Repository.
sp.011 *leu1.32*, *ade6.704*, *ura4.D18*, *h*⁻: from Paul Nurse, Oxford
sp.012 *leu1.32*, *ade6.704*, *ura4.D18*, *h*⁺: from Paul Nurse, Oxford
sp.014 *lys2.97*, *h*⁻: from the National Yeast collection, Norwich.

Strains created for this study:

sp.039 *rad9.192*, *leu1.32*, *ade6.704*, *ura4.D18*, *h*⁻.
sp.016 *lys2.97*, *ade6.704*, *leu1.32*, *ura4.D18*, *h*⁻.
sp.078 *leu1.32*, *ade6.704*, *ura4.D18*, *rad9*⁺/*rad9*⁺ + *sup3.5*, *h*⁺.
sp.101 *leu1.32/leu1.32*, *ade6.704/ade6.704*, *ura4.D18/ura4.D18*, *h*⁺/*h*⁺.

Deletion strains created:

sp.086 *leu1.32/leu1.32*, *ade6.704/ade6.704*, *ura4.D18/ura4.D18*, *rad9/rad9::ura4*, *h*⁺/*h*⁺.
sp.102 *leu1.32/leu1.32*, *ade6.704/ade6.704*, *ura4.D18/ura4.D18*, *rad9/rad9::ura4*, *h*⁹⁰/*h*⁺.
sp.069 *leu1.32*, *ade6.704*, *ura4.D18*, *rad9*⁺/*rad9*⁺::*ura4*, *h*⁻.
sp.070 *leu1.32*, *ade6.704*, *ura4.D18*, *rad9*⁺/*rad9*⁺::*ura4*, *h*⁺.
sp.071 *leu1.32*, *ade6.704*, *ura4.D18*, *h*⁻.
sp.072 *leu1.32*, *ade6.704*, *ura4.D18*, *h*⁺.

sp.069-sp.072 are the products of a single tetrad of sp.102.

Table 2. Plasmids

pUR19	pUC19 based <i>ura4 ars6</i> plasmid (9).
pSTA12	pUC12 based <i>sup3.5</i> plasmid (7).
pDB262	<i>LEU2</i> 2μ plasmid (11).
p9/4	4.3kb <i>HindIII rad9.192</i> complementing insert in DB262.
pRAD9	2.6kb <i>HindIII-EcoRV</i> fragment from p9/4 in pUR19.
pS9	4.3kb <i>HindIII</i> fragment from p9/4 in STA12.
pS9U4	pS9 with <i>Clal-MluI</i> replaced with 1.7kb <i>ura4</i> fragment.
pA4.5	A4.5 <i>ExoIII</i> deletion in pUR19.
pS13	S13 <i>ExoIII</i> deletion in pUR19.
pES13	<i>EcoRI</i> -end S13 deletion in pUR19.

6 cm plates that were lowered directly into the chamber of the source for a time calibrated to deliver a 500 Gy dose.

UV irradiation was carried out in a Stratagene 'Stratalinker' directly on freshly plated cells. For the survival curves, four plates were counted for each point and the analysis repeated at least twice.

RESULTS

Complementation of the *rad9* Mutation

An *S. pombe* genomic library containing partially *Hind*III digested DNA in the shuttle vector pDB262 was used to transform the *rad9.192* strain sp.039. 40 000 transformants were selected for their ability to grow in the absence of leucine and the cells were then irradiated with a dose of 500 Gy in a ⁶⁰Co source, to select for plasmids complementing the radiation sensitivity. At this dose the difference in survival between wild type and *rad9.192* cells was maximal. After a 500 Gy dose, approximately 80% of wild type cells, but less than 1% of *rad9.192* cells, can form colonies




Figure 1. Physical maps of the *rad9* region and complementation data. a. Restriction map of the 4.3kb *Hind*III insert of p9/4 and below this the restriction map of the 2.6kb *Hind*III-*Eco*RV subclone, prad9. H = *Hind*III, E = *Eco*RI, C = *Cla*I, M = *Mlu*I, S = *Ssp*I, V = *Eco*RV. The boxed line represents the genomic sequence, the filled box the exons and the open box the untranslated leader, introns and 3' sequence. The region (*Cla*I-*Mlu*I) replaced by a 1.7kb fragment containing the *ura4* gene in the disruption is indicated below the coding region. b. Diagram to show the position of three cDNA clones with respect to the transcribed region, the filled box the exons and the open box the untranslated leader, introns and 3' sequence. Intron positions are indicated and the complementation analysis is given in the right hand column. c. The functional complementation data are drawn with respect to the prad9 map above. The regions present in the *Exo*III deletions S7, S9, S13, S15, A4, A4.5, A5.5 are represented by bars and the complementation analysis is given in the right hand column. The minimum complementing region (end of A4.5-end of S13) is shown.

(see fig 2a). After three rounds of selection, individual colonies were patched out on 6 cm minimal plates and tested for sensitivity to gamma-radiation. Resistant clones were then tested for co-instability of the *leu*⁺ and *rad*⁺ phenotypes. This indicated that the complementation was due to plasmid-borne sequences rather than to mutations in the genome.

DNA was prepared from the resistant colonies for plasmid rescue into the *E. coli* strain JA226. Plasmids were rescued from two independent radiation resistant colonies. These were retransformed into the *rad9.192* strain sp.039 and found to complement the *rad9.192* phenotype. The complementing plasmids were identical, and contained four *Hind*III fragments of 4.3kb, 1.7kb, 1.5kb and 0.4kb. The *rad9* complementing region was found, on routine subcloning, to be contained within the 4.3kb fragment, contained in plasmid p9/4. Further analysis localised the *rad9* gene to a 2.6kb *Hind*III-*Eco*RV fragment, in plasmid prad9 (see fig. 1A).

Mapping of the Chromosomal Locus of p9/4

The 4.3kb *Hind*III insert from p9/4 was subcloned into the non-*ars* plasmid pSTA12 (7) which contains the *sup3-5* suppressor




Figure 2. Survival curves of *S. pombe* strains following gamma or UV irradiation. a. and b. Survival of the *rad*⁺ strain sp.011 and the *rad9.192* strain sp.039 following gamma- and UV-irradiation. c. Survival of strain the *rad9.192* strain sp.039 containing the complementing plasmids p9/4 and prad9 or the control vectors pUR19 and DB262, following gamma-irradiation. d. Survival of strain the *rad9.192* strain sp.039 containing the complementing plasmids p9/4 and prad9 or the control vector pUR19, following UV-irradiation. e. and f. Gamma- and UV-irradiation survival of the four products of a tetrad of the diploid deletion strain sp.102 and a *rad9.192* control, sp.039. The two null mutants are sp.069 and sp.070 and the two with wild type sensitivity to gamma-irradiation are sp.071 and sp.072.

eliminates the possibility of contaminating DNA obscuring the result (see figure 4). The major PCR product, seen in cDNA and in DNA from the cDNA library, using oligonucleotides O.11 (upstream of intron I) and O.17 (spanning intron II) is about 440bp. This is compatible with the predicted product of 441bp, expected if intron I is spliced out. A band of slightly higher molecular weight (494bp) is seen in the unspliced cDNA clone (cDNA 2, figure 1B) and also as a minor band in cDNA and the cDNA library. This is consistent with intron I being unspliced in a minority of products, as shown by our isolation of a full length cDNA clone containing an unspliced intron I.

In the truncated cDNA clone with a splice at intron I, the AG at 815 is used as splice acceptor (this was determined by sequence analysis of both strands), this splicing does not bring an upstream exon into frame with the main ORF. However, sequence analysis of PCR products derived from two further cDNA clones showed that the splice acceptor was the AG at 811, giving rise to a 445bp PCR product. When this is the case an ORF of 35 amino acids is spliced in frame onto the main ORF. Thus intron I extends from nt 760–813.

It is likely that the start of the complementing cDNA clone (cDNA 2, figure 1b) at 510 is the 5' start of transcription of the *rad9* gene, as the truncated cDNA clone (cDNA 1, figure 1b) started at the same base. No obvious TATA motif is present upstream of this putative start site in the genomic sequence. However, this is not unusual in *S. pombe* genes, especially those that are expressed at low levels (22).

From these studies we conclude that the mRNA starts at base 510 and that initiation of protein synthesis is probably at 655, giving a maximum sized predicted protein of 427 amino acids if all three introns are spliced out. This leaves an untranslated leader region of 145bp upstream of the putative initiation codon. The putative *rad9* polypeptide does not exhibit a codon usage bias as calculated according to the method of Russell and Hall (23). This suggests that the *rad9* gene is not expressed at a high level. Low levels of expression have also been found for most *S. cerevisiae* DNA repair genes (24).

Northern blot analysis indicates that the *rad9* gene is indeed expressed at a low level, as a transcript was only visualised with 5µg polyA⁺ RNA. A single band at 1.6kb was seen with polyA⁺ RNA when probed with the internal 1.5kb *EcoRI*–*SspI* fragment (figure 5). This is consistent with the size of the complementing clone isolated from the cDNA library.

The predicted *rad9* ORF would encode a protein with a molecular weight of 47512D and an isoelectric point at pH 5.15.



Figure 5. Northern blot analysis of the *rad9* transcript. 5µg polyA⁺ RNA was probed with a 1.5kb *EcoRI*–*SspI* fragment from *rad9*. *EcoRI* *HindIII* digested single stranded lambda was used as size markers (not shown). From this the estimated size of the transcript is 1.6kb.

Computer searches, using the DNASTAR programmes and TFASTA at the Daresbury Laboratory, of the EMBL and PIR data bases for homology to the predicted *rad9* protein revealed no significant homology to any known protein or class of protein.

Mapping the Functional Limits of the *rad9* Gene

In order to map the functional limits of the *rad9* gene, the *ExoIII* deletions of the *rad9* sequence in mp19 were cotransformed with pUR19, into the *rad9.192* mutant strain (sp.039). Transformants were selected for their ability to grow in the absence of uracil and then replica plates compared for the ability of colonies to grow after UV irradiation at a Stratalinker setting of 300 Jm⁻². The smallest region that complemented the UV sensitivity was found to extend from nucleotide 496 to nucleotide 1397, in exon III. (see figure 1c for details of the deletions used). This was confirmed by repeating the functional analysis with subclones of deletions A4.5 and S13 (see figure 1c) in pUR19. These plasmids complemented the UV sensitivity when introduced into the *rad9.192* strain. This implies that only the 900bp region containing the first two exons are required to complement the mutant phenotype.

The region from the *EcoRI* site, at nucleotide 658, to the end of the S13 deletion, at nucleotide 1397 (pES13, figure 1c) was found not to complement the UV sensitivity of the *rad9.192* strain. *EcoRI* cuts one base inside the proposed initiating ATG in exon I, supporting the view that this exon is necessary for a functional *rad9* protein.

Gene Deletion

A null allele of the *rad9* gene was constructed in order to test the deletion phenotype of the gene. A plasmid construct was made in which the 0.76kb *ClaI* to *MluI* fragment (see figure 1a) was replaced with a 1.8kb DNA fragment containing the *ura4* gene (25). The resulting 3.5kb *HindIII* to *EcoRV* fragment was then used to transform the diploid strain sp.101 to uracil prototrophy. Colonies not requiring uracil were tested for stability and chromosomal DNA was prepared from stable clones. Southern blot analysis of one such stable transformant, sp.086 (figure 6) showed that the fragment had integrated into the chromosome by homologous recombination. Hybridisation of *HindIII* digested DNA from the *rad*⁺ strain sp.011 with *rad9* revealed a single 4.3kb band. A similar digest of the diploid strain heterozygous for the deletion, contained this band and an additional 5.3kb band.



Figure 6. Southern blot analysis of the null allele of *rad9*. DNA from the *rad*⁺ strain sp.011 (track 1), the diploid deletion strain sp.086 (track 2), and two products of meiosis of a h⁹⁰ derivative of sp.086, one *ura*⁺ (sp.069) (track 3) and the other *ura*⁻ (sp.071) (track 4), was digested with *HindIII*, southern blotted and hybridised with the 2.6kb *HindIII*–*EcoRV* fragment from *rad9*.

Table 4. Random spore analysis of the deletion mutant sp.086

no. of colonies analysed				
rad ⁺ ura ⁺	rad ⁺ ura ⁻	rad ⁻ ura ⁺	rad ⁻ ura ⁻	total
0	857	903	0	1760

This is the result of the disruption of one of the two copies of the *rad9* gene by the *ura4* gene. An h⁹⁰/h⁺ derivative (sp.102) was then selected and used to generate haploid spores.

The spores were then subjected to random spore analysis (Table 4) and tetrad analysis. Ten tetrads were dissected and all showed a 2:2 segregation of uracil prototrophy. Ura⁺ was always associated with radiation sensitivity. This showed that the *rad9* gene is not essential for mitotic growth.

Southern blot analysis (figure 6) showed the 5.3kb *ura4*-containing band was inherited with the radiation-sensitive phenotype in one product of meiosis (sp.069) of the diploid deletion strain, while the *ura*- product (sp.071) inherited the 4.3kb band. This formally shows that the sp.069 deletion strain contains a disrupted copy of the *rad9* gene.

Complementation Data

The *rad9.192* strain is sensitive to both UV and gamma irradiation. At a fluence of 200 Jm⁻² less than 0.01% of *rad9.192* cells survive, compared to about 10% of wild type cells. Similarly, with ionising radiation, at 1000 Gy less than 0.01% of *rad9.192* cells survive, compared to about 50% survival of wild type cells. The survival curves following UV or gamma radiation, of the *rad9.192* strain sp.039 and the equivalent *rad*⁺ control strain, sp.011, are shown in figures 2a and b.

The gamma-radiation and UV irradiation survival curves of the *rad9.192* strain sp.039 containing the complementing plasmids p9/4 and prad9, are shown in figures 2c and 2d. p9/4 is the 4.3kb *HindIII* fragment in DB262 and prad9 is the 2.6kb *HindIII-EcoRV* fragment subcloned from p9/4 in the vector pUR19. The presence of the plasmids increases the UV and gamma survival of cells to almost wild type levels. The gamma-radiation survival curves of the *rad9.192* strain sp.039 containing non-complementing vectors are also shown in figure 2c, for comparison. The equivalent UV irradiation survival curves are shown in figure 2d. No increase in survival after either UV or gamma irradiation was seen with either vector.

The null allele has also been subjected to the same analysis. The four products of a single tetrad were tested for survival after gamma- and UV-irradiation. The two null mutant strains (sp.069 and sp.070) show phenotypes virtually identical to the *rad9.192* allele and the two *ura*- strains (sp.071 and sp.072) show wild type survival (figure 2e and 2f).

DISCUSSION

We have cloned the *rad9* gene of fission yeast by complementation of the *rad9.192* strain with an *S. pombe* gene bank. Selection was carried out over three rounds of gamma-irradiation and the resulting resistant colonies were shown to be co-unstable for radiation resistance and leucine prototrophy. Thus the complementation resulted from plasmid borne sequences. Plasmid rescued from the resistant colonies complemented the radiation sensitive phenotype on reintroduction into the *rad9.191* mutant and the complementing region was localised to a 2.6kb *HindIII-EcoRV* fragment. This sequence was found to integrate

at a genomic locus 3.9cM from the *lys2* gene and to hybridise to the expected 1.2Mb *NotI* fragment. This is consistent with the localisation of *rad9* on the genetic and long range maps of *S. pombe* (19, 20).

DNA sequence analysis of the 2.6kb *HindIII-EcoRV* fragment revealed the presence of an open reading frame which could be extended by the postulation of three introns. The existence of the introns was verified by the isolation of a full length, complementing, cDNA clone and by PCR and sequence analysis of this and other cDNA clones. These studies were necessary as the splicing of the *rad9* transcript proved to be contrary to the accepted consensus data. The results indicate that the *rad9* transcript starts at base 510, with initiation of protein synthesis at base 655. Three introns (I nt 760–813, II nt 1147–1204 and III nt 1818–1874) are spliced out resulting in a predicted protein of 427 amino acids.

The third intron has an unusual branch site to 3' acceptor distance, resulting in a smaller intron than predicted from consensus splice data (22). The consensus branch site, as found in intron II, is CTPuAPy, 11 to 23 bases upstream of the 3' end of the intron (22). Intron III, however, contains the sequence CTAAG 35 bases upstream of the 3' junction. This may be analogous to the situation found in the *cam1* intron which contains the sequence CTGAA starting 34 bases upstream of the 3' junction (22).

The first intron also proved to be unusual. PCR analysis of cDNA using flanking sequences as primers, indicates that splicing occurs at this intron in the majority of cases. The first exon, with an ATG at nucleotide 655, is spliced into frame with the major ORF if the AG at 811 is used as the 3' acceptor. However, sequence analysis of a truncated cDNA clone indicated that in this case the AG at 815 was used, suggesting that the first exon could not be functional. Further analysis of sequences amplified from two other cDNA clones indicates that the AG acceptor at 811 is utilised and we believe that the splicing seen in the truncated clone is aberrant.

Intron I contains a degenerate 5' donor site (GTGTGT) as opposed to the consensus GTANGT (22) and this may explain the high level of message unspliced or mis-spliced at this intron. A transition in the third position has been shown to reduce the efficiency of splicing of a functional artificial intron in the *S. pombe ura4* gene (26). The full length cDNA clone, originally isolated from the cDNA library, is not spliced at this intron but is capable of complementing the *rad9.192* mutant. Therefore, either splicing of the transcript from this cDNA clone is completed *in vivo* or the first exon is not required for the DNA repair function. The non-complementation of a plasmid pES13, missing the ATG at 655 suggests that the first exon is required.

Alternatively, the non-splicing could be specific to a subpopulation of molecules, giving rise to an alternative protein. In this case the first exon could not be used and the *rad9* gene would contain an unusually long leader sequence, with translation starting at the ATG at base 855 in intron II. It is possible that exon I is not translated even when spliced and has a regulatory function as, for example, the untranslated first exon of the oncogene *c-myc* (27). However, this possibility awaits confirmation by N-terminal sequence analysis of the *rad9* protein.

The size of the full length cDNA clone (1.6kb) is consistent with the size of the transcript seen in northern analysis of polyA⁺ RNA when probed with the *rad9* sequence. However, the functional complementation data show that a 900bp fragment spanning the first two exons was sufficient to confer radiation

resistance to the *rad9.192* mutant. This suggests that the mutation in the *rad9.192* strain is contained within the first two exons and that only this region is required for the protein's role in DNA repair. It is possible that the *rad9* protein has more than one role and that different functions are executed by different domains of the protein. However, none of these roles are essential for mitotic growth under normal conditions, as the *rad9* null mutant is viable. The null mutant, which has the same radiation sensitive phenotype as the *rad9.192* mutant, has most of the first exon, all the second exon, and most of the third replaced by the *ura4* gene and therefore the protein is unlikely to be active.

Analysis of the 427 amino acid sequence using the DNASTAR computer program revealed no sequence motifs common to known classes of proteins and no significant homology to any known sequences was found during searches of the EMBL and NBRF-PIR data bases using the TFasta program at the Daresbury Laboratory. Thus, the *rad9* gene, presented here, like the *S. pombe radl* gene (28), is a novel DNA repair gene.

The *S. pombe rad9.192* and *rad1.1* mutants are, like most *S. pombe* repair deficient mutants, highly sensitive to both UV and ionising radiation (1). This is in contrast to the situation in *S. cerevisiae* in which the majority of mutants are sensitive to either UV or ionising radiation (24). This may reflect a difference in response to DNA damage between the two organisms. *S. pombe*, unlike *S. cerevisiae*, spends the greater part of its cell cycle in G2, where a replicated genome is available for recombination repair, and thus, may preferentially repair its DNA damage by a recombination mechanism, while in *S. cerevisiae* the dominant form of repair may be excision repair. This may be reflected in the differences in types of DNA repair mutants isolated in the two organisms. However, there is evidence that repair pathways are conserved between the two highly divergent yeasts, as a homologue to the *S. cerevisiae* gene *RAD6*, which codes for a ubiquitin carrier protein, has been identified in *S. pombe* (29). We also have evidence that at least three further DNA repair genes, all in the excision repair pathway are conserved between the two yeasts (30, 31). It will be of interest to determine whether the *rad9* gene is conserved across species boundaries. The identification of such genes would provide evidence for the evolutionary conservation of pathways other than excision repair.

Further work will include the characterisation of the *rad9* protein with a view to elucidating the role of this gene in DNA repair.

ACKNOWLEDGEMENTS

This work is supported by an MRC project grant G88/03213CB (J.M.M. and F.Z.W.) and by EC grants B16-E-142-UK (A.M.C. and A.R.L.). We are indebted to J. Brannigan and H. Steingrimsdottir for providing the oligonucleotides and PCR techniques and to A. Nasim, Riyadh, Saudi Arabia, for providing the *rad9.192* mutant. J.M.M. and F.Z.W. would like to acknowledge the receipt of travel grants from the Wellcome Trust. We also acknowledge the use of the computer facilities at the Daresbury Laboratory for the homology searches of the EMBL and NBRF data bases.

REFERENCES

- Phipps, J., Nasim, A. and Miller, D.R. (1985) *Adv. Genet.* **23**, 1–72.
- Lieberman, H.B., Riley, R. and Martel, M. (1989) *Mol. Gen. Genet.* **218**, 554–558.
- Lee, M.G. and Nurse, P. (1987) *Nature* **327**, 31–35.
- Hindley, J. and Phear, G.A. (1984) *Gene* **31**, 129–134.
- Kauffer, N.F., Simanis, V. and Nurse, P. (1985) *Nature* **318**, 78–80.
- Loprieno, N. and Shupbach, M. (1971) *Mol. Gen. Genet.* **110**, 348–354.
- Carr, A.M. (1986) D.Phil thesis. University of Sussex.
- Fikes, J., Becker, D., Winston, F. and Guarente, L. (1990) *Nature* **346**, 291–294.
- Barbet, N., Muriel, W.J. and Carr, A.M. (1991) in preparation.
- Hanahan, D. (1983) *J. Mol. Biol.* **166**, 557–568.
- Beach, D.B., Piper, M. and Nurse, P. (1982) *Mol. Gen. Genet.* **187**, 326–329.
- Gutz, H., Heslot, H., Leupold, U. and Loprieno, N. (1974) In King R.C. (ed.), *Handbook of Genetics*, Plenum Press, New York Vol. 1, pp 395–446.
- Maniatis, T., Fritsch, E.F. and Sambrook, J. (1982) *Molecular cloning: A laboratory manual*. Cold Spring Harbor University Press, Cold Spring Harbor.
- Aves, S., Durkacz, B., Carr, A.M. and Nurse, P. (1985) *EMBO J.* **4**, 457–463.
- Durkacz, B., Carr, A.M. and Nurse, P. (1986) *EMBO J.* **5**, 369–373.
- Henikoff, S. (1984) *Gene* **28**, 351–359.
- Yang, J.-L., Maher, V.M. and McCormick, J.J. (1989) *Gene* **83**, 347–354.
- Saiki, R.K., Gelfand, D.H., Stoffel, S., Scharf, S.J., Higuchi, R., Horn, G.T., Mullis, K.B. and Erlich, H.A. (1988) *Science* **239**, 487–494.
- Gygax, A. and Thuriaux, P. (1984) *Curr. Genet.* **8**, 85–92.
- Fan, J.-B., Chikashige, Y., Smith, C., Niwa, O., Yanagida, M. and Cantor, C. (1988) *Nucleic Acids Res.* **17**, 2801–2818.
- Murray, J.M. and Watts, F.Z. (1990) *Nucleic Acids Res.* **18**, 4590.
- Russell, P.R. (1989). In Nasim, A., Young, P., Johnson, B. (eds.), *Molecular Biology of Fission Yeast*. Academic Press Inc., London. p243–271.
- Russell, P.R. and Hall, B.D. (1983) *J. Biol. Chem.* **258**, 143–149.
- Friedberg, E. (1988) *Micro. Revs.* **52**, 70–102.
- Grimm, C., Kohli, J., Murray, J.M. and Maundrell, K. (1988) *Mol. Gen. Genet.* **215**, 81–86.
- Gatermann, K.B., Hoffmann, A., Rosenberg, G.H. and Kauffer, N.F. (1989) *Mol. Cell. Biol.* **9**, 1526–1535.
- Batley, J., Moulding, C., Taub, R., Murphy, W., Stewart, T., Potter, H., Lenoir, G. and Leder, P. (1983) *Cell* **34**, 779–789.
- Sunnerhagen, P., Seaton, B.L., Nasim, A. and Subramani, S. (1990) *Mol. Cell. Biol.* **10**, 3750–3760.
- Reynolds, P., Koken, M.H.M., Hoeijmakers, J.H.J., Prakash, S. and Prakash, L. (1990) *EMBO J.* **9**, 1423–1430.
- Murray, J.M., Doe, C., Schenk, P., Carr, A.M., Lehmann, A.R. and Watts, F.Z. (1991) in preparation.
- Carr, A.M., Barbet, N., Murray, J.M., Muriel, W.J., Watts, F.Z. and Lehmann, A.R. (1991) in preparation.