

Measurement of the underground atmospheric muon charge ratio using the MINOS Near Detector

Article (Published Version)

Adamson, P, Andreopoulos, C, Armstrong, R, Auty, D.J, Ayres, D S, Backhouse, C, Barr, G, Bishai, M, Blake, A, Block, G.J, Devenish, N E, Falk, E, Hartnell, J, Lefevre, G and et al, (2011) Measurement of the underground atmospheric muon charge ratio using the MINOS Near Detector. Physical Review D, D83 (3). 032011. ISSN 1550-7998

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/27077/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Measurement of the underground atmospheric muon charge ratio using the MINOS Near Detector

P. Adamson,⁸ C. Andreopoulos,²³ D. J. Auty,²⁷ D. S. Ayres,¹ C. Backhouse,²¹ G. Barr,²¹ W. L. Barrett,³² P. Bhattarai,¹⁹ M. Bishai,⁴ A. Blake,⁶ G. J. Bock,⁸ D. J. Boehnlein,⁸ D. Bogert,⁸ S. Budd,¹ S. Cavanaugh,¹⁰ D. Cherdack,³⁰ S. Childress,⁸ B. C. Choudhary,^{8,†} J. A. B. Coelho,⁷ S. J. Coleman,³³ L. Corwin,¹³ D. Cronin-Hennessy,¹⁸ D. Damiani,^{33,‡} I. Z. Danko,²² J. K. de Jong,^{21,12} N. E. Devenish,²⁷ M. V. Diwan,⁴ M. Dorman,¹⁷ C. O. Escobar,⁷ J. J. Evans,¹⁷ E. Falk,²⁷ G. J. Feldman,¹⁰ T. H. Fields,¹ M. V. Frohne,^{11,3} H. R. Gallagher,³⁰ R. A. Gomes,⁹ M. C. Goodman,¹ P. Gouffon,²⁴ N. Graf,¹² R. Gran,¹⁹ N. Grant,²³ K. Grzelak,³¹ A. Habig,¹⁹ D. Harris,⁸ P. G. Harris,²⁷ J. Hartnell,^{27,23} R. Hatcher,⁸ A. Himmel,⁵ A. Holin,¹⁷ X. Huang,¹ J. Hylen,⁸ J. Ilic,²³ G. M. Irwin,²⁶ Z. Isvan,²² D. E. Jaffe,⁴ C. James,⁸ D. Jensen,⁸ T. Kafka,³⁰ S. M. S. Kasahara,¹⁸ G. Koizumi,⁸ S. Kopp,²⁹ M. Kordosky,³³ Z. Krahn,¹⁸ A. Kreymer,⁸ K. Lang,²⁹ G. Lefevvre,²⁷ J. Ling,^{4,25} P. J. Litchfield,¹⁸ L. Loiacono,²⁹ P. Lucas,⁸ W. A. Mann,³⁰ M. L. Marshak,¹⁸ N. Mayer,¹³ A. M. McGowan,^{1,§} R. Mehdiev,²⁹ J. R. Meier,¹⁸ M. D. Messier,¹³ D. G. Michael,^{5,*} W. H. Miller,¹⁸ S. R. Mishra,²⁵ J. Mitchell,⁶ C. D. Moore,⁸ J. Morfín,⁸ L. Mualem,⁵ S. Mufson,¹³ J. Musser,¹³ D. Naples,²² J. K. Nelson,³³ H. B. Newman,⁵ R. J. Nichol,¹⁷ J. A. Nowak,¹⁸ W. P. Oliver,³⁰ M. Orchanian,⁵ J. Paley,^{1,13} R. B. Patterson,⁵ G. Pawloski,²⁶ G. F. Pearce,²³ R. Pittam,²¹ R. K. Plunkett,⁸ X. Qiu,²⁶ J. Ratchford,²⁹ T. M. Raufer,²³ B. Rebel,⁸ J. Reichenbacher,^{1,||} P. A. Rodrigues,²¹ C. Rosenfeld,²⁵ H. A. Rubin,¹² V. A. Ryabov,¹⁵ M. C. Sanchez,^{14,1,10} N. Saoulidou,⁸ J. Schneps,³⁰ P. Schreiner,³ P. Shanahan,⁸ A. Sousa,¹⁰ M. Strait,¹⁸ N. Tagg,²⁰ R. L. Talaga,¹ J. Thomas,¹⁷ M. A. Thomson,⁶ G. Tinti,²¹ R. Toner,⁶ G. Tzanakos,² J. Urheim,¹³ P. Vahle,³³ B. Viren,⁴ A. Weber,²¹ R. C. Webb,²⁸ C. White,¹² L. Whitehead,⁴ S. G. Wojcicki,²⁶ D. M. Wright,¹⁶ T. Yang,²⁶ and R. Zwaska⁸

(MINOS Collaboration)

¹Argonne National Laboratory, Argonne, Illinois 60439, USA²Department of Physics, University of Athens, GR-15771 Athens, Greece³Physics Department, Benedictine University, Lisle, Illinois 60532, USA⁴Brookhaven National Laboratory, Upton, New York 11973, USA⁵Lauritsen Laboratory, California Institute of Technology, Pasadena, California 91125, USA⁶Cavendish Laboratory, University of Cambridge, Madingley Road, Cambridge CB3 0HE, United Kingdom⁷Universidade Estadual de Campinas, IFGW-UNICAMP, CP 6165, 13083-970, Campinas, SP, Brazil⁸Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA⁹Instituto de Física, Universidade Federal de Goiás, CP 131, 74001-970, Goiânia, GO, Brazil¹⁰Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA¹¹Holy Cross College, Notre Dame, Indiana 46556, USA¹²Physics Division, Illinois Institute of Technology, Chicago, Illinois 60616, USA¹³Indiana University, Bloomington, Indiana 47405, USA¹⁴Department of Physics and Astronomy, Iowa State University, Ames, Iowa 50011 USA¹⁵Nuclear Physics Department, Lebedev Physical Institute, Leninsky Prospect 53, 119991 Moscow, Russia¹⁶Lawrence Livermore National Laboratory, Livermore, California 94550, USA¹⁷Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, United Kingdom¹⁸University of Minnesota, Minneapolis, Minnesota 55455, USA¹⁹Department of Physics, University of Minnesota–Duluth, Duluth, Minnesota 55812, USA²⁰Otterbein College, Westerville, Ohio 43081, USA²¹Subdepartment of Particle Physics, University of Oxford, Oxford OX1 3RH, United Kingdom²²Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA²³Rutherford Appleton Laboratory, Science and Technologies Facilities Council, OX11 0QX, United Kingdom²⁴Instituto de Física, Universidade de São Paulo, CP 66318, 05315-970, São Paulo, SP, Brazil²⁵Department of Physics and Astronomy, University of South Carolina, Columbia, South Carolina 29208, USA²⁶Department of Physics, Stanford University, Stanford, California 94305, USA

*Deceased.

†Present address: Department of Physics and Astrophysics, University of Delhi, Delhi 110007, India.

‡Present address: Santa Cruz Institute for Particle Physics, Department of Physics, University of California at Santa Cruz, Santa Cruz, CA 95064, USA.

§Present address: Department of Physics and Astronomy, University of Rochester, NY 14627, USA.

||Present address: Department of Physics and Astronomy, University of Alabama, Tuscaloosa, AL 35487, USA.

²⁷*Department of Physics and Astronomy, University of Sussex, Falmer, Brighton BN1 9QH, United Kingdom*

²⁸*Physics Department, Texas A&M University, College Station, Texas 77843, USA*

²⁹*Department of Physics, University of Texas at Austin, 1 University Station C1600, Austin, Texas 78712, USA*

³⁰*Physics Department, Tufts University, Medford, Massachusetts 02155, USA*

³¹*Department of Physics, Warsaw University, Hoża 69, PL-00-681 Warsaw, Poland*

³²*Physics Department, Western Washington University, Bellingham, Washington 98225, USA*

³³*Department of Physics, College of William & Mary, Williamsburg, Virginia 23187, USA*

(Received 16 December 2010; published 28 February 2011)

The magnetized MINOS Near Detector, at a depth of 225 mwe, is used to measure the atmospheric muon charge ratio. The ratio of observed positive to negative atmospheric muon rates, using 301 days of data, is measured to be $1.266 \pm 0.001(\text{stat})_{-0.014}^{+0.015}(\text{syst})$. This measurement is consistent with previous results from other shallow underground detectors and is $0.108 \pm 0.019(\text{stat} + \text{syst})$ lower than the measurement at the functionally identical MINOS Far Detector at a depth of 2070 mwe. This increase in charge ratio as a function of depth is consistent with an increase in the fraction of muons arising from kaon decay for increasing muon surface energies.

DOI: 10.1103/PhysRevD.83.032011

PACS numbers: 13.85.Tp, 95.55.Vj, 95.85.Ry

I. INTRODUCTION

High energy cosmic-ray primaries interact with nuclei in the upper atmosphere and produce showers which contain pions and kaons. These secondary mesons can either interact in further collisions in the atmosphere or decay to produce atmospheric muons. Since the majority of primary cosmic rays are protons, there is an excess of positively charged mesons in the showers, and consequently, the atmospheric muon charge ratio N_{μ^+}/N_{μ^-} is larger than unity. A precise measurement of the atmospheric muon charge ratio can be used to constrain cosmic-ray shower models and, since atmospheric neutrinos are produced in conjunction with atmospheric muons, better determine atmospheric neutrino fluxes.

The differential muon production spectrum in extensive air showers can be parameterized as [1]

$$\frac{dN_{\mu}}{dE_{\mu}} \approx \frac{0.14E_{\mu}^{-2.7}}{\text{cm}^2 \text{ s sr GeV}} \left(\frac{1.0}{1 + \frac{1.1E_{\mu} \cos\theta^*}{\epsilon_{\pi}}} + \frac{0.054}{1 + \frac{1.1E_{\mu} \cos\theta^*}{\epsilon_K}} \right), \quad (1)$$

where E_{μ} is the muon surface energy and θ^* is the zenith angle at the muon production point. Accounting for the curvature of the Earth, this angle is geometrically related to the observed zenith angle θ :

$$\cos\theta^* = \sqrt{1 - \sin^2\theta \left(\frac{R_e}{R_e + h} \right)^2}, \quad (2)$$

where R_e is the radius of the Earth and h is the muon production height for horizontal muons. Assuming the mean column depth for the initial cosmic-ray primary interaction to be 85 g/cm^2 gives $h = 30 \text{ km}$ [2]. The two terms in Eq. (1) represent the contribution to muon production from pion and kaon decay, respectively. The values $\epsilon_{\pi} = 115 \text{ GeV}$ and $\epsilon_K = [850] \text{ GeV}$ are the critical energies at the muon production height above which the pion and kaon interaction probability exceeds the decay probability. The larger value of ϵ_K implies that the kaon

contribution to the muon flux, and therefore to the charge ratio, will be more significant at values of the vertical muon surface energy $E_{\mu} \cos\theta^*$ exceeding ϵ_{π} . The charge ratio is expected to increase as a consequence of the increasing kaon contribution to the muon flux because in high energy interactions, single K^+ 's can be produced in associated production with strange baryons while single K^- 's cannot. The muon charge ratio is expected to decrease at even higher energies due to heavy flavor production [3,4] and changes in composition of the cosmic-ray primaries [5–7]. However, as these latter two processes are expected to only affect the charge ratio at energies greater than those studied here they are not considered further.

Atmospheric muon charge ratio measurements utilizing deep underground detectors are typically higher in value than those produced using detectors which are shallower since they sample muons with a larger value of $E_{\mu} \cos\theta^*$. A previous measurement of the muon charge ratio utilizing the MINOS Far Detector, at a depth of 2070 mwe, reported a value of $1.374 \pm 0.004(\text{stat})_{-0.010}^{+0.012}(\text{syst})$ [8] for surface energies greater than 1 TeV. Another measurement [9] of the charge ratio at TeV energies, conducted using the OPERA experiment located at a depth of 3800 mwe, is in good agreement with the MINOS Far Detector result. Atmospheric muon charge ratio measurements performed by the L3 + C [10], Bess-TEV [11], and CMS [12] Collaborations, for muons with surface energy E_{μ} above 10 GeV and below 300 GeV, are consistent with the 2001 world average of $1.268 \pm [0.008 + 0.0002 \cdot \frac{E_{\mu}}{\text{GeV}}]$ [13].

The MINOS Near and Far Detectors [14] are large underground magnetic calorimeters at depths of 225 and 2070 mwe, respectively. The detectors are designed to study neutrino interactions from the Fermilab Neutrinos at the Main Injector beam [15] but also trigger on atmospheric muons between beam spills. The depths of the MINOS detectors are ideal for making a measurement of the muon charge ratio at values of $E_{\mu} \cos\theta^*$ in the region

dominated by the pion contribution and the transition region where the kaon contribution becomes significant. In this paper we present a measurement of the atmospheric muon charge ratio using the MINOS Near Detector. This result is then compared to the same measurement performed using the MINOS Far Detector [8].

This paper is organized as follows. Sections II and III describe the MINOS Near Detector and the atmospheric muon Monte Carlo simulation, respectively. Section IV describes the criteria for selecting atmospheric muon tracks with correct charge sign identification. Section V outlines the technique used to combine the data collected from two detector magnetic field configurations to determine the atmospheric muon charge ratio. Section VI also elaborates on the evaluation of the systematic uncertainties. Finally, Sec. VI presents the results of this analysis.

II. THE MINOS NEAR DETECTOR

The MINOS Near Detector [14] at Fermilab is a magnetized-steel and scintillator sampling calorimeter. It is located 94 m underground, with an approximately flat overburden of 225 mwe. Its dimensions are $3.8 \times 4.8 \times 16.6 \text{ m}^3$. The detector contains 282 vertical steel planes, each 2.54 cm thick. The scintillator layers are composed of either 64 or 96, 4.1 cm wide and 1 cm thick strips which vary in length from 2.5 to 4 m. The scintillator strips are rotated by 90° with respect to the previous layer to allow for three-dimensional track reconstruction. The first 120 steel planes each have a scintillator layer attached to their surface. The scintillator layer of every fifth steel plane contains 96 strips and is said to be fully instrumented. The remaining four planes in each set of five are partially instrumented and contain 64 strips. Of the last 162 steel planes, only every fifth plane has an attached scintillator layer. These scintillator layers are fully instrumented.

Scintillation light is collected by wavelength shifting fibers embedded in the scintillator strips. Each strip is coupled to a single pixel on a 64-pixel multianode photomultiplier tube [16] by a clear fiber. Each photomultiplier tube pixel is digitized continuously at a frequency of 53 MHz. The detector response for a candidate atmospheric muon event is recorded either when four strips in five sequential planes or when strips from any 20 planes register a signal above the $1/3$ photoelectron dynode threshold within 151 ns.

The detector's approximately toroidal magnetic field [14] varies in strength from 2.1 T near the magnetic coil hole to 0.2 T near the periphery of the steel planes. The magnetic field can be oriented to focus either northerly going μ^- or μ^+ . These magnetic field orientations will be referred to as "forward" and "reverse," respectively. The detector is oriented horizontally and its long axis points 26.5548° west of north. The curvature induced by the magnetic field together with the three-dimensional track

reconstruction allows the determination of the charge sign of a muon traversing the detector.

Detailed information regarding the MINOS Near Detector, the electronics, and the data acquisition systems can be found in [14,17,18].

III. SIMULATED ATMOSPHERIC MUONS

A sample of 42×10^6 simulated atmospheric muon events are used to evaluate the analysis sensitivities and efficiencies, as well as to help assess systematic errors. Surface level muon events were generated by the HEMAS [19] atmospheric cascade simulation. Muons expected to intersect with the detector are propagated through a GEANT3 [20] model of the overburden and the MINOS Near Detector.

IV. EVENT SELECTION

The selection criteria have been chosen to optimize the event selection efficiency observed in the data and the charge sign identification purity obtained from the Monte Carlo simulations. The data sample for this analysis consists of atmospheric muons with well-reconstructed energy and charge sign. Preselections are made to reject events that are not consistent with the passage of an atmospheric muon through the detector. Further selections are applied to ensure track reconstruction quality and good charge sign determination. A summary of the selection criteria, as well as the selection efficiencies and charge identification purities, can be found in Table I.

This analysis uses data collected between 2006 and 2009. During the data taking period, the direction of the magnetic field was periodically changed between the two orientations referred to as forward and reverse magnetic fields. The final reverse magnetic field exposure was 150.5 days. Equivalent exposures of forward magnetic field data were collected immediately following the reverse magnetic field data periods to reduce systematic uncertainties. A total of 7.16×10^8 triggers were collected over a combined 301 days of exposure.

A. Preselections and track quality

For an event to be included in this analysis it must possess a single downward-going atmospheric muon track, have been collected during a period of good detector run conditions, and have a reconstructed initial interaction point within 50 cm of the detector edge. If the reconstructed initial interaction point is further than 3 cm outside the detector volume, the track is rejected. The curvature of the track is known to be poorly determined, and the track is rejected if any scintillator strips hit are further than 3 cm from the reconstructed position of the track or if the track does not pass through a region of the detector where there is a scintillator on each layer of steel.

Track reconstruction errors may occur when the event contains a large amount of activity which is not related to the track. These extra hits degrade the charge sign determination and can be generated by muon bremsstrahlung, natural radioactivity, or electrical and optical cross talk between the channels on the multianode photomultiplier tube [16]. Events are rejected if more than 40% of the strips hit are not track-related. Muon tracks determined to be poorly reconstructed by internal consistency checks of the reconstruction algorithm are also removed from the data sample.

B. Charge sign quality selection

Two selection variables are used to further increase the degree of confidence in the assigned curvature and charge sign of the track. The Kalman filter [21] used in the track curvature fitting provides an uncertainty $\sigma(q/p)$ on the measured value of q/p , where q is the charge and p the momentum of the track. The first charge sign quality selection is based on the value of $(q/p)/\sigma(q/p)$ determined by the track fitter. The second selection variable BdL is defined to be equivalent to $\sum_{i=1}^N |B_i \times dL_i|$, where N is the total number of planes in the muon track and B_i and dL_i are the magnetic field and muon path length vectors, respectively, at plane i . A selection based on BdL is used to ensure that the magnitude of the bending due to curvature is larger than the apparent bending due to multiple scattering. Figure 1 shows the muon charge ratio as a function of $(q/p)/\sigma(q/p)$ and Fig. 2 the charge ratio as a function of BdL , for data collected in both magnetic field orientations as well as the combined data set. The charge ratio for data collected during a single magnetic field orientation is defined as the ratio of positive to negative muons collected only in that orientation. The




FIG. 1. Charge ratio as a function of $(q/p)/\sigma(q/p)$ after all selections and requiring that $BdL > [3.0]$ T m. The vertical line is the $(q/p)/\sigma(q/p)$ threshold value used in the charge sign quality selection.




FIG. 2. Charge ratio as a function of BdL , after all selections and requiring that $(q/p)/\sigma(q/p) > 3.0$. The vertical line is the BdL threshold value used in the charge sign quality selection.

observed variation in the charge ratio above the selection thresholds in a single field orientation, as well as the difference in the charge ratio between the two different field orientations, stems from acceptance effects due to the magnetic field, detector asymmetry, and detector alignment errors. The technique used to combine data taken in the two field orientations and remove these biases is discussed in the next section. For this analysis we required that $BdL > [3.0]$ T m and $(q/p)/\sigma(q/p) > 3.0$. Below these values the charge ratio tends towards unity, indicating a degradation of the charge sign determination. Events which have passed all the selections described in this section are used in the calculation of the atmospheric muon charge ratio described in the next section.

V. CHARGE RATIO DETERMINATION

As demonstrated in the previous section acceptance effects in the Near Detector introduce a bias in the charge ratio when it is calculated using only data from a single magnetic field orientation. Figure 3 shows the charge ratio as a function of azimuthal angle, a variable sensitive to these biases, for the forward and reverse magnetic field data sets. Canceling these biases is done in the same manner as described in [8,22]. If ϵ_1 is the efficiency for the selection of μ^+ and ϵ_2 is the selection efficiency of μ^- in the forward field direction (FF), then the selection efficiencies for μ^+ and μ^- in the reverse field direction (RF) are ϵ_2 and ϵ_1 , respectively. Two independent equations for the charge ratio, r_a and r_b , can be constructed in which the acceptance effects cancel. These ratios, corrected for live time, are

$$r_a = (N_{FF}^{\mu^+} / t_{FF}) / (N_{RF}^{\mu^-} / t_{RF}) \quad (3)$$

and




FIG. 3. The observed charge ratio as a function of azimuthal angle. The charge ratio varies as a function of the azimuthal angle in the forward and reversed data sets due to acceptance effects and alignment errors. When data from the two fields are combined using Eq. (5) a flat distribution is obtained, indicating that the residual uncertainty due to these biases is small. The uncertainties shown are statistical only.

$$r_b = (N_{\text{RF}}^{\mu^+}/t_{\text{RF}})/(N_{\text{FF}}^{\mu^-}/t_{\text{FF}}), \quad (4)$$

where N^{μ^+} (N^{μ^-}) is the number of positive (negative) muons observed during an exposure time t in a particular magnetic field orientation. The geometric mean of r_a and r_b ,

$$\frac{N_{\mu^+}}{N_{\mu^-}} = \sqrt{r_a r_b} = \sqrt{\left(\frac{N_{\text{FF}}^{\mu^+}}{N_{\text{FF}}^{\mu^-}}\right)\left(\frac{N_{\text{RF}}^{\mu^+}}{N_{\text{RF}}^{\mu^-}}\right)}, \quad (5)$$

provides a measurement of the charge ratio that is free of biases due to geometric acceptance effects, alignment errors, and the different magnetic field live times.

Figure 3 illustrates that the significant bias in the charge ratio, that is apparent in a single field orientation data set, is strongly suppressed in the geometric mean.

After the selection criteria in Table I have been applied we obtain a final data set of 3 234 066 events. In the forward field sample we select 893 854 μ^+ and 721 428 μ^- . In the reverse field sample we select 912 944 μ^+ and 705 840 μ^- . The resulting charge ratio obtained by applying Eq. (5) is $N_{\mu^+}/N_{\mu^-} = 1.266 \pm 0.001(\text{stat})$.

A. Systematic uncertainties

The event selection criteria are chosen to remove events in which the charge sign of the muon track has been incorrectly assigned. Systematic uncertainties associated

TABLE I. Summary of the event selection. Each row shows the total number of events remaining after all the applied cuts; in parenthesis, the percentage of events remaining; and last, the percentage of the remaining Monte Carlo events which have their charge sign determined correctly.

Selection	Data selection efficiency	Monte Carlo charge ID purity
Number of triggers	7.16×10^8 (100%)	...
Preselections		
Single track events	3.18×10^8 (44.46%)	70.7%
Detector quality	3.16×10^8 (44.19%)	70.7%
Track quality selections		
Reconstructed initial interaction point and curvature selections	1.01×10^8 (14.19%)	72.9%
Track-related activity	6.13×10^7 (8.56%)	77.5%
Good reconstruction	5.31×10^7 (7.42%)	78.8%
Charge sign quality selections		
$(q/p)/\sigma(q/p) > 3.0$	1.11×10^7 (1.55%)	97.0%
$BdL > 3.0$ T m	3.23×10^6 (0.45%)	99.5%

with event selection are determined by establishing the sensitivity of the measured charge ratio to variations in the selections above their thresholds. Two selections, track-related activity and $(q/p)/\sigma(q/p)$, display variations in the observed charge ratio above their thresholds. These data were divided into three equal statistics samples, corresponding to increasing confidence in the muon charge sign, and their charge ratios calculated. The maximum deviation from the nominal charge ratio, 0.012 and 0.006 for the track-related activity and $(q/p)/\sigma(q/p)$ selections, respectively, is taken as the systematic uncertainty associated with that particular selection.

Another systematic uncertainty relates to the remaining events that have a misidentified charge sign after the selections. Since the atmospheric muon charge ratio is a value greater than unity, more positive than negative muons will have their charge sign misidentified. Thus charge sign misidentification can only decrease the measured charge ratio resulting in a positive one-sided systematic uncertainty. Monte Carlo studies suggest that 0.5% of the events in the final data sample have an incorrect charge sign determination. However, one cannot discount the possibility that the misidentification rate is different in the data than in the Monte Carlo simulation. For this analysis it is assumed that the true charge sign misidentification rate in the data differs from that in the nominal Monte Carlo by an energy independent factor α . The true charge sign misidentification rate in the data can be estimated by exploiting the fact that it influences the shape of the charge ratio versus $(q/p)/\sigma(q/p)$ curve shown in Fig. 1. A toy Monte Carlo model was written to produce similar curves assuming a charge ratio R_a , and a charge sign misidentification rate equivalent to that in the nominal Monte Carlo simulation scaled by a factor α . The difference between the measured charge ratio and R_a for the values of α and R_a that provide the best agreement between the data and the Monte Carlo prediction is taken as a systematic uncertainty on the measured charge ratio. Figure 4 plots the charge ratio as a function of $(q/p)/\sigma(q/p)$, along with the nominal and best-fit Monte Carlo results. The best-fit Monte Carlo result is obtained using a charge ratio R_a of 1.272. A similar study was performed on the curvature of the charge ratio versus *BdL* data in Fig. 2; in that study the best-fit Monte Carlo result is obtained with an actual charge ratio of 1.266. The maximum deviation of R_a from the nominal charge ratio is +0.006, which is taken as the one-sided systematic uncertainty associated with charge sign misidentification uncertainties.

Two alternative charge confidence selection criteria, similar in motivation to the *BdL*, have been investigated. The first alternative required that the muon traverse at least 27 planes at positions covered by the partially instrumented scintillator planes, within 1.75 m of the magnetic coil hole. The second alternative required that at least 37 planes be traversed where the $|B \times dL|$ for each of those planes is




FIG. 4. Charge ratio as a function of $(q/p)/\sigma(q/p)$, compared with the nominal and best-fit charge misidentification Monte Carlo simulations.

greater than 0.03 Tm. These particular selection criteria have been chosen to optimize data selection efficiency while maintaining a charge misidentification rate in the Monte Carlo similar to that of the default *BdL* selection. The maximum deviation from the nominal charge ratio was observed to be 0.003. This value is taken as the systematic uncertainty associated with the *BdL* selection.

An imperfect reversal of the detector's magnetic field would introduce a geometric bias and lead to a systematic uncertainty in the charge ratio stemming from imperfect acceptance asymmetry cancellation. This error can be determined by evaluating the charge ratios calculated using Eq. (3) and (4) since the quantities r_a and r_b will diverge as the magnitude of this bias increases. It is found that r_a and r_b agree to within their statistical uncertainties, indicating that the magnetic field bias is negligible compared to the statistical uncertainty.

Table II lists the systematic errors that are discussed above and their assigned values. Including systematic uncertainties the atmospheric muon charge ratio measured at the MINOS Near Detector is $1.266 \pm 0.001(\text{stat})^{+0.015}_{-0.014} \times$ (syst).

TABLE II. Summary of the systematic uncertainties in this analysis. The total systematic uncertainty is the quadratic sum of the individual uncertainties. The magnitude of these uncertainties are independent of E_μ and $E_\mu \cos\theta^*$.

Uncertainty classification	$\Delta \frac{N_{\mu^+}}{N_{\mu^-}}$
Track-related activity	± 0.012
$(q/p)/\sigma(q/p)$	± 0.006
<i>BdL</i> selection	± 0.003
Charge misidentification	+0.006
Total systematic uncertainty	$+0.015$ -0.014




FIG. 5. The reconstructed momentum of the muon tracks which survive all the selection criteria outlined in Table I. The mean momentum is 15 GeV.

B. The muon charge ratio underground

The maximum momentum for which the charge sign of a muon track can accurately be determined with the MINOS Near Detector is limited by the scintillator granularity and the strength of the magnetic field. The reconstructed momenta $p_{\mu,det}$ of all muon tracks which pass the selections outlined in Table I is shown in Fig. 5. The maximum momentum of muons used in this analysis is approximately 300 GeV; the mean momentum is 15 GeV. The observed charge ratio as a function of reconstructed track momentum underground is shown in Fig. 6. The charge ratio is observed to be flat as a function of track momentum to within the statistical uncertainty of the measurements.




FIG. 6. Charge ratio as a function of reconstructed underground track momentum. The y-axis uncertainties are statistical. The x-axis uncertainties are the width of the momentum bins used; the data are plotted at the median momentum values. The dotted horizontal line is the best-fit charge ratio of 1.266.

VI. THE ATMOSPHERIC MUON CHARGE RATIO AT THE SURFACE

The muon energies measured at the Near Detector depth must now be converted to energies at the surface of the Earth by accounting for energy lost by the muons in the overburden above the detector. For muons, the energy loss in matter can be parameterized by

$$-\frac{dE}{dX} = a(E_{\mu}) + \sum_{n=1}^3 b_n(E_{\mu})E_{\mu}, \quad (6)$$

where X is the slant depth; a is the ionization term and the b_n account for the radiative energy loss from bremsstrahlung, pair production, and photonuclear processes. These parameters have a mild energy dependence, whose values for standard rock can be found in [23]. Using the technique outlined in [24], the energy lost by muons traversing the Near Detector overburden has been calculated as a function of reconstructed track energy and zenith angle. These energies have been calculated assuming a flat vertical




FIG. 7. The extrapolated (a) muon surface energy and (b) $E_{\mu} \cos\theta^*$ distributions of muons passing all the analysis selection criteria. The mean surface energy is 152 GeV and the mean $E_{\mu} \cos\theta^*$ is 66 GeV.

TABLE III. The charge ratio, E_μ , and $E_\mu \cos\theta^*$, in equal bins of $\cos\theta^*$. The range of energies observed in each $\cos\theta^*$ bin is comparable to the energy resolution expected from the surface extrapolation. The charge ratio is observed to be independent of $\cos\theta^*$, E_μ , and $E_\mu \cos\theta^*$ to within the uncertainties of our measurements.

$\cos\theta^*$		Surface energy [GeV]		$E_\mu \cos\theta^*$ [GeV]		Charge ratio $^{+0.015}_{-0.014}$ (syst)	
Interval	Median	Interval	Median	Interval	Median	r	\pm (stat)
>0.9	0.914	55–67	61.5	53.5–61.0	56.5	1.257	0.069
0.8–0.9	0.824	60–77	70.5	54.2–62.9	58.0	1.270	0.013
0.7–0.8	0.735	69–91	80.5	54.6–65.0	59.2	1.277	0.005
0.6–0.7	0.642	79–112	94.5	55.2–69.0	60.8	1.270	0.003
0.5–0.6	0.549	95–142	115	56.2–74.0	62.7	1.269	0.003
0.4–0.5	0.452	115–191	145	57.4–80.9	64.8	1.259	0.003
0.3–0.4	0.356	148–280	195	59.4–91.0	68.0	1.263	0.003
0.2–0.3	0.262	212–484	290	63–109	75	1.263	0.005
0.15–0.2	0.181	382–864	513	76–140	92	1.276	0.013
<0.15	0.139	697–3450	964	104–390	134	1.249	0.032

overburden of 224.6 mwe, which is comprised of two distinct geological layers. Above the cavern hall lies 72.1 m of dolomite-shale bedrock with a density of 2.41 g/cm³, followed by 22.2 m of glacial till with a density of 2.29 g/cm³. It has been suggested in [7] that at the MINOS Near Detector depth the slightly higher rate of energy loss of μ^+ over μ^- [25–30] could lead to a surface charge ratio that is slightly higher than that observed underground. However, as the magnitude of this effect is negligible when compared to the systematic uncertainties of this measurement we have assumed the same energy loss function for both charges.

The surface energy resolution obtained with this extrapolation method is dependent on the accuracy of the overburden model and the muon pointing accuracy. Topographical measurements [31] show that the surface altitude varies by no more than 13 m within 3 km of the detector. This contributes a 14% uncertainty on the surface energy estimation at all zenith angles. The zenith pointing resolution of the Near Detector has been determined to be $1.1 \pm 0.2^\circ$ by measuring the zenith angle separation between the two muons in multimoon cosmic-triggered events for which both muons pass the charge ratio selection criteria. The corresponding error on the surface energy is negligible for vertical muons increasing to 5% at 70° , 14% at 81° , 25% at 85° , and 50% at 87° .

The extrapolated muon surface energy distribution E_μ is plotted in Fig. 7(a). The muons which populate the high energy tail have a large zenith angle and thus pass through the largest amount of matter before reaching the detector. To first approximation the minimum energy a surface muon needs to reach the Near Detector is 52 GeV/ $\cos\theta^*$. The $E_\mu \cos\theta^*$ distribution is much narrower than the E_μ distribution as the $\cos\theta^*$ and $\cos\theta$ terms effectively cancel. Because of this cancellation the muon events which occupy the high energy tail in Fig. 7(b) are those which possess the highest reconstructed momentum at the detector.

Table III presents the measured atmospheric muon charge ratio, E_μ , and $E_\mu \cos\theta^*$ in equal bins of $\cos\theta^*$. The median surface energy increases as $\cos\theta^*$ decreases due to the increasingly large overburden. $E_\mu \cos\theta^*$ increases more slowly than E_μ as the increase is due primarily to the detector's larger analyzable momentum range as a function of zenith angle [32]. The charge ratio is observed to be independent of $\cos\theta^*$, E_μ , and $E_\mu \cos\theta^*$ to within the uncertainties of our measurements. The charge ratio measurement at $E_\mu = 964$ GeV is consistent with the MUTRON spectrograph measurement [22] but is lower than other TeV energy scale measurements [8,9,33]. This difference is due to the range of zenith angles sampled by each of the detectors and the $\cos\theta^*$ dependency of Eq. (1).

The “ π K” [7,8] model is derived from the differential muon production spectrum parameterization given in Eq. (1) and predicts that the muon charge ratio is only dependent on $E_\mu \cos\theta^*$. The π K model is a qualitative model describing the relative contribution of pions and kaons to the atmospheric muon charge ratio. Following the prescription in [8] and defining f_π and f_K as the fraction of all decaying pions and kaons which decay with a detected μ^+ , the atmospheric muon charge ratio N_{μ^+}/N_{μ^-} is given in Eq. (7):

$$N_{\mu^+}/N_{\mu^-} = \frac{\left\{ \frac{f_\pi}{1+1.1E_\mu \cos\theta^*/\epsilon_\pi} + \frac{0.054 \times f_K}{1+1.1E_\mu \cos\theta^*/\epsilon_K} \right\}}{\left\{ \frac{1-f_\pi}{1+1.1E_\mu \cos\theta^*/\epsilon_\pi} + \frac{0.054 \times (1-f_K)}{1+1.1E_\mu \cos\theta^*/\epsilon_K} \right\}}. \quad (7)$$

The charge ratio as a function of $E_\mu \cos\theta^*$ is plotted in Fig. 8 along with the results from L3 + C [10,34], Bess-TEV [11], UTAH [33], MUTRON [22], OPERA [9], CMS [12], and the MINOS Far Detector [7,8]. In instances where the data were not already listed as a function of




FIG. 8. The atmospheric muon charge ratio as a function of $E_\mu \cos\theta^*$. The y-axis uncertainties are the statistical and systematic uncertainties added in quadrature. The MINOS Near Detector data, rebinned in five equal $\cos\theta^*$ intervals, are plotted for each bin at the median $E_\mu \cos\theta^*$ values; the x-axis uncertainties are the bin widths. The dashed line is the best-fit curve to the πK model using only MINOS Near and Far Detector data.

$E_\mu \cos\theta^*$ we convolved the published $\cos\theta$ acceptances with the given E_μ data, with assistance from the original authors in the case of L3 + C [35].

A χ^2 per degree of freedom fit test to the πK model was performed over (f_K, f_π) space using only the charge ratio measurements from the MINOS Near and Far Detectors. The χ^2 minimum was found at $f_\pi = 0.55$ and $f_K = 0.70$. These results are consistent with earlier fits by [7–9]. The best-fit curve to the data is plotted in Fig. 8 and indicates that the kaon contribution to the charge ratio becomes significant at $E_\mu \cos\theta^*$ greater than a few hundred GeV. Using two functionally identical detectors, at two different depths, we have demonstrated that the increase in charge ratio observed at the deeper Far Detector is consistent with an increase in the fraction of observed muons arising from kaon decays in the extensive air shower.

VII. SUMMARY

A charge ratio measurement has been performed on 301 days of atmospheric muon data collected using the MINOS Near Detector. The atmospheric muon charge ratio measured at 225 mwe underground is

$$\frac{N_{\mu^+}}{N_{\mu^-}} = 1.266 \pm 0.001(\text{stat})^{+0.015}_{-0.014}(\text{syst}). \quad (8)$$

The reconstructed underground energy and zenith angle for each muon used in the analysis were used to extrapolate the muon surface energy. No statistically significant change was observed in the charge ratio as a function of either the surface energy E_μ , from 62 to 960 GeV, or in the vertical muon surface energy $E_\mu \cos\theta^*$ from 57 to 134 GeV. This work presents the first demonstration of an increase in the observed muon charge ratio between functionally equivalent shallow and deep underground detectors. The most likely source of this increase is the greater probability that a muon in the deep detector data sample results from kaon production in extensive air showers. A fit has been performed to a simple parametric model of muon production from pion and kaon parents, and the results of the fit support this interpretation of the combined MINOS Near and Far Detector charge ratio measurements.

ACKNOWLEDGMENTS

This work was supported by the U.S. DOE, the United Kingdom STFC, the U.S. NSF, the State and University of Minnesota, the University of Athens, Greece, and Brazil's FAPESP and CNPq. We are grateful to the Minnesota Department of Natural Resources, the crew of Soudan Underground Laboratory, and the staff of Fermilab for their contributions to this effort.

- [1] T.K. Gaisser, *Cosmic Rays and Particle Physics* (Cambridge University Press, Cambridge, England, 1990), p. 279.
 [2] P. Lipari, *Astropart. Phys.* **1**, 195 (1993).

- [3] E. Zas, F. Halzen, and R. A. Vazquez, *Astropart. Phys.* **1**, 297 (1993).
 [4] T.K. Gaisser and M. Honda, *Annu. Rev. Nucl. Part. Sci.* **52**, 153 (2002).

- [5] M. Aglietta *et al.* (EAS-TOP and MACRO Collaborations), *Astropart. Phys.* **21**, 223 (2004).
- [6] V. A. Derbina *et al.* (RUNJOB Collaboration), *Astrophys. J.* **628**, L41 (2005).
- [7] P. Schreiner, J. Reichenbacher, and M. Goodman, *Astropart. Phys.* **32**, 61 (2009).
- [8] P. Adamson *et al.* (MINOS Collaboration), *Phys. Rev. D* **76**, 052003 (2007).
- [9] N. Agafonova *et al.* (OPERA Collaboration), *Eur. Phys. J. C* **67**, 25 (2010).
- [10] P. Achard *et al.* (L3 Collaboration), *Phys. Lett. B* **598**, 15 (2004).
- [11] S. Haino *et al.*, *Phys. Lett. B* **594**, 35 (2004).
- [12] V. Khachatryan *et al.* (CMS Collaboration), *Phys. Lett. B* **692**, 83 (2010).
- [13] T. Hebbeker and C. Timmermans, *Astropart. Phys.* **18**, 107 (2002).
- [14] D.G. Michael *et al.* (MINOS Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **596**, 190 (2008).
- [15] D.A. Crane *et al.* (NuMI Beam Group), Report No. FERMILAB-TM-1946, 1995.
- [16] N. Tagg *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **539**, 668 (2005).
- [17] A. Belias *et al.*, *IEEE Trans. Nucl. Sci.* **51**, 451 (2004).
- [18] T. Cundiff *et al.*, *IEEE Trans. Nucl. Sci.* **53**, 1347 (2006).
- [19] E. Scapparone, arXiv:physics/9902043.
- [20] S. Agostinelli *et al.* (GEANT4 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [21] R. Fruhwirth, *Nucl. Instrum. Methods Phys. Res., Sect. A* **262**, 444 (1987).
- [22] S. Matsuno *et al.*, *Phys. Rev. D* **29**, 1 (1984).
- [23] W.M. Yao *et al.* (Particle Data Group), *J. Phys. G* **33**, 1 (2006), see Chap. 24, *Cosmic Rays*, p. 245.
- [24] J. Reichenbacher and J. de Jong (MINOS Collaboration), in [36], pp. 1241–1245.
- [25] F.M. Smith, W. Birnbaum, and W.H. Barkas, *Phys. Rev.* **91**, 765 (1953).
- [26] W.H. Barkas, W. Birnbaum, and F.M. Smith, *Phys. Rev.* **101**, 778 (1956).
- [27] J.D. Jackson and R.L. McCarthy, *Phys. Rev. B* **6**, 4131 (1972).
- [28] A.R. Clark *et al.*, *Phys. Lett.* **41B**, 229 (1972).
- [29] J.D. Jackson, *Phys. Rev. D* **59**, 017301 (1998).
- [30] R.N. Lee, A.I. Milstein, I.S. Terekhov, and S.G. Karshenboim, *Phys. Rev. A* **71**, 052501 (2005).
- [31] U.S. Department of the Interior Geological Survey, 2010, map ID numbers 13040 and 13615, <http://libremap.org/data/state/illinois/drg/>.
- [32] M. Goodman (MINOS Collaboration), in [36], pp. 1225–1229.
- [33] G. K. Ashley II, J.W. Keuffel, and M.O. Larson, *Phys. Rev. D* **12**, 20 (1975).
- [34] 2010 (unpublished), independent reevaluation of the L3 + C uncertainties [7] suggests that they may have been too conservative in their systematic uncertainty assignment, thus hindering their ability to observe (in their own data set) a rise in the atmospheric muon charge ratio as a function of $E_\mu \cos\theta^*$.
- [35] M. Unger (personal communication).
- [36] *Proceedings of the 30th International Cosmic Ray Conference*, edited by R. Caballero, J.C. D’Olivo, G. Medina-Tanco, L. Nellen, F.A. Sánchez, and J.F. Valdés-Galicia (Universidad Nacional Autónoma de México, Mexico City, 2008).