

Hirschhorn's identities

Article (Published Version)

Hammond, Paul, Lewis, Richard and Liu, Zhi-Guo (1999) Hirschhorn's identities. *Bulletin of the Australian Mathematical Society*, 60 (1). pp. 73-80. ISSN 0004-9727

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/42947/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Bulletin of the Australian Mathematical Society

<http://journals.cambridge.org/BAZ>


Additional services for *Bulletin of the Australian Mathematical Society*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Hirschhorn's identities

Paul Hammond, Richard Lewis and Zhi-Guo Liu

Bulletin of the Australian Mathematical Society / Volume 60 / Issue 01 / August 1999, pp 73 - 80
DOI: 10.1017/S0004972700033347, Published online: 17 April 2009

Link to this article: http://journals.cambridge.org/abstract_S0004972700033347

How to cite this article:

Paul Hammond, Richard Lewis and Zhi-Guo Liu (1999). Hirschhorn's identities. Bulletin of the Australian Mathematical Society, 60, pp 73-80 doi:10.1017/S0004972700033347

Request Permissions : [Click here](#)

HIRSCHHORN'S IDENTITIES

PAUL HAMMOND, RICHARD LEWIS AND ZHI-GUO LIU

We prove a general identity between power series and use this identity to give proofs of a number of identities proposed by M.D. Hirschhorn. We also use the identity to give proofs of a well-known result of Jacobi, the quintuple-product identity and Winquist's identity.

1 INTRODUCTION

We suppose throughout that q is a complex number with $|q| < 1$; this condition ensures that all the sums and products that appear here converge.

In January 1998, Mike Hirschhorn conjectured (written communication) that

(H)

$$\begin{aligned} \sum_{n \in \mathbb{Z}} q^{3n^2} \sum_{n \in \mathbb{Z}} q^{3n^2+n} + q \sum_{n \in \mathbb{Z}} q^{3n^2+2n} \sum_{n \in \mathbb{Z}} q^{3n^2+3n} &= \frac{(q^3, q^6; q^6)_{\infty}^2}{(q, q^5; q^6)_{\infty}^2} \\ \sum_{n \in \mathbb{Z}} q^{5n^2} \sum_{n \in \mathbb{Z}} q^{5n^2+n} + q^2 \sum_{n \in \mathbb{Z}} q^{5n^2+4n} \sum_{n \in \mathbb{Z}} q^{5n^2+5n} &= \frac{(q^4, q^5, q^6, q^{10}; q^{10})_{\infty}^2}{(q^2, q^3, q^7, q^8; q^{10})_{\infty}^2} \\ \sum_{n \in \mathbb{Z}} q^{5n^2} \sum_{n \in \mathbb{Z}} q^{5n^2+3n} + q \sum_{n \in \mathbb{Z}} q^{5n^2+2n} \sum_{n \in \mathbb{Z}} q^{5n^2+5n} &= \frac{(q^2, q^5, q^8, q^{10}; q^{10})_{\infty}^2}{(q, q^4, q^6, q^9; q^{10})_{\infty}^2} \\ \sum_{n \in \mathbb{Z}} q^{7n^2+n} \sum_{n \in \mathbb{Z}} q^{7n^2+2n} + q^2 \sum_{n \in \mathbb{Z}} q^{7n^2+5n} \sum_{n \in \mathbb{Z}} q^{7n^2+6n} &= \frac{(q^6, q^7, q^7, q^8, q^{14}, q^{14}; q^{14})_{\infty}}{(q^2, q^3, q^5, q^9, q^{11}, q^{12}; q^{14})_{\infty}} \\ \sum_{n \in \mathbb{Z}} q^{7n^2+2n} \sum_{n \in \mathbb{Z}} q^{7n^2+3n} + q \sum_{n \in \mathbb{Z}} q^{7n^2+4n} \sum_{n \in \mathbb{Z}} q^{7n^2+5n} &= \frac{(q^2, q^7, q^7, q^{12}, q^{14}, q^{14}; q^{14})_{\infty}}{(q, q^3, q^4, q^{10}, q^{11}, q^{13}; q^{14})_{\infty}} \\ \sum_{n \in \mathbb{Z}} q^{7n^2+n} \sum_{n \in \mathbb{Z}} q^{7n^2+4n} + q \sum_{n \in \mathbb{Z}} q^{7n^2+3n} \sum_{n \in \mathbb{Z}} q^{7n^2+6n} &= \frac{(q^4, q^7, q^7, q^{10}, q^{14}, q^{14}; q^{14})_{\infty}}{(q, q^5, q^6, q^8, q^9, q^{13}; q^{14})_{\infty}} \end{aligned}$$

Received 21st December, 1998

Copyright Clearance Centre, Inc. Serial-fee code: 0004-9729/99 \$A2.00+0.00.

and asked for proofs. Here,

$$(q^{a_1}, q^{a_2}, \dots, q^{a_k}; q^n)_\infty := (q^{a_1}; q^n)_\infty (q^{a_2}; q^n)_\infty \cdots (q^{a_k}; q^n)_\infty,$$

where

$$(z; q)_\infty := \prod_{n=0}^{\infty} (1 - zq^n)$$

We also write

$$[z; q]_\infty := (z; q)_\infty (z^{-1}q; q)_\infty,$$

and

$$(1.1) \quad [z_1, z_2, \dots, z_n; q]_\infty := [z_1; q]_\infty [z_2; q]_\infty \cdots [z_n; q]_\infty$$

for $z_1, z_2, \dots, z_n \neq 0$. (The function $[z; q]_\infty$ is a close relative of the Jacobian theta functions. Indeed, with $z = e^{2\pi i x}$, $q = e^{2\pi i \tau}$, we have $\theta_1(\pi x | \tau) = iq^{1/8} z^{-1/2} (q; q)_\infty [z; q]_\infty$, see [10, p.470]). With this notation, note that, as a function of z ,

$$(1.2) \quad [z; q]_\infty \text{ has simple zeros at } z = q^k, \text{ for } k \in \mathbb{Z},$$

an essential singularity at $z = 0$ and no other zeros and no poles. We have the simple and easily verifiable properties:

$$(1.3) \quad [z^{-1}; q]_\infty = -z^{-1} [z; q]_\infty = [zq; q]_\infty$$

$$(1.4) \quad [z, zq, \dots, zq^{k-1}; q^k]_\infty = [z; q]_\infty$$

$$(1.5) \quad [z, \zeta z, \dots, \zeta^{k-1} z; q]_\infty = [z^k; q^k]_\infty$$

where $\zeta = e^{2\pi i/k}$, and we note that it follows from (1.3) that

$$(1.6) \quad [z^{-1}q; q]_\infty = [z; q]_\infty.$$

Jacobi's well-known triple product identity (see [4, Section 19.8.1] or [6] for an analytic, respectively, combinatorial proof) states that

$$(1.7) \quad (q; q)_\infty [z; q]_\infty = \sum_{n \in \mathbb{Z}} z^n q^{n(n-1)/2}.$$

With this notation (1.1), it may be readily checked that all the identities (H) are

instances of:

$$\begin{aligned} \sum_{n \in \mathbb{Z}} q^{Mn^2 + (x-t)n} \sum_{n \in \mathbb{Z}} q^{Mn(n-1) + (x+t)n} + q^t \sum_{n \in \mathbb{Z}} q^{Mn^2 + (x+t)n} \sum_{n \in \mathbb{Z}} q^{Mn(n-1) + (x-t)n} \\ = \frac{[q^{2t}, q^{2x}, q^M, q^{2M}]_{\infty} (q^{2M}; q^{2M})_{\infty}}{[q^t, q^{M+t}, q^x, q^{M+x}, q^{2M}]_{\infty}} \end{aligned}$$

which, by (1.4) and (1.5),

$$= [-q^t, -q^x; q^M]_{\infty} (q^M; q^M)_{\infty}^2$$

and this, by (1.7),

$$(1.8) \quad = \sum_{n \in \mathbb{Z}} q^{Mn(n-1)/2 + tn} \sum_{n \in \mathbb{Z}} q^{Mn(n-1)/2 + xn}$$

and it is the general identity (1.8) that we prove in Theorem 2 below.

2 THE PROOF OF THE IDENTITIES

Suppose $q \in \mathbb{C} \setminus \{0\}$ (with $|q| < 1$). Following [2], say that $z_0, w_0 \in \mathbb{C} \setminus \{0\}$ are equivalent if $z_0 = q^n w_0$, for some $n \in \mathbb{Z}$.

LEMMA. Suppose $f : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$ is analytic save for some isolated simple poles and suppose that f satisfies

$$(2.1) \quad f(qz) = q^{-1} f(z),$$

for $z \neq 0$. If \mathcal{P} is a complete set of inequivalent poles of f (\mathcal{P} is necessarily finite), then

$$(2.2) \quad \sum_{z \in \mathcal{P}} \text{res}(f; z) = 0.$$

PROOF: First note that, if $f : \mathbb{C} \setminus \{0\} \rightarrow \mathbb{C}$ satisfies (2.1) and if z_0 and w_0 are equivalent points, $z_0 = q^n w_0$, say, then w_0 is a simple pole of f if and only if z_0 is a simple pole of f and

$$\begin{aligned} \text{res}(f; z_0) &= \lim_{z \rightarrow z_0} (z - z_0) f(z) \\ &= \lim_{w \rightarrow w_0} (q^n w - q^n w_0) f(q^n w) \\ &= \lim_{w \rightarrow w_0} q^n (w - w_0) q^{-n} f(w) \\ &= \lim_{w \rightarrow w_0} (w - w_0) f(w) = \text{res}(f; w_0). \end{aligned}$$

Thus the choice of \mathcal{P} is immaterial in evaluating the sum at (2.2). Now let $\mathcal{A} := \{z : r|q| \leq |z| < r\}$ ($r > 0$) be a half-open annulus in \mathbb{C} , centred at the origin. So \mathcal{A} includes its inner boundary circle, $S_2 = \{w : |w| = r|q|\}$, but not its outer boundary circle $S_1 = \{z : |z| = r\}$. The non-zero poles of f are isolated and so we can, changing r , if necessary, suppose that neither S_1 nor S_2 contains any pole or zero of f . It is plain that every $z \in \mathbb{C} \setminus \{0\}$ is equivalent to a unique point in \mathcal{A} and that distinct points in \mathcal{A} are inequivalent and so we may take \mathcal{P} to be the set of poles of f in \mathcal{A} . Now

$$\sum_{z \in \mathcal{P}} \operatorname{res}(f; z) = \frac{1}{2\pi i} \int_{S_1} f(z) dz - \frac{1}{2\pi i} \int_{S_2} f(w) dw.$$

But if $w \in S_2$, then $w = qz$, where $z \in S_1$. We have $dw = qdz$ and $f(w) = f(qz) = q^{-1}f(z)$ and so

$$\sum_{z \in \mathcal{P}} \operatorname{res}(f; z) = \frac{1}{2\pi i} \int_{S_1} f(z) dz - \frac{1}{2\pi i} \int_{S_1} q^{-1}f(z) q dz = 0. \quad \square$$

(This Lemma is really a disguised form of a well-known property of elliptic functions; namely, that the sum of the residues of an elliptic function at its poles in a period parallelogram is zero.)

THEOREM 1. *Suppose $a_1, a_2, \dots, a_n; b_1, b_2, \dots, b_n \in \mathbb{C} \setminus \{0\}$ satisfy:*

- (i) a_i and a_j are inequivalent, for $i \neq j$.
- (ii) $a_1 a_2 \dots a_n = b_1 b_2 \dots b_n$. Then

$$(2.3) \quad \sum_{i=1}^n \frac{\prod_{j=1}^n [a_i b_j^{-1}; q]_{\infty}}{\prod_{j=1, j \neq i}^n [a_i a_j^{-1}; q]_{\infty}} = 0.$$

PROOF: Define

$$\mathcal{F}(z) = \mathcal{F}(z; q) := \frac{\prod_{j=1}^n [z b_j^{-1}; q]_{\infty}}{z \prod_{j=1}^n [z a_j^{-1}; q]_{\infty}}.$$

Condition (i) shows that \mathcal{F} has only simple poles in $\mathbb{C} \setminus \{0\}$ and that $\{a_1, a_2, \dots, a_n\}$ is a complete set of inequivalent poles. Taking into account (1.3), condition (ii) shows that (2.1) is satisfied. Now the residue of \mathcal{F} at a_i is

$$\operatorname{res}(\mathcal{F}; a_i) = \lim_{z \rightarrow a_i} (z - a_i) \mathcal{F}(z)$$

$$\begin{aligned}
&= \frac{\prod_{j=1}^n [a_i b_j^{-1}; q]_{\infty}}{a_i \prod_{j=1, j \neq i}^n [a_i a_j^{-1}; q]_{\infty}} \lim_{z \rightarrow a_i} \frac{(z - a_i)}{[z a_i^{-1}; q]_{\infty}} \\
&= \frac{\prod_{j=1}^n [a_i b_j^{-1}; q]_{\infty}}{a_i \prod_{j=1, j \neq i}^n [a_i a_j^{-1}; q]_{\infty}} \frac{1}{(q; q)_{\infty}^2} \lim_{z \rightarrow a_i} \frac{(z - a_i)}{(1 - z a_i^{-1})} \\
&= -\frac{1}{(q; q)_{\infty}^2} \frac{\prod_{j=1}^n [a_i b_j^{-1}; q]_{\infty}}{\prod_{j=1, j \neq i}^n [a_i a_j^{-1}; q]_{\infty}}.
\end{aligned}$$

(2.3) now follows from (2.2). \square

Theorem 1 appears as [8, (7.4.3)], though no proof is given. The author states that it is a consequence of the 'general theorem' on (Weierstrass) sigma functions and refers to [10, p.451, exercise 3], where again no proof is given. The proof given here appeared in [7]. We can now prove

THEOREM 2. *If $x, t, M \in \mathbb{Z}$ and $0 < t < M$, then*

$$\begin{aligned}
(2.4) \quad \sum_{n \in \mathbb{Z}} q^{Mn^2 + (x-t)n} \sum_{n \in \mathbb{Z}} q^{Mn(n-1) + (x+t)n} + q^t \sum_{n \in \mathbb{Z}} q^{Mn^2 + (x+t)n} \sum_{n \in \mathbb{Z}} q^{Mn(n-1) + (x-t)n} \\
= \sum_{n \in \mathbb{Z}} q^{Mn(n-1)/2 + xn} \sum_{n \in \mathbb{Z}} q^{Mn(n-1)/2 + tn}.
\end{aligned}$$

PROOF: Take $n = 3$ in Theorem 1 and q^{2M} for q and set

$$(a_1, a_2, a_3; b_1, b_2, b_3) = (1, q^{2t}, q^{M+t}, -q^{x+t}, -q^{M-x+t}, q^t).$$

The hypotheses of Theorem 1 are satisfied and (2.3) gives

$$\begin{aligned}
\frac{[-q^{-(x+t)}, -q^{-(M-x+t)}, q^{-t}; q^{2M}]_{\infty}}{[q^{-2t}, q^{-(M+t)}; q^{2M}]_{\infty}} + \frac{[-q^{-(x-t)}, -q^{-(M-x-t)}, q^t; q^{2M}]_{\infty}}{[q^{2t}, q^{-(M-t)}; q^{2M}]_{\infty}} \\
+ \frac{[-q^{M-x}, -q^x, q^M; q^{2M}]_{\infty}}{[q^{(M+t)}, q^{(M-t)}; q^{2M}]_{\infty}} = 0.
\end{aligned}$$

We use (1.3) to write this as

$$\begin{aligned}
-\frac{[-q^{x+t}, -q^{M-x+t}, q^t; q^{2M}]_{\infty}}{[q^{2t}, q^{M+t}; q^{2M}]_{\infty}} - q^t \frac{[-q^{x-t}, -q^{M-x-t}, q^t; q^{2M}]_{\infty}}{[q^{2t}, q^{M-t}; q^{2M}]_{\infty}} \\
+ \frac{[-q^{M-x}, -q^x, q^M; q^{2M}]_{\infty}}{[q^{(M+t)}, q^{(M-t)}; q^{2M}]_{\infty}} = 0,
\end{aligned}$$

or

$$[-q^{x+t}, -q^{M-x+t}; q^{2M}]_{\infty} + q^t[-q^{x-t}, -q^{M-x-t}; q^{2M}]_{\infty} = \frac{[-q^x, -q^{M+x}, q^{2t}, q^M; q^{2M}]_{\infty}}{[q^t, q^{M+t}; q^{2M}]_{\infty}},$$

since, by (1.6), $[q^{M-x}; q^{2M}]_{\infty} = [q^{M+x}; q^{2M}]_{\infty}$. Now multiply each side by $(q^{2M}; q^{2M})_{\infty}^2$ and use (1.4) and (1.5) to get

$$\begin{aligned} [-q^{x+t}, -q^{M-x+t}; q^{2M}]_{\infty} (q^{2M}; q^{2M})_{\infty}^2 + q^t[-q^{x-t}, -q^{M-x-t}; q^{2M}]_{\infty} (q^{2M}; q^{2M})_{\infty}^2 \\ = [-q^x; q^M]_{\infty} [-q^t; q^M]_{\infty} (q^M; q^M)_{\infty}^2 \end{aligned}$$

which, as we pointed out earlier, translates by way of (1.7) into (2.4). \square

Now (2.4) appears in Hirschhorn's work (it is [5, (2.1)] after applying the triple product identity), so he had already proved the identities (H) he was asking about as well as their generalisation (1.8). His proof of (2.4) is very different from that given here. He multiplies together two copies of the triple product identity and sorts the resulting equation into even and odd powers of q . (2.4) also appeared in [3] with more or less the same (that is, Hirschhorn's) proof.

3 THREE MORE IDENTITIES

Theorem 1 can plainly be used to generate a large number of identities of the same type as (2.4). (infinitely many, of course). We shall give three more applications of Theorem 1.

1. Our first application is another of Jacobi's marvellous identities. Taking $n = 3$ and q^2 in place of q and with $(a_1, a_2, a_3; b_1, b_2, b_3) = (1, -q, -q^2; q, q, q)$, Theorem 1 gives

$$\frac{[q^{-1}; q^2]_{\infty}^3}{[-q^{-1}, -q^{-2}; q^2]_{\infty}} + \frac{[-1; q^2]_{\infty}^3}{[-q, q^{-1}; q^2]_{\infty}} + \frac{[-q; q^2]_{\infty}^3}{[-q^2, q; q^2]_{\infty}} = 0$$

from which, with (1.3) and (1.6), we get

$$(3.1) \quad -[q; q^2]_{\infty}^4 - q[-1; q^2]_{\infty}^4 + [-q; q^2]_{\infty}^4 = 0.$$

Noting that $[-1; q^2]_{\infty} = 2(-q^2; q^2)_{\infty}^2$, this translates into Jacobi's identity

$$\left\{ \prod_{n=1}^{\infty} (1 - q^{2n-1}) \right\}^8 + 16q \left\{ \prod_{n=1}^{\infty} (1 + q^{2n}) \right\}^8 = \left\{ \prod_{n=1}^{\infty} (1 + q^{2n-1}) \right\}^8.$$

(According to [10, p.470, footnote], Jacobi said of this identity that it was an 'aequatio identica satis abstrusa'.)

2. We now use Theorem 1 to derive the quintuple product identity [2], which in one form states:

$$(3.2) \quad \frac{[z^2; q]_\infty}{[z; q]_\infty} (q; q)_\infty = \{[z^3 q; q^3]_\infty + z[z^{-3} q; q^3]_\infty\} (q^3; q^3)_\infty.$$

If we set $n = 3$, $\omega = e^{2\pi i/3}$ and $(a_1, a_2, a_3; b_1, b_2, b_3) = (z, z^{-1}, q; q^{1/3}, q^{1/3}\omega, q^{1/3}\omega^{-1})$ in Theorem 1, (2.3) becomes

$$\begin{aligned} & \frac{[zq^{-1/3}, zq^{-1/3}\omega^{-1}, zq^{-1/3}\omega; q]_\infty}{[z^2, zq^{-1}; q]_\infty} + \frac{[z^{-1}q^{-1/3}, z^{-1}q^{-1/3}\omega^{-1}, z^{-1}q^{-1/3}\omega; q]_\infty}{[z^{-2}, z^{-1}q^{-1}; q]_\infty} \\ & + \frac{[q^{2/3}, q^{2/3}\omega^{-1}, q^{2/3}\omega; q]_\infty}{[z^{-1}q, zq; q]_\infty} = 0, \end{aligned}$$

which, with the help of (1.3) and (1.5), gives (3.2).

3. Finally, Winquist's Identity [9], namely:

$$\begin{aligned} & [x; q]_\infty [y; q]_\infty [xy^{-1}; q]_\infty [xy; q]_\infty (q; q)_\infty^2 = xy^{-1} [y^3; q^3]_\infty \\ & \left\{ x[qx^{-3}; q^3]_\infty - [qx^3; q^3]_\infty \right\} (q^3; q^3)_\infty^2 - [x^3; q^3]_\infty \left\{ y[qy^{-3}; q^3]_\infty - [qy^3; q^3]_\infty \right\} (q^3; q^3)_\infty^2, \end{aligned}$$

also follows from Theorem 1, where now we take $n = 6$ and

$$\begin{aligned} & (a_1, a_2, a_3, a_4, a_5, a_6; b_1, b_2, b_3, b_4, b_5, b_6) \\ & = \left(x, x^{-1}, y, y^{-1}, \omega, \omega^{-1}; q^{1/3}, q^{1/3}\omega, q^{1/3}\omega^{-1}, -1, -q^{-1/2}, q^{-1/2} \right). \end{aligned}$$

We omit the details.

4 CONCLUSION

We have seen that Theorem 1 is very powerful and has many applications. A variant of this Theorem was used by Atkin and Swinnerton-Dyer to prove the Dyson conjectures on the ranks of partitions [2] and, in [7], the Theorem was again used to prove some other theorems about ranks. For these reasons, it would be very interesting to have a bijective/combinatorial proof of Theorem 1; such a proof might well lead to a bijective/combinatorial proof of the rank theorems, whose statements are, after all, combinatorial in nature. But even the formulation of a such a version of Theorem 1 seems fraught with difficulties.

Note that, in Theorem 1, the parameters a_1, a_2, \dots, a_n are required to be inequivalent. This condition ensures that the function \mathcal{F} defined in the proof of Theorem 1

has only simple poles, but what would we get if some of the a_i were equivalent and so then the function \mathcal{F} has poles of order ≥ 2 ? Now the function

$$\mathcal{G}(z) = \frac{[az, bz, cz; q]_{\infty}}{z[z, z, abc; q]_{\infty}}$$

satisfies (2.1) and so

$$(4.1) \quad \text{res}(\mathcal{G}; 1) + \text{res}(\mathcal{G}; (abc)^{-1}) = 0.$$

It turns out that (4.1) leads to a striking identity relating a theta product and a sum of Lambert series from which many theorems on sums of squares may be deduced. A paper on this identity is in preparation [1].

REFERENCES

- [1] G.E. Andrews, R.P. Lewis and Z-G. Liu, 'An Identity relating a theta function and a sum of Lambert series', (in preparation).
- [2] A.O.L. Atkin and P. Swinnerton-Dyer, 'Some properties of partitions', *Proc. London Math. Soc.* (3) 4 (1954), 84–106.
- [3] J.A. Ewell, 'Arithmetical consequences of a sextuple product identity', *Rocky Mountain J. Math.* 25 (1995), 1287–1293.
- [4] G.H. Hardy and E.M. Wright, *The theory of numbers*, (3rd edition (Oxford University Press, 1954).
- [5] M.D. Hirschhorn, 'A simple proof of Jacobi's four-square theorem', *J. Austral. Math. Soc. Ser. A* 32 (1982), 61–67.
- [6] R.P. Lewis, 'A combinatorial proof of the triple product identity', *Amer. Math. Monthly* 91 (1984), 420–423.
- [7] R.P. Lewis, 'On the ranks of partitions modulo 9', *Bull. London Math. Soc.* 23 (1991), 417–421.
- [8] L.J. Slater, *Generalized hypergeometric functions* (Cambridge University Press, Cambridge, U.K., 1966).
- [9] L. Winquist, 'An elementary proof of $p(11m + 6) \equiv 0 \pmod{11}$ ', *J. Comb. Theory* 6 (1969), 56–59.
- [10] E.T. Whittaker and G.N. Watson, *A course of modern analysis*, (4th edition) (Cambridge University Press, New York, 1950).

SMS

The University of Sussex
Brighton BN1 9QH
United Kingdom