Global effects of land use on local terrestrial biodiversity

Article (Accepted Version)

Newbold, Tim, Hudson, Lawrence N, Hill, Samantha L L, Contu, Sara, Lysenko, Igor, Senior, Rebecca A, Börger, Luca, Bennett, Dominic J, Choimes, Argyrios, Collen, Ben, Day, Julie, De Palma, Adriana, Díaz, Sandra, Echeverria-Londoño, Susy, Edgar, Melanie J et al. (2015) Global effects of land use on local terrestrial biodiversity. *Nature*, 520 (7545). pp. 45-50. ISSN 0028-0836

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/53593/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

The full, formatted version of this paper can be found at the Nature website:
<http://www.nature.com/nature/journal/v520/n7545/full/nature14324.html>.

To cite this paper: Newbold T., Hudson L.N., Hill S.L.L., Contu S., Lysenko I., Senior R.A., Börger L., Bennett D.J., Choimes A., Collen B., Day J., De Palma A., Díaz S., Echeverria-Londoño S., Edgar M.J., Feldman A., Garon M., Harrison M.L.K., Alhusseini T., Ingram D.J., Itescu Y., Kattge J., Kemp V., Kirkpatrick L., Kleyer M., Laginha Pinto Correia D., Martin C.D., Meiri S., Novosolov M., Pan Y., Phillips H.R.P., Purves D.W., Robinson A., Simpson J., Tuck S.L., Weiher E., White H.J., Ewers R.M., Mace G.M., Scharlemann J.P.W., & Purvis A. (2015) Global effects of land use on local terrestrial biodiversity. *Nature*, **520**, 45–50.

DOI: 10.1038/nature14324

Global effects of land use on local terrestrial biodiversity

Tim Newbold^{1,2*}, Lawrence N. Hudson^{3*}, Samantha L.L. Hill^{1,3}, Sara Contu³, Igor Lysenko⁴, Rebecca A. Senior¹, Luca Börger⁵, Dominic Bennett^{4,†}, Argyrios Choimes^{3,4}, Ben Collen⁶, Julie Day^{4,‡}, Adriana De Palma^{3,4}, Sandra Díaz⁷, Susy Echeverria-Londoño³, Melanie Edgar³, Anat Feldman⁸, Morgan Garon⁴, Michelle L. K. Harrison⁴, Tamera Alhusseini⁴, Daniel J. Ingram^{4,††}, Yuval Itescu⁸, Jens Kattge^{9,10}, Victoria Kemp⁴, Lucinda Kirkpatrick^{4,‡‡}, Michael Kleyer¹¹, David Laginha Pinto Correia³, Callum Martin⁴, Shai Meiri⁸, Maria Novosolov⁸, Yuan Pan⁴, Helen R.P. Phillips^{3,4}, Drew W. Purves², Alexandra Robinson⁴, Jake Simpson⁴, Sean Tuck¹², Evan Weiher¹³, Hannah J. White^{4,†††}, Robert M. Ewers⁴, Georgina M. Mace⁶, Jörn P.W. Scharlemann^{1,14}, Andy Purvis^{3,4}

¹United Nations Environment Programme World Conservation Monitoring Centre, 219 Huntingdon Road, Cambridge CB3 0DL, UK.

²Computational Science Laboratory, Microsoft Research Cambridge, 21 Station Road, Cambridge CB1 2FB, UK.

³Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK.

⁴Department of Life Sciences, Imperial College London, Silwood Park, SL5 7PY, UK.

⁵Department of Biosciences, College of Science, Swansea University, Singleton Park, Swansea SA2 8PP, UK.

⁶Department of Genetics, Evolution and Environment, Centre for Biodiversity and Environment Research, University College London, Gower Street, London WC1E 6BT, UK.

⁷Instituto Multidisciplinario de Biología Vegetal (CONICET-UNC) and FCEfYN, Universidad Nacional de Córdoba, Casilla de Correo 495, 5000 Córdoba, Argentina.

⁸Department of Zoology, Faculty of Life Sciences, Tel-Aviv University, 6997801 Tel Aviv, Israel.

⁹Max Planck Institute for Biogeochemistry, Hans Knöll Straße 10, 07743 Jena, Germany.

¹⁰German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, 04103 Leipzig, Germany.

¹¹Landscape Ecology Group, Institute of Biology and Environmental Sciences, University of Oldenburg, D-26111 Oldenburg, Germany.

¹²Department of Plant Sciences, University of Oxford, Oxford, OX1 3RB, UK.

¹³Biology Department, University of Wisconsin – Eau Claire, Eau Claire, Wisconsin 54701, United States.

¹⁴School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK.

[†]Current address: Department of Earth Science and Engineering, Imperial College London, London SW7 2AZ, UK and Institute of Zoology, Zoological Society of London, London, UK.

[‡]Current address: College of Life and Environmental Sciences, Hatherly Laboratories, University of Exeter, Prince of Wales Road, Exeter, EX4 4PS, UK.

^{††}Current address: School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK.

^{‡‡}Current address: School of Biological and Ecological Sciences, University of Stirling, Stirling FK9 4LA, UK.

^{†††}Current address: School of Biological Sciences, Queen's University Belfast, 97 Lisburn Road, Belfast BT9 7BL, UK.

*These authors contributed equally to this work.

Human activities, especially conversion and degradation of habitats, are causing global biodiversity declines. How local ecological assemblages are responding is less clear – a concern given their importance for many ecosystem functions and services. We analyze a terrestrial assemblage database of unprecedented geographic and taxonomic coverage to quantify local biodiversity responses to land-use and related changes. In the worst-impacted habitats, these pressures reduce within-sample species richness by an average of 76.5%, total abundance by 39.5% and rarefaction-based richness by 40.3%. We estimate that, globally, these pressures have already slightly reduced average within-sample richness (by 13.6%), total abundance (10.7%) and rarefaction-based richness (8.1%), with changes showing marked spatial variation. Rapid further losses are predicted under a business-as-usual land-use scenario: within-sample richness is projected to fall by a further 3.4% globally by 2100, with losses concentrated in biodiverse but economically poor countries. Strong mitigation can deliver much more positive biodiversity changes (up to a 1.9% average increase) that are less strongly related to countries' socioeconomic status.

Biodiversity faces growing pressures from human actions, including habitat conversion and degradation, habitat fragmentation, climate change, harvesting and pollution¹. As a result, global assessments show that species' extinction risk is increasing on average while population sizes are declining^{1,2}. Such assessments have usually focused on data-rich vertebrates, so might not reflect broader biodiversity³. Furthermore, most have concentrated on the global status of species, whereas the long-term security of many ecosystem functions and services – especially in changing environments – is likely to depend upon local biodiversity^{4–6}. Average trends in local diversity remain unclear: analyses of temporal changes in assemblages have suggested no systematic change in species richness^{7,8}, but the available times-series data might under-represent transitions between land-use types⁹ and population time series suggest vertebrate populations have declined sharply in recent decades³.

Spatial comparisons provide an alternative source of evidence on how human pressures affect biodiversity, assuming that differences in pressures have caused observed biodiversity differences between otherwise matched sites^{10–12}. The prevalence of published spatial comparisons makes it possible to go beyond particular taxa or regions^{11,12} to develop global, taxonomically representative models. Furthermore, the willingness of many researchers to share their raw data makes it possible to consider multiple aspects of biodiversity, rather than the single, simple metrics of most existing models¹⁰ which cannot capture all key aspects of diversity¹³.

We present the most geographically and taxonomically representative models to date of how several aspects of the composition and diversity of terrestrial assemblages respond to

multiple human pressures. The explanatory variables in our models most directly measure land use and infrastructure, but might correlate^{14,15} with two other important pressures – harvesting and invasive species – for which comparable high-resolution spatial data are unavailable globally. We exclude climate change effects because they are not captured well by spatial comparisons. We use our models to infer past net changes in assemblages since 1500, project future changes over this century under different socioeconomic scenarios of land use, and relate projected national changes in local biodiversity to socioeconomic variables and natural biodiversity.

Our models of local within-sample species richness (hereafter ‘richness’), rarefaction-based species richness (hereafter ‘rarefied richness’), total abundance, compositional turnover and average organism size are based on among-site comparisons of ecological assemblage composition collated from the literature as part of the PREDICTS project¹⁶. The dataset consisted of 1,130,251 records of abundance and 320,924 of occurrence or species richness at 11,525 sites (2–360 sites per study, median 15; Figure 1a). These data, from 284 publications (see Methods), represent 26,953 species – 1.4% of the number formally described¹⁷ – and 13 of the 14 terrestrial biomes (Extended Data Figure 1). Each site was scored for six putative pressures: land use¹¹ and use intensity¹⁸, land-use history¹⁹, human population density²⁰, proximity to roads²¹, and accessibility from the nearest large town. Random effects in our models accounted for study-level differences in response variables and sampling methods, and for the within-study spatial arrangement of sites.

Effects of pressures on site-level diversity

Local richness, rarefied richness and total abundance were most strongly influenced by land use and land-use intensity: they were substantially lower in most other land-use types than in primary vegetation, especially in intensively-used areas (Figure 1; see Supplementary Information for statistics and coefficient estimates). These results extend those of previous, geographically or taxonomically restricted, meta-analyses (e.g. refs. 11,22). Other variables were weaker as main effects, but showed stronger effects in interaction (Extended Data Figure 2) and were often significant overall (see Supplementary Information). Richness and total abundance tended to be slightly lower at the highest human population densities, and richness was lower nearer to roads and in more accessible sites (Figure 1). Differences in richness were not driven solely by differences in abundance: rarefied richness²³ (see Methods for details) showed weaker but mostly similar patterns, although the effects of variables other than land use and land-use intensity were not significant (Extended Data Figure 3a,b). Under the worst combinations of pressures, our models estimated richness, rarefied richness and total abundance to be 76.5%, 40.3% and 39.5% lower, respectively, than in minimally impacted sites. Effects of pressures on vertebrate, invertebrate and plant richness were statistically indistinguishable ($P > 0.05$; results not shown). The modelled coefficients were robust to efforts to correct for publication bias (Extended Data Figure 4). As with all studies based on data from the literature, unpublished data are almost unrepresented. Coefficients were also robust under cross-validation (Extended Data 3c,d), and the model residuals showed little spatial autocorrelation (Extended Data Figure 5).

The importance of secondary vegetation for conservation is hotly debated^{11,24,25}, and is crucial because this land-use will soon become the most widespread type²⁶. We find that the answer depends strongly on the secondary vegetation’s maturity: early-stage communities tend to be less diverse than those in primary vegetation and are compositionally distinct, but these differences are much reduced in mature secondary vegetation (Figures 1 & 2; we caution though that not all data sources clearly distinguished mature secondary from primary

vegetation). This successional rise in diversity accords with a recent meta-analysis of plant communities over time⁷.

Net changes in diversity provide an incomplete view of the effects of human activities on biodiversity because they ignore replacement of original species by newcomers⁸. We therefore analysed how land use affects similarity in species composition between sites. Communities under the same land use were, unsurprisingly, the most similar (Figure 2a). Across land uses, communities in primary vegetation were most like those in secondary vegetation, while plantation forest, pasture and cropland communities formed a different, human-dominated, cluster (Figure 2b).

Anthropogenic pressures can affect ecosystem functions and services more strongly than changes in species diversity would imply, if species' responses depend on their traits²⁷. Large size is often linked to species' declines^{28,29} and is important for some ecosystem processes³⁰. We combined abundance data with species' average sizes to calculate site-level community-weighted mean plant height and animal mass. As in local studies²⁹, mean plant height was lower in human-dominated land uses than in primary and secondary vegetation, and tended to decline with increasing human population density (Figure 1d). Most field studies focused on particular plant taxa, so this difference does not simply reflect tree removal. Average animal mass did not change consistently with land use or human population density, but increased with proximity to roads (Figure 1d).

Models like ours that substitute space for time ignore time lags in biotic changes, which can be important³¹. We also assume that land uses are situated randomly within studies relative to sites' intrinsic suitability for biodiversity. Adding global data on other important pressures as they become available, and also incorporating climate change, will give a more complete picture of human impacts on local biodiversity.

Global effects on local diversity to date

By applying our model for within-sample species richness – the most widely used and understood biodiversity measure – to maps of current pressure variables, we estimate the global pattern of net local changes to date in plot-level richness (Figure 3; we did not estimate total richness within the $0.5^\circ \times 0.5^\circ$ grid cells). Human-dominated areas are inferred to have lost much more local diversity than have regions where more natural vegetation remains. The worst-affected cells showed a 31% reduction in average local richness – probably enough to impact ecosystem functioning substantially⁵. Local richness increased in 1.7% of cells (by $\leq 4.8\%$). Total abundance and rarefied richness showed broadly similar patterns, although less pronounced in the latter case (Extended Data Figure 6).

We applied our models to global spatial estimates of how land use and human population changed from 1500-2005²⁶ (see Methods) to infer the global history of local biodiversity change. We focus on within-sample species richness because of its wide use and easy interpretation. Our inferences incorporate uncertainty in model parameter estimates, but not in the trajectories of the drivers themselves (which have not been assessed³²) nor effects of changes in roads and accessibility, for which temporal estimates could not be obtained.

Richness is estimated to have declined most rapidly in the 19th and 20th centuries (Figure 4), with other metrics showing similar responses (Extended Data Figure 6). By 2005, we estimate that land use and related pressures had reduced local richness by an average of 13.6% (95% CI: 9.1 – 17.8%) and total abundance by 10.7% (95% CI: 3.8% gain – 23.7% reduction) compared with what they would have been in the absence of human effects. Approximately 60% of the decline in richness was independent of effects on abundance: average rarefied richness has fallen by 8.1% (95% CI: 3.5 – 12.9%). Although these

confidence limits omit uncertainty in the projections of land use and other pressures, there is less uncertainty in estimates of current pressure levels than in changes over time³³.

Our inferences contrast with two recent analyses of community time series^{7,8}, which suggested no overall trend in local diversity, and with the Living Planet Index³ which, based on vertebrate population time series, reports a much more rapid decline in abundance than we infer. Although time series potentially provide a more direct view of temporal trends than our space-for-time approach, the available data might under-represent transitions between land-use types⁹. On the other hand, our approach may underestimate additions of species through climate change and species invasion (although accessibility and proximity to roads may partly capture the latter^{14,15}).

Global and national projections to 2095

Global changes in local diversity from 2005 to 2095 were projected using estimated land use and human population from the four Intergovernmental Panel on Climate Change Representative Concentration Pathway (RCP) scenarios²⁶, which correspond to different intensities of global climate change (Table 1). Although these estimates have limitations³², they are the most consistent available, are widely-used³⁴, and are consistent with the historical estimates²⁶. However, they – like all other global land-use projections – include no estimate of uncertainty; therefore, each of our projections must be viewed as the predicted biodiversity outcome under one particular set of land-use assumptions.

Projected net changes in average local diversity to 2095 vary widely among scenarios (Figure 4; Extended Data Figure 6). The scenario with the least climate change (IMAGE 2.6) yields the second-worst outcome for biodiversity, because it assumes rapid conversion of primary vegetation – especially in the tropics – to crops and biofuels²⁶ (Table 1, Extended Data Figure 7). These projections do not imply that low-emission scenarios must entail large losses of biodiversity, but instead reflect that scenario's mitigation strategy. Indeed, in MiniCAM 4.5 (where mitigation is through carbon markets, crop improvements and diet shifts: Table 1), average richness is projected to increase (though other diversity metrics respond more weakly: Extended Data Figure 6). The worst biodiversity outcomes arise from the scenario with most climate change – MESSAGE 8.5 – in which rapid human population growth drives widespread agricultural expansion (Table 1; Extended Data Figure 7). This scenario, which has been characterised as 'business-as-usual'³⁵, most closely matches recent trends in emissions³⁶ and gives the worst outcomes even though our projections omit direct climate impacts on local assemblages.

The global projections hide wide regional and national variation (Figure 5; Extended Data Figure 8). Projections for 2095 under 'business-as-usual' (MESSAGE 8.5) are strongly inequitable, presenting serious challenges for both sustainable development and global conservation of biodiversity (Figure 5a). Under this scenario, European and North American countries – typically with a high Human Development Index (HDI), low native biodiversity and widespread historical land conversion – are mostly projected to gain in local richness by 2095. More naturally biodiverse but less economically developed Southeast Asian and especially sub-Saharan African countries, with more natural and semi-natural habitat, will suffer the greatest losses (Figure 5a; Extended Data Figure 8f).

Such globally inequitable outcomes might be avoidable: the best scenario for biodiversity (MiniCAM 4.5; Figure 4) yielded country-level outcomes that are relatively independent of HDI, native species richness (Figure 5b) and past changes (Extended Data Figure 8e). For local richness, outcomes under MiniCAM4.5 were better than MESSAGE 8.5 for 93% of countries worldwide (Figure 5c).

Under AIM 6.0, most Afrotropical countries are projected to gain in local richness but heavy losses are inferred for the Indo-Malay region (Extended Data Figure 8). Projections

under IMAGE 2.6 are spatially similar to those under MESSAGE 8.5. The land-use change caused by the biofuels-based strategy in IMAGE 2.6 is projected to have a major negative impact overall on terrestrial biodiversity (Extended Data Figure 8).

Conclusions

Many assessments of the state of biodiversity have focused on global metrics such as rates of species extinction³⁷, but resilient delivery of ecosystem functions and services is more likely to depend on local diversity^{4–6}. Our models suggest land-use changes and associated pressures strongly reduce local terrestrial biodiversity, and we estimate global average reductions to date of 13.6% in within-sample species richness, 10.7% in total abundance and 8.1% in rarefaction-based species richness (Figs. 3 and 4). Climate change, which we could not include in our framework, is likely to exacerbate losses, especially under business-as-usual³⁸, although direct effects of climate change will increase local diversity in some regions⁸.

It is important to remember that the habitat conversion and associated changes that reduced local biodiversity had largely positive consequences for people: agricultural intensification underpinned many countries' development. However, benefits have not been shared equally among or within countries³⁹. Losses of local species richness exceeding 20% are likely to substantially impair the contribution of biodiversity to ecosystem function and services, and thus to human well-being⁵. We estimate that reductions in average plot-level species richness currently exceed this level for 28.4% of grid cells, increasing to 41.5% of cells by 2095 under 'business-as-usual' (note that we do not estimate or project total richness across the cell). Importantly, our projections suggest that such widespread large losses are not inevitable. With concerted action and the right societal choices, global sustainability of local biodiversity may be an achievable goal.

References

1. Tittensor, D. P. *et al.* A mid-term analysis of progress toward international biodiversity targets. *Science* **346**, 241–244 (2014).
2. Pimm, S. L. *et al.* The biodiversity of species and their rates of extinction, distribution, and protection. *Science* **344**, 1246752–1246752 (2014).
3. Collen, B. *et al.* Monitoring change in vertebrate abundance: the Living Planet Index. *Conserv. Biol.* **23**, 317–327 (2009).
4. Isbell, F. *et al.* High plant diversity is needed to maintain ecosystem services. *Nature* **477**, 199–202 (2011).
5. Hooper, D. U. *et al.* A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature* **486**, 105–108 (2012).
6. Cardinale, B. J. *et al.* Biodiversity loss and its impact on humanity. *Nature* **486**, 59–67 (2012).
7. Vellend, M. *et al.* Global meta-analysis reveals no net change in local-scale plant biodiversity over time. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 19456–9 (2013).
8. Dornelas, M. *et al.* Assemblage time series reveal biodiversity change but not systematic loss. *Science* **344**, 296–9 (2014).
9. Cardinale, B. Overlooked local biodiversity loss. *Science* **344**, 1098 (2014).
10. Alkemade, R. *et al.* GLOBIO3: A Framework to Investigate Options for Reducing Global Terrestrial Biodiversity Loss. *Ecosystems* **12**, 374–390 (2009).
11. Gibson, L. *et al.* Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* **478**, 378–381 (2011).

12. Mendenhall, C. D., Karp, D. S., Meyer, C. F. J., Hadly, E. A. & Daily, G. C. Predicting biodiversity change and averting collapse in agricultural landscapes. *Nature* **509**, 213–217 (2014).
13. Pereira, H. M. *et al.* Essential biodiversity variables. *Science* **339**, 277–278 (2013).
14. Weber, E. & Li, B. Plant invasions in China: what is to be expected in the wake of economic development? *Bioscience* **58**, 437–444 (2008).
15. Clements, G. R. *et al.* Where and how are roads endangering mammals in Southeast Asia's forests? *PLoS One* **9**, e115376 (2014).
16. Hudson, L. N. *et al.* The PREDICTS database: a global database of how local terrestrial biodiversity responds to human impacts. *Ecol. Evol.* **4**, 4701–4735 (2014).
17. Chapman, A. D. *Numbers of Living Species in Australia and the World*. (Australian Biological Resources Study, 2009).
18. Phalan, B., Onial, M., Balmford, A. & Green, R. E. Reconciling food production and biodiversity conservation: land sharing and land sparing compared. *Science* **333**, 1289–1291 (2011).
19. Balmford, A. Extinction filters and current resilience: the significance of past selection pressures for conservation biology. *Trends Ecol. Evol.* **11**, 193–196 (1996).
20. Newbold, T. *et al.* A global model of the response of tropical and sub-tropical forest biodiversity to anthropogenic pressures. *Proc. R. Soc. London Ser. B Biol. Sci.* **281**, 20141371 (2014).
21. Benítez-López, A., Alkemade, R. & Verweij, P. A. The impacts of roads and other infrastructure on mammal and bird populations: a meta-analysis. *Biol. Conserv.* **143**, 1307–1316 (2010).
22. Murphy, G. E. P. & Romanuk, T. N. A meta-analysis of declines in local species richness from human disturbances. *Ecol. Evol.* **4**, 91–103 (2014).
23. Magurran, A. E. *Measuring Biological Diversity*. (Wiley-Blackwell, 2004).
24. Barlow, J. *et al.* Quantifying the biodiversity value of tropical primary, secondary, and plantation forests. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 18555–18560 (2007).
25. Dent, D. H. & Wright, S. J. The future of tropical species in secondary forests: A quantitative review. *Biol. Conserv.* **142**, 2833–2843 (2009).
26. Hurtt, G. C. *et al.* Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands. *Clim. Change* **109**, 117–161 (2011).
27. Díaz, S. *et al.* Functional traits, the phylogeny of function, and ecosystem service vulnerability. *Ecol. Evol.* **3**, 2958–75 (2013).
28. Cardillo, M. *et al.* Multiple causes of high extinction risk in large mammal species. *Science* **309**, 1239–1241 (2005).
29. Mayfield, M. M. *et al.* Differences in forest plant functional trait distributions across land-use and productivity gradients. *Am. J. Bot.* **100**, 1356–1368 (2013).
30. Séguin, A., Harvey, É., Archambault, P., Nozais, C. & Gravel, D. Body size as a predictor of species loss effect on ecosystem functioning. *Sci. Rep.* **4**, 4616 (2014).
31. Wearn, O. R., Reuman, D. C. & Ewers, R. M. Extinction debt and windows of conservation opportunity in the Brazilian Amazon. *Science* **337**, 228–232 (2012).
32. Harfoot, M. *et al.* Integrated assessment models for ecologists: the present and the future. *Glob. Ecol. Biogeogr.* **23**, 124–143 (2014).
33. Ellis, E. C. Anthropogenic transformation of the terrestrial biosphere. *Philos. Trans. R. Soc. London Ser. A Math. Phys. Eng. Sci.* **369**, 1010–1035 (2011).
34. Mora, C. *et al.* The projected timing of climate departure from recent variability. *Nature* **502**, 183–187 (2013).

35. Burrows, M. T. *et al.* Geographical limits to species-range shifts are suggested by climate velocity. *Nature* **507**, 492–495 (2014).
36. Oldfield, F. & Steffen, W. Anthropogenic climate change and the nature of Earth System science. *Anthr. Rev.* **1**, 70–75 (2014).
37. Pereira, H. M. *et al.* Scenarios for global biodiversity in the 21st century. *Science* **330**, 1496–1501 (2010).
38. Warren, R. *et al.* Quantifying the benefit of early climate change mitigation in avoiding biodiversity loss. *Nat. Clim. Chang.* **3**, 678–682 (2013).
39. Millennium Ecosystem Assessment. *Ecosystems and Human Well-being: Biodiversity Synthesis*. (World Resources Institute, 2005).
40. Van Vuuren, D. P. *et al.* RCP2.6: exploring the possibility to keep global mean temperature increase below 2°C. *Clim. Change* **109**, 95–116 (2011).
41. Thomson, A. M. *et al.* RCP4.5: a pathway for stabilization of radiative forcing by 2100. *Clim. Change* **109**, 77–94 (2011).
42. Riahi, K. *et al.* RCP 8.5—A scenario of comparatively high greenhouse gas emissions. *Clim. Change* **109**, 33–57 (2011).
43. Rogelj, J., Meinshausen, M. & Knutti, R. Global warming under old and new scenarios using IPCC climate sensitivity range estimates. *Nat. Clim. Chang.* **2**, 248–253 (2012).
44. Bates, D., Maechler, M., Bolker, B. & Walker, S. lme4: Linear mixed-effects models using Eigen and S4. (2013). at <<http://cran.r-project.org/web/packages/lme4/>>
45. Center for International Earth Science Information Network (CIESIN) Columbia University, International Food Policy Research Institute (IFPRI), The World Bank & Centro Internacional de Agricultura Tropical (CIAT). *Global Rural-Urban Mapping Project, Version 1 (GRUMPv1): Population Density Grid*. (NASA Socioeconomic Data and Applications Center (SEDAC), 2011). <http://sedac.ciesin.columbia.edu/data/dataset/grump-v1-population-density> (Accessed 11 July 2012).
46. Center for International Earth Science Information Network (CIESIN) Columbia University & Information Technology Outreach Services (ITOS) University of Georgia. *Global Roads Open Access Data Set, Version 1 (gROADSv1)*. (NASA Socioeconomic Data and Applications Center (SEDAC), 2013). <http://dx.doi.org/10.7927/H4VD6WCT> (Accessed 18 December 2013).
47. Nelson, A. *Estimated travel time to the nearest city of 50,000 or more people in year 2000*. (2008). <http://bioval.jrc.ec.europa.eu/products/gam/index.htm> (Accessed 14 July 2014).
48. Klein Goldewijk, K., Beusen, A., Van Drecht, G. & De Vos, M. The HYDE 3.1 spatially explicit database of human-induced global land-use change over the past 12,000 years. *Glob. Ecol. Biogeogr.* **20**, 73–86 (2011).

Supplementary Information is linked to the online version of the paper at www.nature.com/nature.

Additional references can be found in the Supplementary Information.

Acknowledgements

We thank all the many researchers who have made their data available to us; Stuart Butchart and Birdlife International for sharing bird body-size data; Francis Gilbert for hoverfly body-size data; the IMAGE, HYDE, MESSAGE and MiniCAM teams, especially Rob Alkemade, Michel Bakkenes and Allison Thomson, for sharing additional data from their integrated assessment models; Derek Tittensor for statistical advice; Chris Sleep and Srinivas Patlola at

the Natural History Museum in London for IT support with the database; members of the GARD initiative (<http://www.gardinitiative.org/index.html>) for help with estimating the reptile species richness map; Kate Jones, Jason Tylianakis, Michael Crawley and E.J. Milner-Gulland for discussion, Neil Burgess for comments on a draft of the paper. We also thank Chris D. Thomas and two anonymous reviewers for very helpful comments on the manuscript. This study is part of the PREDICTS (Projecting Responses of Ecological Diversity In Changing Terrestrial Systems) project, which is supported by the UK Natural Environment Research Council (NERC, grant number: NE/J011193/1), the Biotechnology and Biological Sciences Research Council (grant number: BB/F017324/1) a Hans Rausing PhD scholarship. The study was also supported by the TRY initiative on plant traits, whose database is maintained at Max-Planck-Institute for Biogeochemistry, Jena, Germany, and which is supported by DIVERSITAS, IGBP, the Global Land Project, NERC, the French Foundation for Biodiversity Research, and GIS "Climat, Environnement et Société" France. This is a contribution from the Imperial College Grand Challenges in Ecosystem and the Environment Initiative

Author contributions

T.N., L.N.H., S.L.L.H., S.C., I.L., B.C., D.W.P., R.M.E., G.M.M., J.P.W.S. and A.P. designed the project and this study; T.N., L.N.H., I.L., R.A.S., L.B., J.P.W.S. and A.P. performed the analyses; T.N., L.N.H., S.L.L.H., S.C., D.J.B., A.C., B.C., J.D., A.D.P., S.E.-L., M.G., M.L.K.H., T.A., D.J.I., V.K., L.K., D.L.P.C., C.D.M., Y.P., H.R.P.P., A.R., J.S., H.J.W. and A.P. collated the assemblage composition data; T.N., L.N.H., S.L.L.H., S.C., A.D.P., I.L., H.R.P.P., J.P.W.S. and A.P. designed the data-collection protocols and database; R.A.S., S.D., M.E., A.F., Y.I., J.K., M.K., S.M. and E.W. made substantial contributions to the trait data used in the analyses and S.T. to the site-level environmental data; R.A.S., A.F., Y.I., S.M., and M.N. generated the maps of species richness used in the model projections; T.N., L.N.H. and A.P. wrote the manuscript with contributions from G.M.M., L.B., D.W.P., R.M.E., A.D.P., H.R.P.P., S.L.L.H., R.A.S., B.C., S.D., A.F., Y.I., J.K., M.K., S.M., J.P.W.S. and S.T.; T.N. and L.N.H. contributed equally to the study.

Author information

The authors declare no competing financial interests. Correspondence and requests for material should be addressed to T.N. (Tim.Newbold@unep-wcmc.org).

Tables

Table 1. Key features of the four Representative Concentration Pathways (RCP) scenarios (land-use and human-population assumptions are detailed in ref. 26, energy assumptions in refs. 40–42, and climate implications in ref. 43).

Scenario	Land use (see also Extended Data Figure 7)	Climate and energy	Human population
IMAGE 2.6	Agriculture moves from developed to developing countries. Large increase in area of biofuel plantations. Urban extent assumed constant.	Increased energy efficiency. Increased use of carbon capture and storage, nuclear, renewable energy and biofuels. Approximately 1 °C temperature increase by 2100 compared to pre-industrial.	10.1 billion by 2100 (UN Medium variant, 2010)
MiniCAM 4.5	Carbon pricing leads to preservation of primary forest and expansion of secondary forest. Crop yield increases, improved agricultural efficiency and dietary shifts lead to decreases in cropland and pasture areas. Small increase in area of biofuel plantations. Urban extent assumed constant.	Decline in overall energy use. Decreased use of fossil fuels and increase in nuclear and renewable energy, and in carbon capture and storage. Moderate increase in use of biofuels, but limited by availability of biomass. Approximately 1.75 °C temperature increase by 2100.	Peaks above 9 billion around 2065, then declines to 8.7 billion by 2095.
AIM 6.0	Urban area increases owing to human population growth. Cropland area increases to meet food demand. Pasture area declines strongly.	Approximately 2.5 °C temperature increase by 2100.	9.1 billion by 2100 (UN Medium variant, 2004)
MESSAGE 8.5	Increasing crop yields and intensification account for much of the increased production required, but area of cropland and, to a lesser extent, pasture increases rapidly. Small increase in area of biofuel plantations. Urban area increases owing to increased population.	Small improvements in efficiency leading to high demand for energy. Conventional oil and gas become scarce, leading to shift in favour of unconventional and carbon-intensive fossil fuels. Moderate increase in use of biofuels. Approximately 4 °C increase in temperature by 2100.	12 billion by 2100.

Figure legends

Figure 1. Locations of sites and responses of four metrics of local diversity to human pressures. **a**, Sites used in the models. Responses⁴⁴ of richness, **b**, total abundance, **c**, and community-weighted mean organism size – plant height (crosses) and animal mass (triangles), **d**, to anthropogenic variables. Error bars show 95% confidence intervals. Primary = primary vegetation, YSV = young secondary vegetation, ISV = intermediate secondary vegetation, MSV = mature secondary vegetation, Plantation = plantation forest. Land-use intensity: minimal (circle), light (triangle), intense (diamond), and combined light and intense (square). HPD = human population density⁴⁵, PR = proximity to roads⁴⁶ (as $-\log(\text{distance to nearest road})$), ACC = accessibility to humans⁴⁷ (as $-\log(\text{travel time to nearest major city})$) are shown as fitted effects from a model with no interactions between continuous effects and land use, at the lowest (L), median (M) and highest (H) values in the dataset. Sample sizes are given in full in the Methods (Table S2).

Figure 2. Similarity in assemblage composition as a function of land use. **a**, Average dissimilarity of species composition ($1 - \text{Sørensen Index}$) between pairs of sites within and among land uses (shown relative to the similarity between pairs of primary-vegetation sites); blue and red colours indicate, respectively, more and less similar composition; numbers indicate numbers of studies within which comparisons could be made. **b**, Clustering of land-use types based on average compositional dissimilarity; urban sites were excluded owing to the small sample size. Land-use labels as in Figure 1.

Figure 3. Net change in local richness caused by land use and related pressures by 2000. Projections used an IMAGE reference scenario¹⁰. The baseline landscape was assumed to be entirely uninhabited, unused primary vegetation. Shown using a Lambert Cylindrical Equal-Area projection at $0.5^\circ \times 0.5^\circ$ resolution.

Figure 4. Projected net change in local richness from 1500 to 2095. Future projections were based on the four RCP scenarios (Table 1). Historical (shading) and future (error bars) uncertainty is shown as 95% confidence intervals, rescaled to zero in 2005. The baseline for projections is a world entirely composed of uninhabited, unused primary vegetation; thus, the value at 1500 is not constrained to be zero because by then non-primary land uses were present – and in some regions widespread. The global average projection for MESSAGE 8.5 does not join the historical reconstruction because that scenario's human-population projections start in 2010 and because human population and plantation-forest extent have not been harmonized among scenarios.

Figure 5. Biodiversity projections at the country level. **a-b**, Country-level projections of average net local richness change between 2005 and 2095 under the worst (**a**, MESSAGE 8.5) and best (**b**, MiniCAM 4.5) RCP scenarios for biodiversity, shown in relation to the Human Development Index. Colours indicate biogeographic realms; colour intensity reflects natural vertebrate species richness (more intense = higher richness); point size is proportional to (log) country area. **c**, Correlation between projected richness changes under the MiniCAM 4.5 and MESSAGE 8.5 scenarios, with dashed line showing equality; colours as in **a** and **b**; colour intensity is proportional to the HDI (more intense = higher index).

Methods

Data collation

Between March 2012 and April 2014 we collated among-site comparisons of ecological assemblage composition from published studies (or from unpublished datasets where the methods have been published) examining the effect of human activities on more than one named taxon. A full description of how the dataset was assembled and curated is presented elsewhere¹⁶. We define sites to be in the same study if they were sampled using the same methodology and the data were reported in the same paper; therefore, some publications contain multiple studies. After six months of broad searching, we targeted efforts towards under-represented taxa, habitat types, biomes and regions. We accepted data only from published or in-press papers, or data collected using a published methodology, and we required that the data providers agreed to our making their data publicly available at the end of the PREDICTS (Projecting Responses of Ecological Diversity In Changing Terrestrial Systems) project in 2015. We accepted data only where abundance, occurrence or richness had been measured at two or more sampling locations and/or times, and where all sites were sampled using the same sampling procedure and with either the same effort or site-specific data on effort. We used geographical coordinates preferentially from the paper or supplied by data providers; but where coordinates were not thus available, we georeferenced them from maps in the papers. The final dataset came from 378 studies⁴⁹⁻³³² and two unpublished datasets (M. E. Hanley, 2005 and 2011) that were collected with published methods¹⁴⁶.

Studies compared from 2 to 360 sites (median = 15; 82% had ≥ 5 sites); most sampled species from multiple families but fewer than half sampled multiple orders. Over 70% of sites were from studies that sampled entire communities within a taxonomic group rather than a target list of species. Removing studies having a target list did not substantially alter model coefficients (results not shown) and increased the projected global net average loss of local species richness until 2005 by 0.6%. Sites varied in the maximum linear extent sampled (median 106 m; interquartile range 50 m to 354 m). Model coefficients for the approximately 50% of studies that reported maximum linear extent were robust to its inclusion in the models (results not shown).

The great majority of listed taxa were species-level, although many could not be given explicit species identifications (e.g. morpho-species)¹⁶; henceforth we refer to distinct taxa in our dataset as species. We matched taxonomic names given in the source paper to the Catalogue of Life 2013 Annual Checklist (COL)³³³, obtaining the full taxonomic classification. In order to relate the taxonomic names to species-level trait databases, we generated, for each taxon, a 'best-guess' Latin binomial as: a) the taxon name from COL if the COL query returned a species-level identification; b) the first two words of the text returned by the COL query if this was a sub-species designation; c) the first two words of the taxon name in the source publication if the COL query returned neither a species or sub-species name, and the taxon name in the source publication contained two or more words. Taxa that met none of these criteria were not matched to trait data, but were included in the calculation of richness and total abundance, and for estimating turnover in community composition among sites.

The resulting data set contained data for 26,953 species at 11,525 sites. For many high-diversity taxa, the database contains data for more than 1% of the number of species thought to have been formally described (Extended Figure 1a). The distribution of sites among major biomes is roughly proportional to the amount of terrestrial net primary productivity (NPP) fixed within each biome (Extended Figure 1b).

Site-level composition and diversity

We computed four site-level biodiversity metrics: within-sample species richness, total abundance, rarefaction-based richness and community-weighted mean organism size. These were calculated as follows.

Within-sample species richness was calculated as the number of differently-named taxa recorded at a given site in a standardised sampling unit (a measure also known as species density³³⁴). We gave precedence to the author's classification of species, even where a search of global databases revealed potential synonymies, because only certain taxonomic groups could be reliably matched to accepted taxonomies.

This measure of richness is appropriate for conservation questions but among site-differences could be due to effects on numbers of individuals as well as to changes in the shape of the species accumulation curve³³⁴. We therefore also calculated **rarefaction-based species richness** by taking 1000 random samples of n individuals from each site, where n is the smallest total number of individuals recorded at any site within its study, and calculating the mean species richness across samples. This index could only be calculated for sites where, in addition to the criteria above being met, abundance was recorded as number of individuals. Rarefied species richness was rounded to the nearest integer for analysis with Poisson errors.

Total abundance was calculated as the sum of the measures of abundance of all taxa at a site; we were thus unable to estimate abundance for sites where only species occurrence or overall richness or diversity had been recorded (17% of sites). Some abundance metrics – those not reported as densities per unit time, distance, area or volume sampled – were sensitive to sampling effort. When a study reported any of these metrics and sampling effort varied among sites within a study, we corrected the raw abundance measurements for the sampling effort expended at each sampling location and time; this was done by rescaling the sampling efforts within each study, so that the most heavily sampled site had a value of one, and then dividing the raw abundance measurements by this relative sampling effort. The rescaling was performed to prevent introducing additional heterogeneity in the modelled abundance values.

Community-weighted mean organism size was calculated as the arithmetic mean of log-transformed height of plants (available for 4,235 species in our dataset) or the log-transformed body mass or volume of vertebrates, beetles and hoverflies (5,236 species) present at a site, weighted by abundance³³⁵. Plant height data were taken from the TRY database³³⁶; for 61 species where plant vegetative height data were unavailable, we estimated it from generative height from a regression across the 2,554 species with estimates of both traits ($R^2 = 0.91$). Data on vertebrate body mass were taken from the PanTHERIA database for mammals³³⁷, from BirdLife International's World Bird Database for birds, and from a wide range of published and grey-literature sources for amphibians^{338–384}. Length data for reptiles were taken from published^{385,386} and unpublished (S. Meiri & A. Feldman, unpublished data) sources, and converted to estimates of body mass using published length-mass allometries^{387,388}. Arthropod size data (beetles and hoverflies) were collated from published sources^{389,390}. Beetle length and amphibian snout-vent length values were raised to the power three so that they had the same dimensionality as the other animal size measures. For both plant height and vertebrate body mass, missing values were interpolated as the average values for congeners, since both of these traits are strongly conserved phylogenetically (Pagel's $\lambda = 0.98, 0.997, 0.93, 0.89$ for plant height, vertebrate body mass, beetle body length and hoverfly thorax volume, respectively).

Human pressure data

While many human pressures can impact local biodiversity, we focus on those that can be obtained for sites around the world and for which, as far as possible, spatiotemporal data are available for 1500-2095; this focus enables us to use our statistical models as a basis for projecting responses through time. Each site was assigned to one of eight land-use classes based on the description of the habitat given in the source paper (see Extended Data Table 1 for definitions): primary vegetation, secondary vegetation (subdivided into mature, intermediate or young secondary vegetation), plantation forest, cropland, pasture and urban¹⁶. These classes were selected to match the land-use classification adopted in the Intergovernmental Panel on Climate Change Representative Concentration Pathways scenarios²⁶ in order to facilitate the projection of our models onto these scenarios. Sites were also assigned to a level of human intensity of use – minimal, light or intense – within each major land-use class, also based on the description of the habitat in the source paper (see Extended Data Table 1 for definitions). The factors that determined this level depended on the land-use class (e.g. bushmeat extraction and limited logging in primary and secondary vegetation; or stocking density and chemical inputs in pasture; Extended Data Table 1). Sites that could not be classified for land-use and use intensity were excluded from the analyses. The final dataset contained the following numbers of sites in each land use and land-use intensity level: Primary vegetation, minimal use – 1546 (from 183 studies), light use – 860 (76 studies), intense use – 449 (33 studies); mature secondary vegetation, minimal use – 198 (52 studies), light/intense use – 213 (23 studies); intermediate secondary vegetation, minimal use – 404 (55 studies), light/intense use – 269 (30 studies); young secondary vegetation, minimal use – 431 (50 studies), light/intense use – 331 (34 studies); plantation forest, minimal use – 356 (47 studies), light use – 402 (42 studies), intense use – 238 (29 studies); cropland, minimal use – 427 (45 studies), light use – 632 (43 studies), intense use – 703 (36 studies); pasture, minimal use – 525 (43 studies), light use – 434 (52 studies), intense use – 174 (23 studies); and urban, minimal use – 174 (23 studies), light use – 244 (26 studies), intense use – 195 (18 studies).

We overlaid our sites with available global data sets to obtain site-level estimates of human population density⁴⁵, distance to the nearest road⁴⁶ and estimated travel time to nearest population centre with greater than 50,000 inhabitants⁴⁷. For distance to nearest road, the map of roads was first projected onto a Berhmann equal-area projection. These operations were carried out using Python code implemented using the *arcpy* Python module in ArcMap Version 10.0³⁹¹. In the main figures, the inverses of distance to roads and travel time to major population centre (proximity to roads and accessibility) were presented so that high values corresponded to higher hypothesized human impact. To estimate the history of human use of the landscapes within which sites were located, we calculated the number of years since the 30-arc-second grid cell containing each site became 30% covered by human land uses (cropland, pasture and urban), according to the HYDE model⁴⁸. Collinearity among variables describing anthropogenic change was low: the highest correlation was between land use and human population density (Pearson $R^2 = 0.31$).

Modelling site-level diversity, composition and turnover

The response of site-level diversity to the measures of anthropogenic change was modelled using generalized linear mixed effects models, implemented in the *lme4* package Version 1.0-5⁴⁴ in R Version 3.0.2³⁹². We first compared candidate random-effects structures using the full candidate fixed-effects structure³⁹³. Random-intercept terms considered in all

models were the identity of study from which data were taken, to account for study-level differences in the response variables and sampling methods used, and – within-study – the spatial block in which the site was located, to account for the spatial arrangement of sites. For models of species richness (within-sample and rarefied), we also fitted an observation-level random effect (i.e. site identity) to account for the overdispersion present³⁹⁴. We also considered random slopes, with respect to study, of each of the main fixed effects (land use, land-use intensity, human population density, distance to nearest road, travel time to nearest major city and time since the landscape was majority converted to human uses). Random effects were retained or discarded based on Akaike Information Criterion values.

Once the best random-effects structure had been selected, we performed backward stepwise model simplification to select the best fixed-effects structure (see ‘Full Statistical Results’)³⁹³. Human population density, distance to roads, travel time to nearest major city and time since major human use of the landscape were log transformed in the analyses, with a value of 1 added to human population density, travel time to nearest major city and time since major landscape conversion to deal with zero values. These four variables were fitted as continuous effects, with quadratic polynomials for human population, distance to roads and travel time to nearest major city, and as a linear effect for time since human landscape conversion. For variables fitted as quadratic polynomials, we also tested linear effects during the backward stepwise model selection. All continuous variables were rescaled prior to analysis so that values ranged between zero and one. Interaction terms were tested first, and then removed to test the main effects. All main effects that were part of significant interaction terms were retained in the final models regardless of their significance as main effects. For the model of community-weighted mean body mass and plant height, because the number of sites with data was smaller than for the other metrics, only land use (excluding urban sites, which were few), human population density and distance to roads, and no interactions, were fitted (for the model of plant height, sample sizes in each land use were: primary vegetation – 634 sites, secondary vegetation – 851 sites, plantation forest – 222 sites, cropland – 72 sites, pasture – 412 sites; and for the model of animal mass: primary vegetation – 1728 sites, secondary vegetation – 805 sites, plantation forest – 602 sites, cropland – 641 sites, pasture – 440 sites). The decision whether or not to retain terms was based on likelihood ratio tests. The coefficient estimates of the best models are shown in Figure 1b-d and Extended Data Figure 2, and the formulae and statistical results are shown in the Supplementary Information. To test for spatial autocorrelation in the residuals of the final best models, we calculated Moran’s I values and associated P-values, separately for each study considered in the models, using the spdep package Version 0.5-68³⁹⁵ in R; the distribution of P-values across studies was used as an indication of whether spatial autocorrelation was likely to cause a problem. This revealed that the residuals showed little spatial autocorrelation (Extended Data Figure 5). We used cross validation to assess the robustness of model parameter estimates, first based on dividing the studies randomly into 10 equal-sized sets and dropping each set in turn (Extended Data Figure 3c), and secondly based on leaving out the studies from each biome in turn (Extended Data Figure 3d).

Publication bias is a potential problem for any large-scale synthesis of data from many publications. In standard meta-analyses, funnel plots³⁹⁶ can be used to test for any relationship between standard error and effect size, as a bias in effect sizes at high standard error toward more positive or more negative effects indicates a likely effect of publication bias. Creating funnel plots for our data was more complicated because ours was a site-level analysis of raw diversity estimates rather than a traditional meta-analysis. Instead we generated individual models relating diversity to land use for each study that sampled at least two sites within each of at least two land-use types. We focused on land use because: a) there were a small number of sites included in most within-study models; and 2) the original

studies focused on effects of land use – not generally on land-use intensity, human population density or distance to roads – and thus any effect of publication bias would likely be seen in the land-use coefficients. Funnel plots were generated by plotting, for each land-use type, the estimated model coefficients against the associated standard errors (Extended Data Figure 4). There were some indications of an effect of publication bias, with less certain coefficient estimates tending to have more negative estimates for some of the land uses (Extended Data Figure 4). On the other hand, study-level random slopes of human-dominated land uses tended to be more negative for studies that sampled more sites (Extended Data Figure 4). It is important to emphasize that in a site-level analysis like ours, studies with fewer sites have less weight in the models. Modelled coefficient estimates were generally robust to the removal of these studies (Extended Data Figure 4). Basing projections on coefficient estimates from models where small studies were excluded led to a less than 1% change in the estimated global richness values (results not shown). As with all studies based on data from the literature, we under-represent unpublished data.

To model turnover of species composition between pairs of sites, we calculated average dissimilarity²³ in the lists of present species (1 – Sørensen index) between all pairs of sites within each study. For this analysis, we were only able to consider studies with more than one site in at least one of the land-use types considered. Once compositional similarity had been calculated for every pair of sites within each study, the average compositional similarity was calculated for every pair of land-use types considered within each study (including comparisons between sites in the same land-use type). Finally, the average compositional similarity was calculated for each pair of land-use types across all studies. To visualize the clustering of different land-use types in terms of community composition, we performed a hierarchical complete-linkage cluster analysis on the compositional dissimilarity (i.e. 1 – similarity) matrix, using the hclust function in R Version 3.0.2³⁹². To test whether differences in the average geographic distance between pairs of sites in different land-use combinations affected these results, we correlated average compositional similarity with average distance between sites, for all pairwise combinations of land use (including comparisons of a land-use type with itself). Correlations between average distance and average community similarity were only very weakly negative ($R^2 = 0.001$), suggesting they do not strongly distort the comparisons of community composition. However, the fact that some land uses tend to occur more closely together than others could influence the diversity patterns seen in our models, if some land uses are typically close to high-diversity habitats and so are more likely to benefit from dispersal. For example, sites in secondary vegetation and plantation forest were closer, on average, to primary vegetation sites than were those in cropland, pasture and urban (average distances to sites in primary vegetation were: other primary vegetation sites = 7.38 km, mature secondary vegetation = 4.4 km, intermediate secondary vegetation = 3.9 km, young secondary vegetation = 6.9 km, plantation forest = 4.2 km, cropland = 16.4 km, pasture = 10.1 km, and urban = 11.4 km). Accounting for distance in such already-complex models is not computationally tractable. In making the projections, we therefore implicitly assume that the average distances will not change (i.e. that secondary vegetation and plantation forests will remain closer to primary vegetation than cropland, pasture and urban habitats).

Projecting the Models onto Spatial Estimates of Anthropogenic Variables

We projected the best overall models of richness (within-sample and rarefied), abundance and community-weighted mean organism size onto estimates of land use, land-use intensity and human population density at $0.5^\circ \times 0.5^\circ$ resolution, using historical estimates for 1500 to 2005, and four Representative Concentration Pathways scenarios of future

changes (IMAGE 2.6, MiniCAM 4.5, AIM 6.0 and MESSAGE 8.5; the names refer to the integrated assessment models used and the numbers to the amount of radiative forcing assumed in 2100)³⁹⁷. In the absence of global projections, proximity to roads and accessibility were omitted from our projections.

Estimates of land use for both the historical reconstruction and the future scenarios were taken from the harmonized land-use data accompanying the scenarios²⁶. Estimates of the stage of secondary vegetation (young, intermediate or mature) are not available directly in the RCP land-use data. However, these data contain estimates of the transition each year between secondary vegetation and all other land-use types. To convert this into an estimate of the proportion of secondary vegetation in each of the stages of maturity, we considered any transition to secondary vegetation to result in secondary vegetation of age zero. Each year, this age was then incremented by one. In the absence of better information, any transitions from secondary vegetation to any other land-use type were assumed to be drawn evenly from the ages currently represented. For the purposes of the projections, secondary vegetation was considered to be young until an age of 30 years, intermediate between 30 years and 100 years, and mature thereafter. We developed C# code to convert land-use transitions into estimates of the stage structure of secondary vegetation.

Gridded temporal estimates of human population density were directly available for the HYDE historical scenario and MESSAGE future scenario. Human population trajectories in the MiniCAM model were resolved only to the level of United Nations regions⁴¹; we therefore downscaled these to grid cells assuming no temporal change in the spatial pattern of relative population density within regions compared to present day patterns⁴⁵, which is the method used in other RCP-scenario land-use models lacking human population data resolved to grid cells²⁶. Gridded estimates of human population from the MESSAGE model were downloaded from <http://www.iiasa.ac.at/web-apps/ggi/GgiDb/>. For the scenarios for which human population projections were not available (IMAGE and AIM), we used country-level estimates from the ‘medium’ scenario of the United Nations population division³⁹⁸, which gives the closest global predictions of future human population to those assumed by IMAGE and AIM²⁶. These country-level estimates were downscaled to grid cells using the same method as for MiniCAM’s regional projections.

Land-use intensity was an important explanatory variable in our models, but global maps of land-use intensity are not available. We therefore generated global estimates of current land-use intensity based on a map of ‘Global Land Systems’³⁹⁹, which divides coarse land-use types into sub-categories based on levels of cropland intensity, livestock densities and human population density. We mapped each Global Land Systems class onto one or more relevant combinations of our classes of land use and land-use intensity (Extended Data Table 2). The Global Land Systems dataset has a spatial resolution of 5 arc-minutes. To calculate the proportion of each 0.5° cell occupied by each land use and land-use intensity combination we calculated the proportion of 5-arc-minute cells within each 0.5° cell containing matching Global Land Systems categories (see legend of Extended Data Table 2 for details).

To generate past and future estimates of land-use intensity, we modelled the current proportion of each land-use type estimated to be under minimal, light or intense levels of intensity within each grid cell (one model for each intensity level), as a function of the prevalence of the land-use type within each cell and human population density, with the relationships allowed to vary among the 23 United Nations (UN) sub-regions (i.e. we fitted interaction terms between UN sub-region and both the prevalence of each land-use type and human population density). UN sub-region data were taken from the world borders shapefile Version 0.3 (http://thematicmapping.org/downloads/world_borders.php) and converted to a 0.5° × 0.5° raster using ArcMap Version 10.0³⁹¹. The models were developed using

generalized linear models with a binomial distribution of errors, implemented in the lme4 package Version 1.0-5⁴⁴ in R Version 3.0.2³⁹². The resulting models explained between 30.6% and 76.7% of the deviance in estimated current levels of intensity. Past and future land-use intensities were estimated by applying the models to the same past and future estimates of land use and human population density as above.

The scenarios gave the proportion of each grid cell estimated to be occupied by each combination of land use and land-use intensity. We did not attempt to resolve human population density within grid cells for our historical estimates or forecasts, thereby assuming it to be spatially (not temporally) constant within each cell. The coefficients from the models of site-level diversity were thus applied to each combination of land use and intensity within each cell, with the same human population density estimate across all combinations. All predictions were expressed as a percentage net change compared with a baseline before human land-use impacts on biodiversity, in which all land use was assumed to be primary vegetation of minimal intensity of use, and with a human population density of zero. Each cell's average value of net biodiversity change was calculated as the area-weighted mean value across all land uses and intensities. Global average values were calculated as mean values across all cells, weighted by cell area and an appropriate weighting factor to account for the fact that cells have different baseline levels of diversity. The weighting factors applied were: terrestrial vertebrate species richness in the case of richness, and net primary production (NPP) in the case of total abundance. No weighting factor was applied for projections of community-weighted mean plant height. Terrestrial vertebrate species richness was estimated by overlaying extent-of-occurrence range maps for mammals, birds, amphibians and reptiles, using Python code written by ourselves and implemented in ArcMap Version 10.0³⁹¹. Data on Net Primary Production were estimates of potential NPP (i.e. in the absence of human impacts) from the Lund-Postdam-Jena (LPJ) Dynamic Global Vegetation Model⁴⁰⁰.

The 95% confidence intervals around the projected values of biodiversity for each combination of pressure variables were estimated based on uncertainty in the modelled coefficients. We were unable to conduct multi-model averaging to account for uncertainty in the structure of the models (i.e. projections were based only on the final best model) because applying such complex mixed-effects models, based on such large datasets, to multiple scenarios of human pressure at a global scale was intractable both in terms of time and computer-memory requirements. We were also unable to account for uncertainty in the trajectories of the human pressure variables, because uncertainty estimates are not available for any of the variables considered.

To estimate average biodiversity change in individual countries, we intersected the gridded projections with the world borders shapefile (see above) using the extract function in the raster package Version 2.2-12⁴⁰¹ in R Version 3.0.2³⁹². Mean values across the cells associated with each country were calculated, weighted by cell area. To interpret the outcomes for countries in terms of their natural biodiversity, we related the country-level projections to estimates of average natural vertebrate species richness (see above). To interpret the outcomes for countries in terms of their socio-economy, we related the projections to estimates of the Human Development Index, which is an indicator of education, life expectancy, wealth and standard of living (<https://data.undp.org/>).

Extended Data Legends

Extended Data Figure 1. Taxonomic and geographic representativeness of the dataset used. **a,** The relationship between the number of species represented in our data and the number estimated to have been described¹⁷, for 47 major taxonomic groups; lines show (from

bottom to top) 0.1%, 1% and 10% representation of described species in our dataset; magenta = invertebrates, red = vertebrates, green = plants, blue = fungi, and grey = all other taxonomic groups. **b**, the relationship across biomes⁴⁰² between the percentage of global terrestrial net primary production and the number of sites in our dataset; A = tundra, B = boreal forests and taiga, C = temperate conifer forests, D = temperate broadleaf and mixed forests, E = montane grasslands and shrublands, F = temperate grasslands, savannas and shrublands, G = Mediterranean forests, woodlands and scrub, H = deserts and xeric shrublands, J = tropical and subtropical grasslands, savannas and shrublands, K = tropical and subtropical coniferous forests, M = tropical and subtropical dry broadleaf forests, N = tropical and subtropical moist broadleaf forests, P = mangroves; note that the flooded grasslands and savanna biome is not represented in the dataset; grey line show 1:1 line.

Extended Data Figure 2. Detailed response of local diversity to human pressures.

Modelled effects (controlling for land use) of human population density (HPD), distance to nearest road, time since 30% conversion of a landscape to human uses (TSC) and time to nearest population centre with greater than 50,000 inhabitants (**a-d**), interactions between pairs of these variables (**e**), and interactions between these variables and land use (**f-i**) on site-level diversity: **a-c, f, g**, within-sample species richness; **e, h, i**, total abundance; and **d**, community-weighted mean vertebrate body mass. Shaded polygons in **a-d** show 95% confidence intervals. For clarity, shaded polygons in **f-i** are shown as $\pm 0.5 \times \text{SEM}$. Confidence intervals in **e** are omitted. Rugs along the x axes in the line graphs show the values of the explanatory variables represented in the dataset used for modelling. Only significant effects are shown. Note that distance to nearest road and travel time to major population centre measures are the raw (log-transformed) values fitted in the models rather than the proximity to roads and accessibility values (obtained as 1 minus the former values) presented in Figure 1. Sample sizes are given in full in the Methods.

Extended Data Figure 3. Robustness of modelled effects of human pressures. a, Effects of land use, human population density (HPD) and proximity to nearest road (PR) on rarefaction-based species richness. **b**, To test that any differences between these results and the results for within-sample species richness presented in the main manuscript were not because rarefied species richness could only be calculated with a smaller dataset, we also show modelled effects on within-sample species richness with the same reduced dataset. **c-d**, Cross-validated robustness of coefficient estimates for land use and land-use intensity: crosses show 95% confidence intervals around the coefficient estimates under ten-fold cross-validation, excluding data from approximately 10% of studies at a time (**c**), and under geographical cross-validation, excluding data from one biome at a time (**d**); colours, points, error bars and land-use labels are as in Figure 1 in the main text. Primary, primary vegetation; YSV, young secondary vegetation; ISV, intermediate secondary vegetation; MSV, mature secondary vegetation; plantation, plantation forest. Sample sizes are given in full in the Methods.

Extended Data Figure 4. Tests of the potential for publication bias to influence the richness models and projections. Left-hand panels (**a, d, g, j, m**) show funnel plots of the relationship between the standard error around coefficient estimates (inversely related to the size of studies) and the coefficient estimates themselves for each coarse land-use type; there is evidence for publication bias with respect to some of the land-use types, as indicated by an absence of points on one or other side of zero for studies with large standard errors (but note that small studies are down-weighted in the model); red points show studies with more than 5 sites in the land use in question (10 for secondary vegetation and plantation forest because

there were more sites for these land uses and some studies with between 5 and 10 sites showed variable responses); horizontal dashed lines show the modelled coefficients for each land use. Central panels (**b, e, h, k, n**) show the relationship between study size (log-transformed total number of sites) and the random slope of the land use in question with respect to study identity, from a random-slopes-and-intercepts model; where a significant relationship was detected using a linear model, fitted values and 95% confidence intervals are shown as a red dashed line and red dotted lines, respectively; conversely to what would be expected if publication bias was present, where significant relationships between study size and random slopes were detected, these were negative (i.e. larger studies detected more negative effects). Right-hand panels (**c, f, i, l, o**) show the robustness of modelled coefficients to removal of studies with few sites in a given land use (black points in the left-hand panels); left-hand error bars show coefficient estimates for all studies and right-hand error bars show coefficient estimates for studies with more than 5 sites in that land use (10 for secondary vegetation and plantation forest).

Extended Data Figure 5. Tests for spatial autocorrelation in the model residuals. For the four main modelled metrics of site-level diversity — **a**, within-sample species richness; **b**, total abundance; **c**, community-weighted mean plant-height; and **d**, community-weighted mean animal mass — histograms of P-values from sets of Moran's tests for spatial autocorrelation in the residuals of the best models for individual studies. The percentage of studies with significant spatial autocorrelation ($P < 0.05$; indicated by vertical red line) is shown.

Extended Data Figure 6. Current, past and future projections of all metrics of local biodiversity. Net change in local diversity caused by land use and related pressures by 2000 under an IMAGE reference scenario¹⁰: **a**, richness; **b**, rarefied richness; **c**, total abundance; and **d**, community-weighted mean plant height. Note that the values used to divide the colours are the same in all panels, but that the maximum and minimum values are different, as indicated in the legends. Historical and future estimates of net change in local diversity from 1500-2095, based on estimates of land-use, land-use intensity and human population density from the four Representative Concentration Pathways (RCP) scenarios (Table 1): **e**, richness; **f**, total abundance; and **g**, community-weighted mean plant height; historical (shading) and future (error bars) uncertainty shown as 95% confidence intervals, with uncertainty rescaled to be zero in 2005 to show uncertainty in past and future change separately; the global average projection for the MESSAGE scenario does not directly join the historical reconstruction because projections start in 2010 (human population estimates are available at 15-year intervals) and because human population (and thus land-use intensity) and plantation-forest extent have not been harmonized among scenarios. In panel **e**, the dashed line shows projected diversity change under land-use change only (i.e. without land-use intensity and human population density, the projections of which involved simplifying assumptions), and the dotted line shows projections of rarefaction-based species richness.

Extended Data Figure 7. Reconstructed and projected total global land-use areas under the Representative Concentration Pathways (RCP) scenarios. **a**, estimated total area of the major land-use types. **b-f**, estimated total area of secondary vegetation in different stages of recovery.

Extended Data Figure 8. Biodiversity projections at the country level. **a-d**, Country-level projections of net change in local richness between 2005 and 2095 under the four RCP scenarios (**a**, IMAGE 2.6; **b**, MiniCAM 4.5; **c**, AIM 6.0; and **d**, MESSAGE 8.5), shown in

relation to the Human Development Index (an indicator of education, life expectancy, wealth and standard of living) in the most recent year for which data are available. **e-f**, Country-level projections of net change in local richness between 2005 and 2095 under the best- and worst-performing RCP scenarios in terms of biodiversity (**e**. MiniCAM 4.5; **f**, MESSAGE 8.5), shown in relation to past change in biodiversity from a baseline with no human land-use effects to 2005 according to the HYDE land-use reconstruction. Colours indicate biogeographic realms (key in **b**); colour intensity reflects native vertebrate species richness (more intense = higher species richness); point size is proportional to (log) country area.

Extended Data Table 1. Land use and land-use intensity classification definitions (from ref. 16).

Extended Data Table 2. Conversion between Global Land Systems dataset and our intensity classification for each major land-use type. To estimate proportional coverage of each intensity class for each land-use type in the $0.5^\circ \times 0.5^\circ$ grid cells used for projection, we calculated the number of finer-resolution Global Land Systems³⁹⁹ cells with a matching intensity class for the land-use type in question, as a proportion of Global Land Systems cells matching any intensity class for the land-use type in question. For example, to calculate the proportion of urban land that is under intense use, we divided the number of cells with a Global Land Systems classification of ‘urban’ by the number of cells classified as ‘urban’ or ‘peri-urban and villages’. None of the Global Land Systems classes could inform about the intensity of plantation forest, and so we assumed that any plantation forest was composed of equal proportions under minimal, light and intense use.

Additional references

49. Aben, J., Dorenbosch, M., Herzog, S. K., Smolders, A. J. P. & Van Der Velde, G. Human Disturbance affects a Deciduous Forest Bird Community in the Andean Foothills of Central Bolivia. *Bird Conservation International* **18**, 363-380, doi:10.1017/s0959270908007326 (2008).
50. Adum, G. B., Eichhorn, M. P., Oduro, W., Ofori-Boateng, C. & Rodel, M. O. Two-Stage Recovery of Amphibian Assemblages Following Selective Logging of Tropical Forests. *Conservation Biology* **27**, 354-363, doi:10.1111/cobi.12006 (2013).
51. Aguilar Barquero, V. & Jiménez Hernández, F. Diversidad y distribución de palmas (Arecaceae) en tres fragmentos de bosque muy húmedo en Costa Rica. *Revista de Biología Tropical* **57**, 83-92 (2009).
52. Alberta Biodiversity Monitoring Institute (ABMI). The raw soil arthropods dataset and the raw trees & snags dataset from Prototype Phase (2003-2006) and Rotation 1 (2007-2012). (2013).
53. Alcala, E. L., Alcala, A. C. & Dolino, C. N. Amphibians and reptiles in tropical rainforest fragments on Negros Island, the Philippines. *Environmental Conservation* **31**, 254-261, doi:10.1017/s0376892904001407 (2004).
54. Alcayaga, O. E., Pizarro-Araya, J., Alfaro, F. M. & Cepeda-Pizarro, J. Spiders (Arachnida, Araneae) associated to agroecosystems in the Elqui Valley (Coquimbo Region, Chile). *Revista Colombiana De Entomologia* **39**, 150-154 (2013).
55. Ancrenaz, M., Goossens, B., Gimenez, O., Sawang, A. & Lackman-Ancrenaz, I. Determination of ape distribution and population size using ground and aerial surveys:

- 988 a case study with orang-utans in lower Kinabatangan, Sabah, Malaysia. *Animal*
 989 *Conservation* **7**, 375-385, doi:10.1017/s136794300400157x (2004).
- 990 56. Arbeláez-Cortés, E., Rodríguez-Correa, H. A. & Restrepo-Chica, M. Mixed bird
 991 flocks: patterns of activity and species composition in a region of the Central Andes
 992 of Colombia. *Revista Mexicana De Biodiversidad* **82**, 639-651 (2011).
- 993 57. Armbrrecht, I., Perfecto, I. & Silverman, E. Limitation of nesting resources for ants in
 994 Colombian forests and coffee plantations. *Ecological Entomology* **31**, 403-410,
 995 doi:10.1111/j.1365-2311.2006.00802.x (2006).
- 996 58. Arroyo, J., Iturrondobeitia, J. C., Rad, C. & Gonzalez-Carcedo, S. Oribatid mite
 997 (Acari) community structure in steppic habitats of Burgos Province, central northern
 998 Spain. *Journal of Natural History* **39**, 3453-3470, doi:10.1080/00222930500240346
 999 (2005).
- 1000 59. Azhar, B. *et al.* The influence of agricultural system, stand structural complexity and
 1001 landscape context on foraging birds in oil palm landscapes. *Ibis* **155**, 297-312,
 1002 doi:10.1111/ibi.12025 (2013).
- 1003 60. Azpiroz, A. B. & Blake, J. G. Avian assemblages in altered and natural grasslands in
 1004 the northern Campos of Uruguay. *Condor* **111**, 21-35, doi:10.1525/cond.2009.080111
 1005 (2009).
- 1006 61. Baeten, L. *et al.* Early Trajectories of Spontaneous Vegetation Recovery after
 1007 Intensive Agricultural Land Use. *Restoration Ecology* **18**, 379-386,
 1008 doi:10.1111/j.1526-100X.2009.00627.x (2010).
- 1009 62. Baeten, L., Hermy, M., Van Daele, S. & Verheyen, K. Unexpected understorey
 1010 community development after 30 years in ancient and post-agricultural forests.
 1011 *Journal of Ecology* **98**, 1447-1453, doi:10.1111/j.1365-2745.2010.01711.x (2010).
- 1012 63. Báldi, A., Batáry, P. & Erdős, S. Effects of grazing intensity on bird assemblages and
 1013 populations of Hungarian grasslands. *Agriculture Ecosystems & Environment* **108**,
 1014 251-263, doi:10.1016/j.agee.2005.02.006 (2005).
- 1015 64. Banks, J. E., Sandvik, P. & Keesecker, L. Beetle (Coleoptera) and spider (Araneae)
 1016 diversity in a mosaic of farmland, edge, and tropical forest habitats in western Costa
 1017 Rica. *Pan-Pacific Entomologist* **83**, 152-160, doi:10.3956/0031-0603-83.2.152
 1018 (2007).
- 1019 65. Barlow, J. *et al.* Quantifying the biodiversity value of tropical primary, secondary, and
 1020 plantation forests. *Proceedings of the National Academy of Sciences of the United*
 1021 *States of America* **104**, 18555-18560, doi:10.1073/pnas.0703333104 (2007).
- 1022 66. Bartolommei, P., Mortelliti, A., Pezzo, F. & Puglisi, L. Distribution of nocturnal birds
 1023 (Strigiformes and Caprimulgidae) in relation to land-use types, extent and
 1024 configuration in agricultural landscapes of Central Italy. *Rendiconti Lincei-Scienze*
 1025 *Fisiche E Naturali* **24**, 13-21, doi:10.1007/s12210-012-0211-3 (2013).
- 1026 67. Basset, Y. *et al.* Changes in Arthropod Assemblages along a Wide Gradient of
 1027 Disturbance in Gabon. *Conservation Biology* **22**, 1552-1563, doi:10.1111/j.1523-
 1028 1739.2008.01017.x (2008).
- 1029 68. Bates, A. J. *et al.* Changing Bee and Hoverfly Pollinator Assemblages along an
 1030 Urban-Rural Gradient. *Plos One* **6**, doi:10.1371/journal.pone.0023459 (2011).
- 1031 69. Baur, B. *et al.* Effects of abandonment of subalpine hay meadows on plant and
 1032 invertebrate diversity in Transylvania, Romania. *Biological Conservation* **132**, 261-
 1033 273, doi:10.1016/j.biocon.2006.04.018 (2006).
- 1034 70. Berg, A., Ahrne, K., Ockinger, E., Svensson, R. & Soderstrom, B. Butterfly
 1035 distribution and abundance is affected by variation in the Swedish forest-farmland
 1036 landscape. *Biological Conservation* **144**, 2819-2831,
 1037 doi:10.1016/j.biocon.2011.07.035 (2011).

- 1038 71. Bernard, H., Fjeldsa, J. & Mohamed, M. A case study on the effects of disturbance
1039 and conversion of tropical lowland rain forest on the non-volant small mammals in
1040 north Borneo: Management implications. *Mammal Study* **34**, 85-96,
1041 doi:10.3106/041.034.0204 (2009).
- 1042 72. Berry, N. J. *et al.* The high value of logged tropical forests: lessons from northern
1043 Borneo. *Biodiversity and Conservation* **19**, 985-997, doi:10.1007/s10531-010-9779-z
1044 (2010).
- 1045 73. Bicknell, J. & Peres, C. A. Vertebrate population responses to reduced-impact logging
1046 in a neotropical forest. *Forest Ecology and Management* **259**, 2267-2275,
1047 doi:10.1016/j.foreco.2010.02.027 (2010).
- 1048 74. Bihn, J. H., Verhaagh, M., Braendle, M. & Brandl, R. Do secondary forests act as
1049 refuges for old growth forest animals? Recovery of ant diversity in the Atlantic forest
1050 of Brazil. *Biological Conservation* **141**, 733-743, doi:10.1016/j.biocon.2007.12.028
1051 (2008).
- 1052 75. Billeter, R. *et al.* Indicators for biodiversity in agricultural landscapes: a pan-
1053 European study. *Journal of Applied Ecology* **45**, 141-150, doi:10.1111/j.1365-
1054 2664.2007.01393.x (2008).
- 1055 76. Bóçon, R. *Riqueza e abundância de aves em três estágios sucessionais da floresta*
1056 *ombrófila densa submontana, Antonina, Paraná* PhD thesis, Universidade Federal do
1057 Paraná, (2010).
- 1058 77. Borges, S. H. Bird assemblages in secondary forests developing after slash-and-burn
1059 agriculture in the Brazilian Amazon. *Journal of Tropical Ecology* **23**, 469-477,
1060 doi:10.1017/s0266467407004105 (2007).
- 1061 78. Boutin, C., Baril, A. & Martin, P. A. Plant diversity in crop fields and woody
1062 hedgerows of organic and conventional farms in contrasting landscapes. *Agriculture*
1063 *Ecosystems & Environment* **123**, 185-193, doi:10.1016/j.agee.2007.05.010 (2008).
- 1064 79. Bouyer, J. *et al.* Identification of ecological indicators for monitoring ecosystem
1065 health in the trans-boundary W Regional park: A pilot study. *Biological*
1066 *Conservation* **138**, 73-88, doi:10.1016/j.biocon.2007.04.001 (2007).
- 1067 80. Bragagnolo, C., Nogueira, A. A., Pinto-da-Rocha, R. & Pardini, R. Harvestmen in an
1068 Atlantic forest fragmented landscape: Evaluating assemblage response to habitat
1069 quality and quantity. *Biological Conservation* **139**, 389-400,
1070 doi:10.1016/j.biocon.2007.07.008 (2007).
- 1071 81. Brearley, F. Q. Below-ground secondary succession in tropical forests of
1072 Borneo. *Journal of Tropical Ecology* **27**, 413-420, doi:10.1017/s0266467411000149
1073 (2011).
- 1074 82. Brito, I., Goss, M. J., de Carvalho, M., Chatagnier, O. & van Tuinen, D. Impact of
1075 tillage system on arbuscular mycorrhiza fungal communities in the soil under
1076 Mediterranean conditions. *Soil & Tillage Research* **121**, 63-67,
1077 doi:10.1016/j.still.2012.01.012 (2012).
- 1078 83. Brunet, J. *et al.* Understory succession in post-agricultural oak forests: Habitat
1079 fragmentation affects forest specialists and generalists differently. *Forest Ecology and*
1080 *Management* **262**, 1863-1871, doi:10.1016/j.foreco.2011.08.007 (2011).
- 1081 84. Buczkowski, G. Extreme life history plasticity and the evolution of invasive
1082 characteristics in a native ant. *Biological Invasions* **12**, 3343-3349,
1083 doi:10.1007/s10530-010-9727-6 (2010).
- 1084 85. Buczkowski, G. & Richmond, D. S. The effect of urbanization on ant abundance and
1085 diversity: a temporal examination of factors affecting biodiversity. *Plos One* **7**,
1086 doi:10.1371/journal.pone.0041729 (2012).

- 1087 86. Buddle, C. M. & Shorthouse, D. P. Effects of experimental harvesting on spider
1088 (Araneae) assemblages in boreal deciduous forests. *Canadian Entomologist* **140**, 437-
1089 452, doi:10.4039/n07-LS01 (2008).
- 1090 87. Buscardo, E. *et al.* The early effects of afforestation on biodiversity of grasslands in
1091 Ireland. *Biodiversity and Conservation* **17**, 1057-1072, doi:10.1007/s10531-007-9275-
1092 2 (2008).
- 1093 88. Cabra-García, J., Bermúdez-Rivas, C., Osorio, A. M. & Chacón, P. Cross-taxon
1094 congruence of alpha and beta diversity among five leaf litter arthropod groups in
1095 Colombia. *Biodiversity and Conservation* **21**, 1493-1508, doi:10.1007/s10531-012-
1096 0259-5 (2012).
- 1097 89. Caceres, N. C., Napoli, R. P., Casella, J. & Hannibal, W. Mammals in a fragmented
1098 savannah landscape in south-western Brazil. *Journal of Natural History* **44**, 491-512,
1099 doi:10.1080/00222930903477768 (2010).
- 1100 90. Cagle, N. L. Snake species distributions and temperate grasslands: A case study from
1101 the American tallgrass prairie. *Biological Conservation* **141**, 744-755,
1102 doi:10.1016/j.biocon.2008.01.003 (2008).
- 1103 91. Calviño-Cancela, M., Rubido-Bará, M. & van Etten, E. J. B. Do eucalypt plantations
1104 provide habitat for native forest biodiversity? *Forest Ecology and Management* **270**,
1105 153-162, doi:10.1016/j.foreco.2012.01.019 (2012).
- 1106 92. Cameron, S. A. *et al.* Patterns of widespread decline in North American bumble
1107 bees. *Proceedings of the National Academy of Sciences of the United States of*
1108 *America* **108**, 662-667, doi:10.1073/pnas.1014743108 (2011).
- 1109 93. Carrijo, T. F., Brandao, D., de Oliveira, D. E., Costa, D. A. & Santos, T. Effects of
1110 pasture implantation on the termite (Isoptera) fauna in the Central Brazilian Savanna
1111 (Cerrado). *Journal of Insect Conservation* **13**, 575-581, doi:10.1007/s10841-008-
1112 9205-y (2009).
- 1113 94. Carvalho, A. L. d., Ferreira, E. J. L., Lima, J. M. T. & de Carvalho, A. L. Floristic and
1114 structural comparisons among palm communities in primary and secondary forest
1115 fragments of the Raimundo Irineu Serra Environmental Protection Area - Rio Branco,
1116 Acre, Brazil. *Acta Amazonica* **40**, 657-666, doi:10.1590/s0044-59672010000400004
1117 (2010).
- 1118 95. Castro, H., Lehsten, V., Lavorel, S. & Freitas, H. Functional response traits in relation
1119 to land use change in the Montado. *Agriculture Ecosystems & Environment* **137**, 183-
1120 191, doi:10.1016/j.agee.2010.02.002 (2010).
- 1121 96. Castro-Luna, A. A., Sosa, V. J. & Castillo-Campos, G. Bat diversity and abundance
1122 associated with the degree of secondary succession in a tropical forest mosaic in
1123 south-eastern Mexico. *Animal Conservation* **10**, 219-228, doi:10.1111/j.1469-
1124 1795.2007.00097.x (2007).
- 1125 97. Center For International Forestry Research (CIFOR). Multidisciplinary Landscape
1126 Assessment - Cameroon. (2013).
- 1127 98. Center For International Forestry Research (CIFOR). Multidisciplinary Landscape
1128 Assessment - Philippines. (2013).
- 1129 99. Centro Agronómico Tropical de Investigación y Enseñanza(CATIE). *Unpublished*
1130 *data of reptilian and amphibian diversity in six countries in Central America* (Centro
1131 Agronómico Tropical de Investigación y Enseñanza (CATIE), 2010).
- 1132 100. Cerezo, A., Conde, M. & Poggio, S. Pasture area and landscape heterogeneity are key
1133 determinants of bird diversity in intensively managed farmland. *Biodiversity and*
1134 *Conservation* **20**, 2649-2667, doi:10.1007/s10531-011-0096-y (2011).
- 1135 101. Chapman, K. & Reich, P. Land use and habitat gradients determine bird community
1136 diversity and abundance in suburban, rural and reserve landscapes of Minnesota,

- 1137 USA. *Biological Conservation* **135**, 527-541, doi:10.1016/j.biocon.2006.10.050
 1138 (2007).
- 1139 102. Chauvat, M., Wolters, V. & Dauber, J. Response of collembolan communities to land-
 1140 use change and grassland succession. *Ecography* **30**, 183-192,
 1141 doi:10.1111/j.2007.0906-7590.04888.x (2007).
- 1142 103. Clarke, F. M., Rostant, L. V. & Racey, P. A. Life after logging: post-logging recovery
 1143 of a neotropical bat community. *Journal of Applied Ecology* **42**, 409-420,
 1144 doi:10.1111/j.1365-2664.2005.01024.x (2005).
- 1145 104. Cleary, D. F. R. *et al.* Diversity and community composition of butterflies and
 1146 odonates in an ENSO-induced fire affected habitat mosaic: a case study from East
 1147 Kalimantan, Indonesia. *Oikos* **105**, 426-446, doi:10.1111/j.0030-1299.2004.12219.x
 1148 (2004).
- 1149 105. Cleary, D. F. R. & Mooers, A. O. Burning and logging differentially affect endemic
 1150 vs. widely distributed butterfly species in Borneo. *Diversity and Distributions* **12**,
 1151 409-416, doi:10.1111/j.1366-9516.2006.00256.x (2006).
- 1152 106. Cockle, K. L., Leonard, M. L. & Bodrati, A. A. Presence and abundance of birds in an
 1153 Atlantic forest reserve and adjacent plantation of shade-grown yerba mate, in
 1154 Paraguay. *Biodiversity and Conservation* **14**, 3265-3288, doi:10.1007/s10531-004-
 1155 0446-0 (2005).
- 1156 107. Connop, S., Hill, T., Steer, J. & Shaw, P. Microsatellite analysis reveals the spatial
 1157 dynamics of *Bombus humilis* and *Bombus sylvarum*. *Insect Conservation and*
 1158 *Diversity* **4**, 212-221, doi:10.1111/j.1752-4598.2010.00116.x (2011).
- 1159 108. D'Aniello, B., Stanislao, I., Bonelli, S. & Balletto, E. Haying and grazing effects on
 1160 the butterfly communities of two Mediterranean-area grasslands. *Biodiversity and*
 1161 *Conservation* **20**, 1731-1744, doi:10.1007/s10531-011-0058-4 (2011).
- 1162 109. Darvill, B., Knight, M. E. & Goulson, D. Use of genetic markers to quantify
 1163 bumblebee foraging range and nest density. *Oikos* **107**, 471-478, doi:10.1111/j.0030-
 1164 1299.2004.13510.x (2004).
- 1165 110. Davis, A. L. V. & Philips, T. K. Effect of deforestation on a southwest Ghana dung
 1166 beetle assemblage (Coleoptera : Scarabaeidae) at the periphery of Ankasa
 1167 conservation area. *Environmental Entomology* **34**, 1081-1088, doi:10.1603/0046-
 1168 225x(2005)034[1081:eodoas]2.0.co;2 (2005).
- 1169 111. Davis, E. S., Murray, T. E., Fitzpatrick, U., Brown, M. J. F. & Paxton, R. J.
 1170 Landscape effects on extremely fragmented populations of a rare solitary bee,
 1171 *Colletes floralis*. *Molecular Ecology* **19**, 4922-4935, doi:10.1111/j.1365-
 1172 294X.2010.04868.x (2010).
- 1173 112. Dawson, J. *et al.* Bird communities of the lower Waria Valley, Morobe Province,
 1174 Papua New Guinea: a comparison between habitat types. *Tropical Conservation*
 1175 *Science* **4**, 317-348 (2011).
- 1176 113. Delabie, J. H. C. *et al.* Ants as biological indicators of Wayana Amerindian land use
 1177 in French Guiana. *Comptes Rendus Biologies* **332**, 673-684,
 1178 doi:10.1016/j.crv.2009.01.006 (2009).
- 1179 114. Diekötter, T., Walther-Hellwig, K., Conradi, M., Suter, M. & Frankl, R. Effects of
 1180 landscape elements on the distribution of the rare bumblebee species *Bombus*
 1181 *muscorum* in an agricultural landscape. *Biodiversity and Conservation* **15**, 57-68,
 1182 doi:10.1007/s10531-004-2932-9 (2006).
- 1183 115. Domínguez, E., Bahamonde, N. & Muñoz-Escobar, C. Efectos de la extracción de
 1184 turba sobre la composición y estructura de una turbera de *Sphagnum* explotada y
 1185 abandonada hace 20 años, Chile. *Anales Instituto Patagonia (Chile)* **40**, 37-45,
 1186 doi:10.4067/S0718-686X2012000200003 (2012).

116. Dominguez-Haydar, Y. & Armbrrecht, I. Response of ants and their seed removal in rehabilitation areas and forests at El Cerrejon coal mine in Colombia. *Restoration Ecology* **19**, 178-184, doi:10.1111/j.1526-100X.2010.00735.x (2010).
117. Dumont, B. *et al.* How does grazing intensity influence the diversity of plants and insects in a species-rich upland grassland on basalt soils? *Grass and Forage Science* **64**, 92-105, doi:10.1111/j.1365-2494.2008.00674.x (2009).
118. Dures, S. G. & Cumming, G. S. The confounding influence of homogenising invasive species in a globally endangered and largely urban biome: Does habitat quality dominate avian biodiversity? *Biological Conservation* **143**, 768-777, doi:10.1016/j.biocon.2009.12.019 (2010).
119. Edenius, L., Mikusinski, G. & Bergh, J. Can Repeated Fertilizer Applications to Young Norway Spruce Enhance Avian Diversity in Intensively Managed Forests? *Ambio* **40**, 521-527, doi:10.1007/s13280-011-0137-5 (2011).
120. Elek, Z. & Lovei, G. L. Patterns in ground beetle (Coleoptera : Carabidae) assemblages along an urbanisation gradient in Denmark. *Acta Oecologica-International Journal of Ecology* **32**, 104-111, doi:10.1016/j.actao.2007.03.008 (2007).
121. Endo, W. *et al.* Game vertebrate densities in hunted and nonhunted forest sites in Manu National Park, Peru. *Biotropica* **42**, 251-261, doi:10.1111/j.1744-7429.2009.00546.x (2010).
122. Faruk, A., Belabut, D., Ahmad, N., Knell, R. J. & Garner, T. W. J. Effects of oil-palm plantations on diversity of tropical anurans. *Conservation Biology* **27**, 615-624, doi:10.1111/cobi.12062 (2013).
123. Farwig, N., Sajita, N. & Boehning-Gaese, K. Conservation value of forest plantations for bird communities in western Kenya. *Forest Ecology and Management* **255**, 3885-3892, doi:10.1016/j.foreco.2008.03.042 (2008).
124. Fayle, T. M. *et al.* Oil palm expansion into rain forest greatly reduces ant biodiversity in canopy, epiphytes and leaf-litter. *Basic and Applied Ecology* **11**, 337-345, doi:10.1016/j.baae.2009.12.009 (2010).
125. Felton, A. M., Engstrom, L. M., Felton, A. & Knott, C. D. Orangutan population density, forest structure and fruit availability in hand-logged and unlogged peat swamp forests in West Kalimantan, Indonesia. *Biological Conservation* **114**, 91-101, doi:10.1016/s0006-3207(03)00013-2 (2003).
126. Fensham, R., Dwyer, J., Eyre, T., Fairfax, R. & Wang, J. The effect of clearing on plant composition in mulga (*Acacia aneura*) dry forest, Australia. *Austral Ecology* **37**, 183-192, doi:10.1111/j.1442-9993.2011.02261.x (2012).
127. Fermon, H., Waltert, M., Vane-Wright, R. I. & Muhlenberg, M. Forest use and vertical stratification in fruit-feeding butterflies of Sulawesi, Indonesia: impacts for conservation. *Biodiversity and Conservation* **14**, 333-350, doi:10.1007/s10531-004-5054-9 (2005).
128. Ferreira, C. & Alves, P. C. Impacto da implementação de medidas de gestão do habitat nas populações de coelho-bravo (*Oryctolagus cuniculus algirus*) no Parque Natural do Sudoeste Alentejano e Costa Vicentina. (Centro de Investigação em Biodiversidade e Recursos Genéticos (CIBIO), 2005).
129. Fierro, M. M., Cruz-Lopez, L., Sanchez, D., Villanueva-Gutierrez, R. & Vandame, R. Effect of biotic factors on the spatial distribution of stingless bees (Hymenoptera: Apidae, Meliponini) in fragmented neotropical habitats. *Neotropical Entomology* **41**, 95-104, doi:10.1007/s13744-011-0009-5 (2012).

- 1235 130. Filgueiras, B., Iannuzzi, L. & Leal, I. Habitat fragmentation alters the structure of
1236 dung beetle communities in the Atlantic Forest. *Biological Conservation* **144**, 362-
1237 369, doi:10.1016/j.biocon.2010.09.013 (2011).
- 1238 131. Flaspohler, D. J. *et al.* Long-term effects of fragmentation and fragment properties on
1239 bird species richness in Hawaiian forests. *Biological Conservation* **143**, 280-288,
1240 doi:10.1016/j.biocon.2009.10.009 (2010).
- 1241 132. Fukuda, D., Tisen, O. B., Momose, K. & Sakai, S. Bat diversity in the vegetation
1242 mosaic around a lowland dipterocarp forest of Borneo. *Raffles Bulletin of Zoology* **57**,
1243 213-221 (2009).
- 1244 133. Furlani, D., Ficetola, G. F., Colombo, G., Ugurlucan, M. & De Bernardi, F.
1245 Deforestation and the structure of frog communities in the Humedale Terraba-Sierpe,
1246 Costa Rica. *Zoological Science* **26**, 197-202, doi:10.2108/zsj.26.197 (2009).
- 1247 134. Garden, J. G., McAlpine, C. A. & Possingham, H. P. Multi-scaled habitat
1248 considerations for conserving urban biodiversity: native reptiles and small mammals
1249 in Brisbane, Australia. *Landscape Ecology* **25**, 1013-1028, doi:10.1007/s10980-010-
1250 9476-z (2010).
- 1251 135. Gardner, T. A., Hernandez, M. I. M., Barlow, J. & Peres, C. A. Understanding the
1252 biodiversity consequences of habitat change: the value of secondary and plantation
1253 forests for neotropical dung beetles. *Journal of Applied Ecology* **45**, 883-893,
1254 doi:10.1111/j.1365-2664.2008.01454.x (2008).
- 1255 136. Gheler-Costa, C., Vettorazzi, C. A., Pardini, R. & Verdade, L. M. The distribution and
1256 abundance of small mammals in agroecosystems of southeastern Brazil.
1257 *Mammalia* **76**, 185-191, doi:10.1515/mammalia-2011-0109 (2012).
- 1258 137. Giordani, P. Assessing the effects of forest management on epiphytic lichens in
1259 coppiced forests using different indicators. *Plant Biosystems* **146**, 628-637,
1260 doi:10.1080/11263504.2011.654136 (2012).
- 1261 138. Giordano, S. *et al.* Biodiversity and trace element content of epiphytic bryophytes in
1262 urban and extraurban sites of southern Italy. *Plant Ecology* **170**, 1-14,
1263 doi:10.1023/B:VEGE.0000019025.36121.5d (2004).
- 1264 139. Golodets, C., Kigel, J. & Sternberg, M. Recovery of plant species composition and
1265 ecosystem function after cessation of grazing in a Mediterranean grassland. *Plant and*
1266 *Soil* **329**, 365-378, doi:10.1007/s11104-009-0164-1 (2010).
- 1267 140. Gottschalk, M. S., De Toni, D. C., Valente, V. L. S. & Hofmann, P. R. P. Changes in
1268 Brazilian Drosophilidae (Diptera) assemblages across an urbanisation gradient.
1269 *Neotropical Entomology* **36**, 848-862, doi:10.1590/S1519-566X2007000600005
1270 (2007).
- 1271 141. Goulson, D. *et al.* Effects of land use at a landscape scale on bumblebee nest density
1272 and survival. *Journal of Applied Ecology* **47**, 1207-1215, doi:10.1111/j.1365-
1273 2664.2010.01872.x (2010).
- 1274 142. Goulson, D., Lye, G. C. & Darvill, B. Diet breadth, coexistence and rarity in
1275 bumblebees. *Biodiversity and Conservation* **17**, 3269-3288, doi:10.1007/s10531-008-
1276 9428-y (2008).
- 1277 143. Gove, A. D., Majer, J. D. & Rico-Gray, V. Methods for conservation outside of
1278 formal reserve systems: The case of ants in the seasonally dry tropics of Veracruz,
1279 Mexico. *Biological Conservation* **126**, 328-338, doi:10.1016/j.biocon.2005.06.008
1280 (2005).
- 1281 144. Grogan, J. *et al.* What loggers leave behind: Impacts on big-leaf mahogany (*Swietenia*
1282 *macrophylla*) commercial populations and potential for post-logging recovery in the
1283 Brazilian Amazon. *Forest Ecology and Management* **255**, 269-281,
1284 doi:10.1016/j.foreco.2007.09.048 (2008).

- 1285 145. Gu, W.-B., Zhen-Rong, Y. & Dun-Xiao, H. Carabid community and its fluctuation in
1286 farmland of salinity transforming area in the North China Plain: a case study in
1287 Quzhou County, Hebei Province. *Biodiversity Science* **12**, 262-268 (2004).
- 1288 146. Gutierrez-Lamus, D. L. Composition and abundance of Anura in two forest types
1289 (natural and planted) in the eastern Cordillera of Colombia. *Caldasia* **26**, 245-264
1290 (2004).
- 1291 147. Hanley, M. E. *et al.* Increased bumblebee abundance along the margins of a mass
1292 flowering crop: evidence for pollinator spill-over. *Oikos* **120**, 1618-1624,
1293 doi:10.1111/j.1600-0706.2011.19233.x (2011).
- 1294 148. Hanley, M. E. *Unpublished data of bee diversity in UK croplands* (2005).
- 1295 149. Hanley, M. E. *Unpublished data of bee diversity in UK croplands and urban*
1296 *habitats* (2011).
- 1297 150. Hanson, T. R., Brunsfeld, S. J., Finegan, B. & Waits, L. P. Pollen dispersal and
1298 genetic structure of the tropical tree *Dipteryx panamensis* in a fragmented Costa Rican
1299 landscape. *Molecular Ecology* **17**, 2060-2073, doi:10.1111/j.1365-294X.2008.03726.x
1300 (2008).
- 1301 151. Hashim, N., Akmal, W., Jusoh, W. & Nasir, M. Ant diversity in a Peninsular
1302 Malaysian mangrove forest and oil palm plantation. *Asian Myrmecology* **3**, 5-8
1303 (2010).
- 1304 152. Hatfield, R. G. & LeBuhn, G. Patch and landscape factors shape community
1305 assemblage of bumble bees, *Bombus* spp. (Hymenoptera : Apidae), in montane
1306 meadows. *Biological Conservation* **139**, 150-158, doi:10.1016/j.biocon.2007.06.019
1307 (2007).
- 1308 153. Hawes, J. *et al.* Diversity and composition of Amazonian moths in primary, secondary
1309 and plantation forests. *Journal of Tropical Ecology* **25**, 281-300,
1310 doi:10.1017/s0266467409006038 (2009).
- 1311 154. Helden, A. J. & Leather, S. R. Biodiversity on urban roundabouts - Hemiptera,
1312 management and the species-area relationship. *Basic and Applied Ecology* **5**, 367-377,
1313 doi:10.1016/j.baae.2004.06.004 (2004).
- 1314 155. Hernández, L., Delgado, L., Meier, W. & Duran, C. Empobrecimiento de bosques
1315 fragmentados en el norte de la Gran Sabana, Venezuela. *Interciencia* **37**, 891-898
1316 (2012).
- 1317 156. Herrmann, F., Westphal, C., Moritz, R. F. A. & Steffan-Dewenter, I. Genetic diversity
1318 and mass resources promote colony size and forager densities of a social bee (*Bombus*
1319 *pascuorum*) in agricultural landscapes. *Molecular Ecology* **16**, 1167-1178,
1320 doi:10.1111/j.1365-294X.2007.03226.x (2007).
- 1321 157. Hietz, P. Conservation of vascular epiphyte diversity in Mexican coffee
1322 plantations. *Conservation Biology* **19**, 391-399, doi:10.1111/j.1523-
1323 1739.2005.00145.x (2005).
- 1324 158. Higuera, D. & Wolf, J. H. D. Vascular epiphytes in dry oak forests show resilience to
1325 anthropogenic disturbance, Cordillera Oriental, Colombia. *Caldasia* **32**, 161-174
1326 (2010).
- 1327 159. Hilje, B. & Aide, T. M. Recovery of amphibian species richness and composition in a
1328 chronosequence of secondary forests, northeastern Costa Rica. *Biological*
1329 *Conservation* **146**, 170-176, doi:10.1016/j.biocon.2011.12.007 (2012).
- 1330 160. Hoffmann, A. & Zeller, U. Influence of variations in land use intensity on species
1331 diversity and abundance of small mammals in the Nama Karoo, Namibia. *Belgian*
1332 *Journal of Zoology* **135**, 91-96 (2005).

- 1333 161. Horgan, F. G. Invasion and retreat: shifting assemblages of dung beetles amidst
1334 changing agricultural landscapes in central Peru. *Biodiversity and Conservation* **18**,
1335 3519-3541, doi:10.1007/s10531-009-9658-7 (2009).
- 1336 162. Hu, C. & Cao, Z. P. Nematode community structure under compost and chemical
1337 fertilizer management practice, in the north China plain. *Experimental Agriculture***44**,
1338 485-496, doi:10.1017/s0014479708006716 (2008).
- 1339 163. Hylander, K. & Weibull, H. Do time-lagged extinctions and colonizations change the
1340 interpretation of buffer strip effectiveness? - a study of riparian bryophytes in the first
1341 decade after logging. *Journal of Applied Ecology* **49**, 1316-1324, doi:10.1111/j.1365-
1342 2664.2012.02218.x (2012).
- 1343 164. Hylander, K. & Nemomissa, S. Complementary Roles of Home Gardens and Exotic
1344 Tree Plantations as Alternative Habitats for Plants of the Ethiopian Montane
1345 Rainforest. *Conservation Biology* **23**, 400-409, doi:10.1111/j.1523-
1346 1739.2008.01097.x (2009).
- 1347 165. Ims, R. A. & Henden, J. A. Collapse of an arctic bird community resulting from
1348 ungulate-induced loss of erect shrubs. *Biological Conservation* **149**, 2-5,
1349 doi:10.1016/j.biocon.2012.02.008 (2012).
- 1350 166. Isaacs Cubides, P. J. & Urbina Cardona, J. N. Anthropogenic disturbance and edge
1351 effects on anuran assemblages inhabiting cloud forest fragments in Colombia.
1352 *Natureza & Conservacao* **9**, 39-46, doi:10.4322/natcon.2011.004 (2011).
- 1353 167. Ishitani, M., Kotze, D. J. & Niemela, J. Changes in carabid beetle assemblages across
1354 an urban-rural gradient in Japan. *Ecography* **26**, 481-489, doi:10.1034/j.1600-
1355 0587.2003.03436.x (2003).
- 1356 168. Jacobs, C. T., Scholtz, C. H., Escobar, F. & Davis, A. L. V. How might intensification
1357 of farming influence dung beetle diversity (Coleoptera: Scarabaeidae) in Maputo
1358 Special Reserve (Mozambique)? *Journal of Insect Conservation* **14**, 389-399,
1359 doi:10.1007/s10841-010-9270-x (2010).
- 1360 169. Johnson, M. F., Gómez, A. & Pinedo-Vasquez, M. Land use and mosquito diversity in
1361 the Peruvian Amazon. *Journal of Medical Entomology* **45**, 1023-1030,
1362 doi:10.1603/0022-2585(2008)45[1023:luamdi]2.0.co;2 (2008).
- 1363 170. Jonsell, M. Old park trees as habitat for saproxylic beetle species. *Biodiversity and*
1364 *Conservation* **21**, 619-642, doi:10.1007/s10531-011-0203-0 (2012).
- 1365 171. Julier, H. E. & Roulston, T. a. H. Wild bee abundance and pollination service in
1366 cultivated pumpkins: farm management, nesting behavior and landscape effects.
1367 *Journal of Economic Entomology* **102**, 563-573, doi:10.1603/029.102.0214 (2009).
- 1368 172. Jung, T. S. & Powell, T. Spatial distribution of meadow jumping mice (*Zapus*
1369 *hudsonius*) in logged boreal forest of northwestern Canada. *Mammalian Biology* **76**,
1370 678-682, doi:10.1016/j.mambio.2011.08.002 (2011).
- 1371 173. Kapoor, V. Effects of rainforest fragmentation and shade-coffee plantations on spider
1372 communities in the Western Ghats, India. *Journal of Insect Conservation* **12**, 53-68,
1373 doi:10.1007/s10841-006-9062-5 (2008).
- 1374 174. Kappes, H., Katzschner, L. & Nowak, C. Urban summer heat load: meteorological
1375 data as a proxy for metropolitan biodiversity. *Meteorologische Zeitschrift* **21**, 525-
1376 528, doi:10.1127/0941-2948/2012/0361 (2012).
- 1377 175. Kati, V., Zografou, K., Tzirkalli, E., Chitos, T. & Willemse, L. Butterfly and
1378 grasshopper diversity patterns in humid Mediterranean grasslands: the roles of
1379 disturbance and environmental factors. *Journal of Insect Conservation* **16**, 807-818,
1380 doi:10.1007/s10841-012-9467-2 (2012).
- 1381 176. Katovai, E., Burley, A. L. & Mayfield, M. M. Understory plant species and functional
1382 diversity in the degraded wet tropical forests of Kolombangara Island, Solomon

- Islands. *Biological Conservation* **145**, 214-224, doi:10.1016/j.biocon.2011.11.008 (2012).
177. Kessler, M. *et al.* Tree diversity in primary forest and different land use systems in Central Sulawesi, Indonesia. *Biodiversity and Conservation* **14**, 547-560, doi:10.1007/s10531-004-3914-7 (2005).
178. Kessler, M. *et al.* Alpha and beta diversity of plants and animals along a tropical land-use gradient. *Ecological Applications* **19**, 2142-2156, doi:10.1890/08-1074.1 (2009).
179. Knight, M. E. *et al.* Bumblebee nest density and the scale of available forage in arable landscapes. *Insect Conservation and Diversity* **2**, 116-124, doi:10.1111/j.1752-4598.2009.00049.x (2009).
180. Knop, E., Ward, P. I. & Wich, S. A. A comparison of orang-utan density in a logged and unlogged forest on Sumatra. *Biological Conservation* **120**, 183-188, doi:10.1016/j.biocon.2004.02.010 (2004).
181. Kohler, F., Verhulst, J., van Klink, R. & Kleijn, D. At what spatial scale do high-quality habitats enhance the diversity of forbs and pollinators in intensively farmed landscapes? *Journal of Applied Ecology* **45**, 753-762, doi:10.1111/j.1365-2664.2007.01394.x (2008).
182. Koivula, M., Hyrylainen, V. & Soininen, E. Carabid beetles (Coleoptera: Carabidae) at forest-farmland edges in southern Finland. *Journal of Insect Conservation* **8**, 297-309, doi:10.1007/s10841-004-0296-9 (2004).
183. Kolb, A. & Diekmann, M. Effects of environment, habitat configuration and forest continuity on the distribution of forest plant species. *Journal of Vegetation Science* **15**, 199-208, doi:10.1111/j.1654-1103.2004.tb02255.x (2004).
184. Kőrösi, Á., Batáry, P., Orosz, A., Rédei, D. & Báldi, A. Effects of grazing, vegetation structure and landscape complexity on grassland leafhoppers (Hemiptera: Auchenorrhyncha) and true bugs (Hemiptera: Heteroptera) in Hungary. *Insect Conservation and Diversity* **5**, 57-66, doi:10.1111/j.1752-4598.2011.00153.x (2012).
185. Krauss, J., Klein, A. M., Steffan-Dewenter, I. & Tscharntke, T. Effects of habitat area, isolation, and landscape diversity on plant species richness of calcareous grasslands. *Biodiversity and Conservation* **13**, 1427-1439, doi:10.1023/b:bioc.0000021323.18165.58 (2004).
186. Krauss, J., Steffan-Dewenter, I. & Tscharntke, T. How does landscape context contribute to effects of habitat fragmentation on diversity and population density of butterflies? *Journal of Biogeography* **30**, 889-900, doi:10.1046/j.1365-2699.2003.00878.x (2003).
187. Kumar, R. & Shahabuddin, G. Effects of biomass extraction on vegetation structure, diversity and composition of forests in Sariska Tiger Reserve, India. *Environmental Conservation* **32**, 248-259, doi:10.1017/s0376892905002316 (2005).
188. Lachat, T. *et al.* Arthropod diversity in Lama forest reserve (South Benin), a mosaic of natural, degraded and plantation forests. *Biodiversity and Conservation* **15**, 3-23, doi:10.1007/s10531-004-1234-6 (2006).
189. Lantschner, M. V., Rusch, V. & Hayes, J. P. Habitat use by carnivores at different spatial scales in a plantation forest landscape in Patagonia, Argentina. *Forest Ecology and Management* **269**, 271-278, doi:10.1016/j.foreco.2011.12.045 (2012).
190. Lantschner, M. V., Rusch, V. & Peyrou, C. Bird assemblages in pine plantations replacing native ecosystems in NW Patagonia. *Biodiversity and Conservation* **17**, 969-989, doi:10.1007/s10531-007-9243-x (2008).
191. Latta, S. C., Tinoco, B. A., Astudillo, P. X. & Graham, C. H. Patterns and magnitude of temporal change in avian communities in the Ecuadorian Andes. *Condor* **113**, 24-40, doi:10.1525/cond.2011.090252 (2011).

- 1433 192. Légaré, J.-P., Hébert, C. & Ruel, J.-C. Alternative Silvicultural Practices in Irregular
1434 Boreal Forests: Response of Beetle Assemblages. *Silva Fennica* **45**, 937-956 (2011).
- 1435 193. Letcher, S. G. & Chazdon, R. L. Rapid recovery of biomass, species richness, and
1436 species composition in a forest chronosequence in northeastern Costa
1437 Rica. *Biotropica* **41**, 608-617, doi:10.1111/j.1744-7429.2009.00517.x (2009).
- 1438 194. Littlewood, N. A., Pakeman, R. J. & Pozsgai, G. Grazing impacts on
1439 Auchenorrhyncha diversity and abundance on a Scottish upland estate. *Insect*
1440 *Conservation and Diversity* **5**, 67-74, doi:10.1111/j.1752-4598.2011.00135.x (2012).
- 1441 195. Liu, Y. H., Axmacher, J. C., Wang, C. L., Li, L. T. & Yu, Z. R. Ground beetle
1442 (Coleoptera: Carabidae) assemblages of restored semi-natural habitats and intensively
1443 cultivated fields in northern China. *Restoration Ecology* **20**, 234-239,
1444 doi:10.1111/j.1526-100X.2010.00755.x (2012).
- 1445 196. Lo-Man-Hung, N. F., Gardner, T. A., Ribeiro-Júnior, M. A., Barlow, J. & Bonaldo, A.
1446 B. The value of primary, secondary, and plantation forests for Neotropical epigeic
1447 arachnids. *Journal of Arachnology* **36**, 394-401, doi:10.1636/CT07-136.1 (2008).
- 1448 197. López-Quintero, C. A., Straatsma, G., Franco-Molano, A. E. & Boekhout, T.
1449 Macrofungal diversity in Colombian Amazon forests varies with regions and regimes
1450 of disturbance. *Biodiversity Conservation*, 2221–2243, doi:10.1007/s10531-012-0280-
1451 8 (2012).
- 1452 198. Louhaichi, M., Salkini, A. K. & Petersen, S. L. Effect of small ruminant grazing on
1453 the plant community characteristics of semiarid Mediterranean ecosystems.
1454 *International Journal of Agriculture and Biology* **11**, 681-689 (2009).
- 1455 199. Lucas-Borja, M. E. *et al.* The effects of human trampling on the microbiological
1456 properties of soil and vegetation in Mediterranean mountain areas. *Land Degradation*
1457 *& Development* **22**, 383-394, doi:10.1002/ldr.1014 (2011).
- 1458 200. Luja, V., Herrando-Perez, S., Gonzalez-Solis, D. & Luiselli, L. Secondary rain forests
1459 are not havens for reptile species in tropical Mexico. *Biotropica* **40**, 747-757,
1460 doi:10.1111/j.1744-7429.2008.00439.x (2008).
- 1461 201. Luskin, M. S. Flying Foxes Prefer to Forage in Farmland in a Tropical Dry Forest
1462 Landscape Mosaic in Fiji. *Biotropica* **42**, 246-250, doi:10.1111/j.1744-
1463 7429.2009.00577.x (2010).
- 1464 202. MacSwiney, M. C. G., Vilchis, P. L., Clarke, F. M. & Racey, P. A. The importance of
1465 cenotes in conserving bat assemblages in the Yucatan, Mexico. *Biological*
1466 *Conservation* **136**, 499-509, doi:10.1016/j.biocon.2006.12.021 (2007).
- 1467 203. Maeto, K. & Sato, S. Impacts of forestry on ant species richness and composition in
1468 warm-temperate forests of Japan. *Forest Ecology and Management* **187**, 213-223,
1469 doi:10.1016/s0378-1127(03)00333-5 (2004).
- 1470 204. Magura, T., Horvath, R. & Tothmeresz, B. Effects of urbanization on ground-
1471 dwelling spiders in forest patches, in Hungary. *Landscape Ecology* **25**, 621-629,
1472 doi:10.1007/s10980-009-9445-6 (2010).
- 1473 205. Mallari, N. A. D. *et al.* Population densities of understorey birds across a habitat
1474 gradient in Palawan, Philippines: implications for conservation. *Oryx* **45**, 234-242,
1475 doi:10.1017/s0030605310001031 (2011).
- 1476 206. Malone, L. *et al.* Observations on bee species visiting white clover in New Zealand
1477 pastures. *Journal of Apicultural Research* **49**, 284-286, doi:10.3896/ibra.1.49.3.09
1478 (2010).
- 1479 207. Marin-Spiotta, E., Ostertag, R. & Silver, W. L. Long-term patterns in tropical
1480 reforestation: Plant community composition and aboveground biomass accumulation.
1481 *Ecological Applications* **17**, 828-839, doi:10.1890/06-1268 (2007).

- 1482 208. Marshall, E. J. P., West, T. M. & Kleijn, D. Impacts of an agri-environment field
1483 margin prescription on the flora and fauna of arable farmland in different
1484 landscapes. *Agriculture Ecosystems & Environment* **113**, 36-44,
1485 doi:10.1016/j.agee.2005.08.036 (2006).
- 1486 209. Martin, P. S., Gheler-Costa, C., Lopes, P. C., Rosalino, L. M. & Verdade, L. M.
1487 Terrestrial non-volant small mammals in agro-silvicultural landscapes of Southeastern
1488 Brazil. *Forest Ecology and Management* **282**, 185-195,
1489 doi:10.1016/j.foreco.2012.07.002 (2012).
- 1490 210. Matsumoto, T., Itioka, T., Yamane, S. & Momose, K. Traditional land use associated
1491 with swidden agriculture changes encounter rates of the top predator, the army ant, in
1492 Southeast Asian tropical rain forests. *Biodiversity and Conservation* **18**, 3139-3151,
1493 doi:10.1007/s10531-009-9632-4 (2009).
- 1494 211. Mayfield, M. M., Ackerly, D. & Daily, G. C. The diversity and conservation of plant
1495 reproductive and dispersal functional traits in human-dominated tropical
1496 landscapes. *Journal of Ecology* **94**, 522-536, doi:10.1111/j.1365-2745.2006.01108.x
1497 (2006).
- 1498 212. McFrederick, Q. S. & LeBuhn, G. Are urban parks refuges for bumble bees *Bombus*
1499 spp. (Hymenoptera : Apidae)? *Biological Conservation* **129**, 372-382,
1500 doi:10.1016/j.biocon.2005.11.004 (2006).
- 1501 213. McNamara, S., Erskine, P. D., Lamb, D., Chantalangsy, L. & Boyle, S. Primary tree
1502 species diversity in secondary fallow forests of Laos. *Forest Ecology and*
1503 *Management* **281**, 93-99, doi:10.1016/j.foreco.2012.06.004 (2012).
- 1504 214. Meyer, B., Gaebele, V. & Steffan-Dewenter, I. D. Patch size and landscape effects on
1505 pollinators and seed set of the horseshoe vetch, *hippocrepis comosa*, in an agricultural
1506 landscape of central Europe. *Entomologia Generalis* **30**, 173-185 (2007).
- 1507 215. Meyer, B., Jauker, F. & Steffan-Dewenter, I. Contrasting resource-dependent
1508 responses of hoverfly richness and density to landscape structure. *Basic and Applied*
1509 *Ecology* **10**, 178-186, doi:10.1016/j.baae.2008.01.001 (2009).
- 1510 216. Mico, E., Garcia-Lopez, A., Brustel, H., Padilla, A. & Galante, E. Explaining the
1511 saproxylic beetle diversity of a protected Mediterranean area. *Biodiversity and*
1512 *Conservation* **22**, 889-904, doi:10.1007/s10531-013-0456-x (2013).
- 1513 217. Milder, J. C. *et al.* Effects of farm and landscape management on bird and butterfly
1514 conservation in western Honduras. *Ecosphere* **1**, art2, doi:10.1890/es10-00003.1
1515 (2010).
- 1516 218. Miranda, M. V., Politi, N. & Rivera, L. O. Unexpected changes in the bird assemblage
1517 in areas under selective logging in piedmont forest in northwestern Argentina.
1518 *Ornitologia Neotropical* **21**, 323-337 (2010).
- 1519 219. Moreno-Mateos, D. *et al.* Effects of land use on nocturnal birds in a Mediterranean
1520 agricultural landscape. *Acta Ornithologica* **46**, 173-182,
1521 doi:10.3161/000164511x625946 (2011).
- 1522 220. Muchane, M. N. *et al.* Land use practices and their implications on soil macro-fauna
1523 in Maasai Mara ecosystem. *International Journal of Biodiversity and Conservation* **4**,
1524 500-514, doi:10.5897/IJBC12.030 (2012).
- 1525 221. Mudri-Stojnic, S., Andric, A., Jozan, Z. & Vujic, A. Pollinator diversity
1526 (Hymenoptera and Diptera) in semi-natural habitats in Serbia during
1527 summer. *Archives of Biological Sciences* **64**, 777-786, doi:10.2298/abs1202777s
1528 (2012).
- 1529 222. Naidoo, R. Species richness and community composition of songbirds in a tropical
1530 forest-agricultural landscape. *Animal Conservation* **7**, 93-105,
1531 doi:10.1017/s1367943003001185 (2004).

- 1532 223. Nakamura, A., Proctor, H. & Catterall, C. P. Using soil and litter arthropods to assess
1533 the state of rainforest restoration. *Ecological Management & Restoration* **4**, S20-S28,
1534 doi:10.1046/j.1442-8903.4.s.3.x (2003).
- 1535 224. Naoe, S., Sakai, S. & Masaki, T. Effect of forest shape on habitat selection of birds in
1536 a plantation-dominant landscape across seasons: comparison between continuous and
1537 strip forests. *Journal of Forest Research* **17**, 219-223, doi:10.1007/s10310-011-0296-
1538 z (2012).
- 1539 225. Navarrete, D. & Halffter, G. Dung beetle (Coleoptera : Scarabaeidae : Scarabaeinae)
1540 diversity in continuous forest, forest fragments and cattle pastures in a landscape of
1541 Chiapas, Mexico: the effects of anthropogenic changes. *Biodiversity and*
1542 *Conservation* **17**, 2869-2898, doi:10.1007/s10531-008-9402-8 (2008).
- 1543 226. Navarro, I. L., Roman, A. K., Gomez, F. H. & Perez, H. A. Seasonal variation in dung
1544 beetles (Coleoptera: Scarabaeidae: Scarabaeinae) from Serrania de Coraza, Sucre
1545 (Colombia). *Revista Colombiana de Ciencia Animal* **3**, 102-110 (2011).
- 1546 227. Neuschulz, E. L., Botzat, A. & Farwig, N. Effects of forest modification on bird
1547 community composition and seed removal in a heterogeneous landscape in South
1548 Africa. *Oikos* **120**, 1371-1379, doi:10.1111/j.1600-0706.2011.19097.x (2011).
- 1549 228. Nicolas, V., Barriere, P., Tapiero, A. & Colyn, M. Shrew species diversity and
1550 abundance in Ziam Biosphere Reserve, Guinea: comparison among primary forest,
1551 degraded forest and restoration plots. *Biodiversity and Conservation* **18**, 2043-2061,
1552 doi:10.1007/s10531-008-9572-4 (2009).
- 1553 229. Nielsen, A. *et al.* Assessing bee species richness in two Mediterranean communities:
1554 importance of habitat type and sampling techniques. *Ecological Research* **26**, 969-
1555 983, doi:10.1007/s11284-011-0852-1 (2011).
- 1556 230. Noreika, N. & Kotze, D. J. Forest edge contrasts have a predictable effect on the
1557 spatial distribution of carabid beetles in urban forests. *Journal of Insect*
1558 *Conservation* **16**, 867-881, doi:10.1007/s10841-012-9474-3 (2012).
- 1559 231. Noreika, N. New records of rare species of Coleoptera found in Ukmergė district in
1560 2004-2005. *New and Rare for Lithuania Insect Species* **21**, 68-71 (2009).
- 1561 232. Norfolk, O., Abdel-Dayem, M. & Gilbert, F. Rainwater harvesting and arthropod
1562 biodiversity within an arid agro-ecosystem. *Agriculture Ecosystems &*
1563 *Environment* **162**, 8-14, doi:10.1016/j.agee.2012.08.007 (2012).
- 1564 233. Noriega, J. A., Realpe, E. & Fagua, G. Diversidad de escarabajos coprofagos
1565 (Coleoptera: Scarabaeidae) en un bosque de galeria con tres estadios de alteracion.
1566 *Universitas Scientiarum* **12**, 51-63 (2007).
- 1567 234. Noriega, J. A., Palacio, J. M., Monroy-G, J. D. & Valencia, E. Estructura de un
1568 ensamblaje de escarabajos coprofagos (Coleoptera: Scarabaeinae) en tres sitios con
1569 diferente uso del suelo en Antioquia, Colombia. *Actualidades Biologicas*
1570 *(Medellin)* **34**, 43-54 (2012).
- 1571 235. Nöske, N. M. *et al.* Disturbance effects on diversity of epiphytes and moths in a
1572 montane forest in Ecuador. *Basic and Applied Ecology* **9**, 4-12,
1573 doi:10.1016/j.baae.2007.06.014 (2008).
- 1574 236. Numa, C., Verdu, J. R., Rueda, C. & Galante, E. Comparing dung beetle species
1575 assemblages between protected areas and adjacent pasturelands in a Mediterranean
1576 savanna landscape. *Rangeland Ecology & Management* **65**, 137-143,
1577 doi:10.2111/rem-d-10-00050.1 (2012).
- 1578 237. O'Connor, T. G. Influence of land use on plant community composition and diversity
1579 in Highland Sourveld grassland in the southern Drakensberg, South Africa. *Journal of*
1580 *Applied Ecology* **42**, 975-988, doi:10.1111/j.1365-2664.2005.01065.x (2005).

- 1581 238. O'Dea, N. & Whittaker, R. J. How resilient are Andean montane forest bird
1582 communities to habitat degradation? *Biodiversity and Conservation* **16**, 1131-1159,
1583 doi:10.1007/s10531-006-9095-9 (2007).
- 1584 239. Ofori-Boateng, C. *et al.* Differences in the effects of selective logging on amphibian
1585 assemblages in three west African forest types. *Biotropica* **45**, 94-101,
1586 doi:10.1111/j.1744-7429.2012.00887.x (2013).
- 1587 240. Oke, C. Land snail diversity in post extraction secondary forest reserves in Edo State,
1588 Nigeria. *African Journal of Ecology* **51**, 244-254, doi:10.1111/aje.12029 (2013).
- 1589 241. Oke, O. C. & Chokor, J. U. The effect of land use on snail species richness and
1590 diversity in the tropical rainforest of south-western Nigeria. *African Scientist* **10**, 95-
1591 108 (2009).
- 1592 242. Oliveira, D. E., Carrijo, T. F. & Brandão, D. Species composition of termites
1593 (Isoptera) in different Cerrado vegetation physiognomies. *Sociobiology* **60**, 190-197,
1594 doi:10.13102/sociobiology.v60i2.190-197 (2013).
- 1595 243. Osgathorpe, L. M., Park, K. & Goulson, D. The use of off-farm habitats by foraging
1596 bumblebees in agricultural landscapes: implications for conservation
1597 management. *Apidologie* **43**, 113-127, doi:10.1007/s13592-011-0083-z (2012).
- 1598 244. Otavo, S. E., Parrado-Rosselli, A. & Noriega, J. A. Scarabaeoidea superfamily
1599 (Insecta: Coleoptera) as a bioindicator element of anthropogenic disturbance in an
1600 amazon national park. *Revista De Biologia Tropical* **61**, 735-752 (2013).
- 1601 245. Otto, C. R. V. & Roloff, G. J. Songbird response to green-tree retention prescriptions
1602 in clearcut forests. *Forest Ecology and Management* **284**, 241-250,
1603 doi:10.1016/j.foreco.2012.07.016 (2012).
- 1604 246. Paradis, S. & Work, T. T. Partial cutting does not maintain spider assemblages within
1605 the observed range of natural variability in Eastern Canadian black spruce
1606 forests. *Forest Ecology and Management* **262**, 2079-2093,
1607 doi:10.1016/j.foreco.2011.08.032 (2011).
- 1608 247. Paritsis, J. & Aizen, M. A. Effects of exotic conifer plantations on the biodiversity of
1609 understory plants, epigeal beetles and birds in *Nothofagus dombeyi* forests. *Forest
1610 Ecology and Management* **255**, 1575-1583, doi:10.1016/j.foreco.2007.11.015 (2008).
- 1611 248. Parra-H, A. & Nates-Parra, G. Variation of the orchid bees community (Hymenoptera
1612 : Apidae) in three altered habitats of the Colombian "llano" piedmont. *Revista De
1613 Biologia Tropical* **55**, 931-941 (2007).
- 1614 249. Pelegrin, N. & Bucher, E. H. Effects of habitat degradation on the lizard assemblage
1615 in the Arid Chaco, central Argentina. *Journal of Arid Environments* **79**, 13-19,
1616 doi:10.1016/j.jaridenv.2011.11.004 (2012).
- 1617 250. Phalan, B., Onial, M., Balmford, A. & Green, R. Reconciling food production and
1618 biodiversity conservation: land sharing and land sparing compared. *Science* **333**,
1619 1289-1291, doi:10.1126/science.1208742 (2011).
- 1620 251. Pillsbury, F. C. & Miller, J. R. Habitat and landscape characteristics underlying
1621 anuran community structure along an urban-rural gradient. *Ecological
1622 Applications* **18**, 1107-1118, doi:10.1890/07-1899.1 (2008).
- 1623 252. Pineda, E. & Halffter, G. Species diversity and habitat fragmentation: frogs in a
1624 tropical montane landscape in Mexico. *Biological Conservation* **117**, 499-508,
1625 doi:10.1016/j.biocon.2003.08.009 (2004).
- 1626 253. Politi, N., Hunter, M., Jr. & Rivera, L. Assessing the effects of selective logging on
1627 birds in Neotropical piedmont and cloud montane forests. *Biodiversity and
1628 Conservation* **21**, 3131-3155, doi:10.1007/s10531-012-0358-3 (2012).
- 1629 254. Poveda, K., Martinez, E., Kersch-Becker, M., Bonilla, M. & Tschardtke, T.
1630 Landscape simplification and altitude affect biodiversity, herbivory and Andean

- potato yield. *Journal of Applied Ecology* **49**, 513-522, doi:10.1111/j.1365-2664.2012.02120.x (2012).
255. Power, E. F., Kelly, D. L. & Stout, J. C. Organic farming and landscape structure: effects on insect-pollinated plant diversity in intensively managed grasslands. *Plos One* **7**, doi:10.1371/journal.pone.0038073 (2012).
 256. Power, E. F. & Stout, J. C. Organic dairy farming: impacts on insect-flower interaction networks and pollination. *Journal of Applied Ecology* **48**, 561-569, doi:10.1111/j.1365-2664.2010.01949.x (2011).
 257. Presley, S. J., Willig, M. R., Wunderle, J. M., Jr. & Saldanha, L. N. Effects of reduced-impact logging and forest physiognomy on bat populations of lowland Amazonian forest. *Journal of Applied Ecology* **45**, 14-25, doi:10.1111/j.1365-2664.2007.01373.x (2008).
 258. Proenca, V. M., Pereira, H. M., Guilherme, J. & Vicente, L. Plant and bird diversity in natural forests and in native and exotic plantations in NW Portugal. *Acta Oecologica-International Journal of Ecology* **36**, 219-226, doi:10.1016/j.actao.2010.01.002 (2010).
 259. Quaranta, M. *et al.* Wild bees in agroecosystems and semi-natural landscapes. 1997-2000 collection period in Italy. *Bulletin of Insectology* **57**, 11-62 (2004).
 260. Quintero, C., Laura Morales, C. & Adrian Aizen, M. Effects of anthropogenic habitat disturbance on local pollinator diversity and species turnover across a precipitation gradient. *Biodiversity and Conservation* **19**, 257-274, doi:10.1007/s10531-009-9720-5 (2010).
 261. Redpath, N., Osgathorpe, L. M., Park, K. & Goulson, D. Crofting and bumblebee conservation: The impact of land management practices on bumblebee populations in northwest Scotland. *Biological Conservation* **143**, 492-500, doi:10.1016/j.biocon.2009.11.019 (2010).
 262. Reid, J. L., Harris, J. B. C. & Zahawi, R. A. Avian habitat preference in tropical forest restoration in southern Costa Rica. *Biotropica* **44**, 350-359, doi:10.1111/j.1744-7429.2011.00814.x (2012).
 263. Reis, Y. T. & Cancellato, E. M. Termite (Insecta, Isoptera) richness in primary and secondary Atlantic Forest in southeastern Bahia. *Iheringia Serie Zoologia* **97**, 229-234 (2007).
 264. Rey-Velasco, J. C. & Miranda-Esquivel, D. R. *Habitat modification in Andean forest: the response of ground beetles (Coleoptera: Carabidae) on the northeastern Colombian Andes* (2012).
 265. Ribeiro, D. B. & Freitas, A. V. L. The effect of reduced-impact logging on fruit-feeding butterflies in Central Amazon, Brazil. *Journal of Insect Conservation* **16**, 733-744, doi:10.1007/s10841-012-9458-3 (2012).
 266. Richardson, B. A., Richardson, M. J. & Soto-Adames, F. N. Separating the effects of forest type and elevation on the diversity of litter invertebrate communities in a humid tropical forest in Puerto Rico. *Journal of Animal Ecology* **74**, 926-936, doi:10.1111/j.1365-2656.2005.00990.x (2005).
 267. Robles, C. A., Carmaran, C. C. & Lopez, S. E. Screening of xylophagous fungi associated with *Platanus acerifolia* in urban landscapes: Biodiversity and potential biodeterioration. *Landscape and Urban Planning* **100**, 129-135, doi:10.1016/j.landurbplan.2010.12.003 (2011).
 268. Rodrigues, M. M., Uchoa, M. A. & Ide, S. Dung beetles (Coleoptera: Scarabaeoidea) in three landscapes in Mato Grosso do Sul, Brazil. *Brazilian Journal of Biology* **73**, 211-220 (2013).

- 1680 269. Römbke, J., Schmidt, P. & Höfer, H. The earthworm fauna of regenerating forests and
1681 anthropogenic habitats in the coastal region of Paraná. *Pesquisa Agropecuaria*
1682 *Brasileira* **44**, 1040-1049, doi:10.1590/S0100-204X2009000800037 (2009).
- 1683 270. Romero-Duque, L. P., Jaramillo, V. J. & Perez-Jimenez, A. Structure and diversity of
1684 secondary tropical dry forests in Mexico, differing in their prior land-use history.
1685 *Forest Ecology and Management* **253**, 38-47, doi:10.1016/j.foreco.2007.07.002
1686 (2007).
- 1687 271. Rosselli, L. *Factores ambientales relacionados con la presencia y abundancia de las*
1688 *aves de los humedales de la Sabana de Bogotá* PhD thesis, Universidad Nacional de
1689 Colombia, (2011).
- 1690 272. Rousseau, L., Fonte, S. J., Tellez, O., van der Hoek, R. & Lavelle, P. Soil macrofauna
1691 as indicators of soil quality and land use impacts in smallholder agroecosystems of
1692 western Nicaragua. *Ecological Indicators* **27**, 71-82,
1693 doi:10.1016/j.ecolind.2012.11.020 (2013).
- 1694 273. Safian, S., Csontos, G. & Winkler, D. Butterfly community recovery in degraded
1695 rainforest habitats in the Upper Guinean Forest Zone (Kakum forest, Ghana). *Journal*
1696 *of Insect Conservation* **15**, 351-359, doi:10.1007/s10841-010-9343-x (2011).
- 1697 274. Sakchoowong, W., Nomura, S., Ogata, K. & Chanpaisaeng, J. Diversity of pselaphine
1698 beetles (Coleoptera : Staphylinidae : Pselaphinae) in eastern Thailand. *Entomological*
1699 *Science* **11**, 301-313, doi:10.1111/j.1479-8298.2008.00281.x (2008).
- 1700 275. Saldana-Vazquez, R. A., Sosa, V. J., Hernandez-Montero, J. R. & Lopez-Barrera, F.
1701 Abundance responses of frugivorous bats (Stenodermatinae) to coffee cultivation and
1702 selective logging practices in mountainous central Veracruz, Mexico. *Biodiversity and*
1703 *Conservation* **19**, 2111-2124, doi:10.1007/s10531-010-9829-6 (2010).
- 1704 276. Samnegård, U., Persson, A. S. & Smith, H. G. Gardens benefit bees and enhance
1705 pollination in intensively managed farmland. *Biological Conservation* **144**, 2602-
1706 2606, doi:10.1016/j.biocon.2011.07.008 (2011).
- 1707 277. Santana, J., Porto, M., Gordinho, L., Reino, L. & Beja, P. Long-term responses of
1708 Mediterranean birds to forest fuel management. *Journal of Applied Ecology* **49**, 632-
1709 643, doi:10.1111/j.1365-2664.2012.02141.x (2012).
- 1710 278. Savage, J., Wheeler, T. A., Moores, A. M. A. & Taillefer, A. G. Effects of habitat
1711 size, vegetation cover, and surrounding land use on diptera diversity in temperate
1712 nearctic bogs. *Wetlands* **31**, 125-134, doi:10.1007/s13157-010-0133-8 (2011).
- 1713 279. Schmidt, A. C., Fraser, L. H., Carlyle, C. N. & Bassett, E. R. L. Does cattle grazing
1714 affect ant abundance and diversity in temperate grasslands? *Rangeland Ecology &*
1715 *Management* **65**, 292-298, doi:10.2111/rem-d-11-00100.1 (2012).
- 1716 280. Schon, N. L., Mackay, A. D. & Minor, M. A. Soil fauna in sheep-grazed hill pastures
1717 under organic and conventional livestock management and in an adjacent ungrazed
1718 pasture. *Pedobiologia* **54**, 161-168, doi:10.1016/j.pedobi.2011.01.001 (2011).
- 1719 281. Schüepp, C., Herrmann, J. D., Herzog, F. & Schmidt-Entling, M. H. Differential
1720 effects of habitat isolation and landscape composition on wasps, bees, and their
1721 enemies. *Oecologia* **165**, 713-721, doi:10.1007/s00442-010-1746-6 (2011).
- 1722 282. Schüepp, C., Rittiner, S. & Entling, M. H. High Bee and Wasp Diversity in a
1723 Heterogeneous Tropical Farming System Compared to Protected Forest. *Plos One* **7**,
1724 doi:10.1371/journal.pone.0052109 (2012).
- 1725 283. Scott, D. M. *et al.* The impacts of forest clearance on lizard, small mammal and bird
1726 communities in the arid spiny forest, southern Madagascar. *Biological*
1727 *Conservation* **127**, 72-87, doi:10.1016/j.biocon.2005.07.014 (2006).

- 1728 284. Sedlock, J. L. *et al.* Bat diversity in tropical forest and agro-pastoral habitats within a
 1729 protected area in the Philippines. *Acta Chiropterologica* **10**, 349-358,
 1730 doi:10.3161/150811008x414926 (2008).
- 1731 285. Shafie, N. J., Sah, S. A. M., Latip, N. S. A., Azman, N. M. & Khairuddin, N. L.
 1732 Diversity pattern of bats at two contrasting habitat types along Kerian River, Perak,
 1733 Malaysia. *Tropical Life Sciences Research* **22**, 13-22 (2011).
- 1734 286. Shahabuddin, G. & Kumar, R. Effects of extractive disturbance on bird assemblages,
 1735 vegetation structure and floristics in tropical scrub forest, Sariska Tiger Reserve,
 1736 India. *Forest Ecology and Management* **246**, 175-185,
 1737 doi:10.1016/j.foreco.2007.03.061 (2007).
- 1738 287. Sheil, D. *et al.* Exploring biological diversity, environment and local people's
 1739 perspectives in forest landscapes: Methods for a multidisciplinary landscape
 1740 assessment. (Center for International Forestry Research (CIFOR), Jakarta, 2002).
- 1741 288. Sheldon, F., Styring, A. & Hosner, P. Bird species richness in a Bornean exotic tree
 1742 plantation: A long-term perspective. *Biological Conservation* **143**, 399-407,
 1743 doi:10.1016/j.biocon.2009.11.004 (2010).
- 1744 289. Shuler, R. E., Roulston, T. H. & Farris, G. E. Farming practices influence wild
 1745 pollinator populations on squash and pumpkin. *Journal of Economic Entomology* **98**,
 1746 790-795, doi:10.1603/0022-0493-98.3.790 (2005).
- 1747 290. Silva, F. A. B., Costa, C. M. Q., Moura, R. C. & Farias, A. I. Study of the dung beetle
 1748 (Coleoptera: Scarabaeidae) community at two sites: atlantic forest and clear-cut,
 1749 Pernambuco, Brazil. *Environmental Entomology* **39**, 359-367, doi:10.1603/en09180
 1750 (2010).
- 1751 291. da Silva, P. G. *Espécies de Scarabaeinae (Coleoptera: Scarabaeidae) de fragmentos*
 1752 *florestais com diferentes níveis de alteração em Santa Maria, Rio Grande do Sul* MSc
 1753 thesis, Universidade Federal de Santa Maria, (2011).
- 1754 292. Slade, E. M., Mann, D. J. & Lewis, O. T. Biodiversity and ecosystem function of
 1755 tropical forest dung beetles under contrasting logging regimes. *Biological*
 1756 *Conservation* **144**, 166-174, doi:10.1016/j.biocon.2010.08.011 (2011).
- 1757 293. Smith-Pardo, A. & Gonzalez, V. H. Bee diversity (Hymenoptera : Apoidea) in a
 1758 tropical rainforest succession. *Acta Biologica Colombiana* **12**, 43-55 (2007).
- 1759 294. Sodhi, N. S. *et al.* Deforestation and avian extinction on tropical landbridge
 1760 islands. *Conservation Biology* **24**, 1290-1298, doi:10.1111/j.1523-1739.2010.01495.x
 1761 (2010).
- 1762 295. Sosa, R. A., Benz, V. A., Galea, J. M. & Poggio Herrero, I. V. Efecto del grado de
 1763 disturbio sobre el ensamble de aves en la reserva provincial Parque Luro, La Pampa,
 1764 Argentina. *Revista de la Asociación Argentina de Ecología de Paisajes* **1**, 101-110
 1765 (2010).
- 1766 296. de Souza, V. M., de Souza, B. & Morato, E. F. Effect of the forest succession on the
 1767 anurans (Amphibia: Anura) of the Reserve Catuaba and its periphery, Acre,
 1768 southwestern Amazonia. *Revista Brasileira De Zoologia* **25**, 49-57 (2008).
- 1769 297. Sridhar, H., Raman, T. R. S. & Mudappa, D. Mammal persistence and abundance in
 1770 tropical rainforest remnants in the southern Western Ghats, India. *Current Science* **94**,
 1771 748-757 (2008).
- 1772 298. St-Laurent, M. H., Ferron, J., Hins, C. & Gagnon, R. Effects of stand structure and
 1773 landscape characteristics an habitat use by birds and small mammals in managed
 1774 boreal forest of eastern Canada. *Canadian Journal of Forest Research-Revue*
 1775 *Canadienne De Recherche Forestiere* **37**, 1298-1309, doi:10.1139/xo6-295 (2007).

- 1776 299. Ström, L., Hylander, K. & Dynesius, M. Different long-term and short-term responses
1777 of land snails to clear-cutting of boreal stream-side forests. *Biological*
1778 *Conservation* **142**, 1580-1587, doi:10.1016/j.biocon.2009.02.028 (2009).
- 1779 300. Struebig, M. J., Kingston, T., Zubaid, A., Mohd-Adnan, A. & Rossiter, S. J.
1780 Conservation value of forest fragments to Palaeotropical bats. *Biological*
1781 *Conservation* **141**, 2112-2126, doi:10.1016/j.biocon.2008.06.009 (2008).
- 1782 301. Su, Z. M., Zhang, R. Z. & Qiu, J. X. Decline in the diversity of willow trunk-dwelling
1783 weevils (Coleoptera: Curculionoidea) as a result of urban expansion in Beijing,
1784 China. *Journal of Insect Conservation* **15**, 367-377, doi:10.1007/s10841-010-9310-6
1785 (2011).
- 1786 302. Sugiura, S., Tsuru, T., Yamaura, Y. & Makihara, H. Small off-shore islands can serve
1787 as important refuges for endemic beetle conservation. *Journal of Insect*
1788 *Conservation* **13**, 377-385, doi:10.1007/s10841-008-9185-y (2009).
- 1789 303. Summerville, K. S. Managing the forest for more than the trees: effects of
1790 experimental timber harvest on forest Lepidoptera. *Ecological Applications* **21**, 806-
1791 816, doi:10.1890/10-0715.1 (2011).
- 1792 304. Summerville, K. S., Conoan, C. J. & Steichen, R. M. Species traits as predictors of
1793 lepidopteran composition in restored and remnant tallgrass prairies. *Ecological*
1794 *Applications* **16**, 891-900, doi:10.1890/1051-0761(2006)016[0891:stapol]2.0.co;2
1795 (2006).
- 1796 305. Sung, Y. H., Karraker, N. E. & Hau, B. C. H. Terrestrial herpetofaunal assemblages in
1797 secondary forests and exotic *Lophostemon confertus* plantations in South
1798 China. *Forest Ecology and Management* **270**, 71-77,
1799 doi:10.1016/j.foreco.2012.01.011 (2012).
- 1800 306. Threlfall, C. G., Law, B. & Banks, P. B. Sensitivity of insectivorous bats to
1801 urbanization: Implications for suburban conservation planning. *Biological*
1802 *Conservation* **146**, 41-52, doi:10.1016/j.biocon.2011.11.026 (2012).
- 1803 307. Tonietto, R., Fant, J., Ascher, J., Ellis, K. & Larkin, D. A comparison of bee
1804 communities of Chicago green roofs, parks and prairies. *Landscape and Urban*
1805 *Planning* **103**, 102-108, doi:10.1016/j.landurbplan.2011.07.004 (2011).
- 1806 308. Turner, E. C. & Foster, W. A. The impact of forest conversion to oil palm on
1807 arthropod abundance and biomass in Sabah, Malaysia. *Journal of Tropical*
1808 *Ecology* **25**, 23-30, doi:10.1017/s0266467408005658 (2009).
- 1809 309. Tylianakis, J. M., Klein, A. M. & Tscharntke, T. Spatiotemporal variation in the
1810 diversity of hymenoptera across a tropical habitat gradient. *Ecology* **86**, 3296-3302,
1811 doi:10.1890/05-0371 (2005).
- 1812 310. Vanbergen, A. J., Woodcock, B. A., Watt, A. D. & Niemela, J. Effect of land-use
1813 heterogeneity on carabid communities at the landscape scale. *Ecography* **28**, 3-16,
1814 doi:10.1111/j.0906-7590.2005.03991.x (2005).
- 1815 311. Vassilev, K., Pedashenko, H., Nikolov, S. C., Apostolova, I. & Dengler, J. Effect of
1816 land abandonment on the vegetation of upland semi-natural grasslands in the Western
1817 Balkan Mts., Bulgaria. *Plant Biosystems* **145**, 654-665,
1818 doi:10.1080/11263504.2011.601337 (2011).
- 1819 312. Vázquez, D. P. & Simberloff, D. Ecological specialization and susceptibility to
1820 disturbance: Conjectures and refutations. *American Naturalist* **159**, 606-623,
1821 doi:10.1086/339991 (2002).
- 1822 313. Verboven, H. A. F., Brys, R. & Hermy, M. Sex in the city: Reproductive success of
1823 *Digitalis purpurea* in a gradient from urban to rural sites. *Landscape and Urban*
1824 *Planning* **106**, 158-164, doi:10.1016/j.landurbplan.2012.02.015 (2012).

- 1825 314. Verdasca, M. J. *et al.* Forest fuel management as a conservation tool for early
1826 successional species under agricultural abandonment: The case of Mediterranean
1827 butterflies. *Biological Conservation* **146**, 14-23, doi:10.1016/j.biocon.2011.10.031
1828 (2012).
- 1829 315. Verdú, J. R. *et al.* Grazing promotes dung beetle diversity in the xeric landscape of a
1830 Mexican Biosphere Reserve. *Biological Conservation* **140**, 308-317,
1831 doi:10.1016/j.biocon.2007.08.015 (2007).
- 1832 316. Vergara, C. H. & Badano, E. I. Pollinator diversity increases fruit production in
1833 Mexican coffee plantations: The importance of rustic management systems.
1834 *Agriculture Ecosystems & Environment* **129**, 117-123,
1835 doi:10.1016/j.agee.2008.08.001 (2009).
- 1836 317. Vergara, P. M. & Simonetti, J. A. Avian responses to fragmentation of the Maulino
1837 Forest in central Chile. *Oryx* **38**, 383-388, doi:10.1017/s0030605304000742 (2004).
- 1838 318. Walker, T. R., Crittenden, P. D., Young, S. D. & Prystina, T. An assessment of
1839 pollution impacts due to the oil and gas industries in the Pechora basin, north-eastern
1840 European Russia. *Ecological Indicators* **6**, 369-387,
1841 doi:10.1016/j.ecolind.2005.03.015 (2006).
- 1842 319. Wang, Y., Bao, Y., Yu, M., Xu, G. & Ding, P. Nestedness for different reasons: the
1843 distributions of birds, lizards and small mammals on islands of an inundated lake.
1844 *Diversity and Distributions* **16**, 862-873, doi:10.1111/j.1472-4642.2010.00682.x
1845 (2010).
- 1846 320. Watling, J. I., Gerow, K. & Donnelly, M. A. Nested species subsets of amphibians
1847 and reptiles on Neotropical forest islands. *Animal Conservation* **12**, 467-476,
1848 doi:10.1111/j.1469-1795.2009.00274.x (2009).
- 1849 321. Weller, B. & Ganzhorn, J. U. Carabid beetle community composition, body size, and
1850 fluctuating asymmetry along an urban-rural gradient. *Basic and Applied Ecology* **5**,
1851 193-201, doi:10.1078/1439-1791-00220 (2004).
- 1852 322. Wells, K., Kalko, E. K. V., Lakim, M. B. & Pfeiffer, M. Effects of rain forest logging
1853 on species richness and assemblage composition of small mammals in Southeast
1854 Asia. *Journal of Biogeography* **34**, 1087-1099, doi:10.1111/j.1365-
1855 2699.2006.01677.x (2007).
- 1856 323. Williams, C. D., Sheahan, J. & Gormally, M. J. Hydrology and management of
1857 turloughs (temporary lakes) affect marsh fly (Sciomyzidae: Diptera) communities.
1858 *Insect Conservation and Diversity* **2**, 270-283, doi:10.1111/j.1752-4598.2009.00064.x
1859 (2009).
- 1860 324. Willig, M. R. *et al.* Phyllostomid bats of lowland Amazonia: Effects of habitat
1861 alteration on abundance. *Biotropica* **39**, 737-746, doi:10.1111/j.1744-
1862 7429.2007.00322.x (2007).
- 1863 325. Winfree, R., Griswold, T. & Kremen, C. Effect of human disturbance on bee
1864 communities in a forested ecosystem. *Conservation Biology* **21**, 213-223,
1865 doi:10.1111/j.1523-1739.2006.00574.x (2007).
- 1866 326. Woinarski, J. C. Z. *et al.* Fauna assemblages in regrowth vegetation in tropical open
1867 forests of the Northern Territory, Australia. *Wildlife Research* **36**, 675-690,
1868 doi:10.1071/wr08128 (2009).
- 1869 327. Woodcock, B. A. *et al.* The potential of grass field margin management for enhancing
1870 beetle diversity in intensive livestock farms. *Journal of Applied Ecology* **44**, 60-69,
1871 doi:10.1111/j.1365-2664.2006.01258.x (2007).
- 1872 328. Wunderle, J. M., Henriques, L. M. P. & Willig, M. R. Short-term responses of birds to
1873 forest gaps and understory: An assessment of reduced-impact logging in a lowland

- 1874 Amazon forest. *Biotropica* **38**, 235-255, doi:10.1111/j.1744-7429.2006.00138.x
 1875 (2006).
- 1876 329. Yoshikura, S., Yasui, S. & Kamijo, T. Comparative study of forest-dwelling bats'
 1877 abundances and species richness between old-growth forests and conifer plantations
 1878 in Nikko National Park, central Japan. *Mammal Study* **36**, 189-198,
 1879 doi:10.3106/041.036.0402 (2011).
- 1880 330. Zaitsev, A. S., Chauvat, M., Pflug, A. & Wolters, V. Oribatid mite diversity and
 1881 community dynamics in a spruce chronosequence. *Soil Biology & Biochemistry* **34**,
 1882 1919-1927, doi:10.1016/s0038-0717(02)00208-0 (2002).
- 1883 331. Zaitsev, A. S., Wolters, V., Waldhardt, R. & Dauber, J. Long-term succession of
 1884 oribatid mites after conversion of croplands to grasslands. *Applied Soil Ecology* **34**,
 1885 230-239, doi:10.1016/j.apsoil.2006.01.005 (2006).
- 1886 332. Zimmerman, G., Bell, F. W., Woodcock, J., Palmer, A. & Paloniemi, J. Response of
 1887 breeding songbirds to vegetation management in conifer plantations established in
 1888 boreal mixedwoods. *The Forestry Chronicle* **87**, 217-224 (2011).
- 1889 333. Roskov, Y. *et al.* *Species 2000 & ITIS Catalogue of Life, 2013 Annual Checklist*.
 1890 (2013).
- 1891 334. Gotelli, N. J. & Colwell, R. K. Quantifying biodiversity: procedures and pitfalls in the
 1892 measurement and comparison of species richness. *Ecol. Lett.* **4**, 379-391 (2001).
- 1893 335. Violle, C. *et al.* Let the concept of trait be functional! *Oikos* **116**, 882-892 (2007).
- 1894 336. Kattge, J. *et al.* TRY - a global database of plant traits. *Glob. Chang. Biol.* **17**, 2905-
 1895 2935 (2011).
- 1896 337. Jones, K. E. *et al.* PanTHERIA: a species-level database of life history, ecology, and
 1897 geography of extant and recently extinct mammals. *Ecology* **90**, 2648 (2009).
- 1898 338. Cooper, N., Bielby, J., Thomas, G. H. & Purvis, A. Macroecology and extinction risk
 1899 correlates of frogs. *Glob. Ecol. Biogeogr.* **17**, 211-221 (2008).
- 1900 339. AmphibiaWeb. <http://amphibiaweb.org/> (2013).
- 1901 340. Sunyer, J., Páiz, G., Dehling, D. M. & Köhler, G. A collection of amphibians from
 1902 Río San Juan, southeastern Nicaragua. *Herpetol. Notes* **2**, 189-202 (2009).
- 1903 341. Zug, G. R. & Zug, P. B. The marine toad *Bufo marinus*: a natural history resumé of
 1904 native populations. *Smithson. Contrib. to Zool.* **284**, 1-58 (1979).
- 1905 342. Amphibians & Reptiles of Peninsular Malaysia. <http://www.amphibia.my/> (2009)
- 1906 343. Shahriza, S., Ibrahim, H. J. & Shahrul Anuar, M. S. The correlation between total
 1907 rainfall and breeding parameters of white-lipped frog, *Rana labialis* (Anura: Ranidae)
 1908 in Kedah, Malaysia. *Trop. Nat. Hist.* **10**, 131-139 (2010).
- 1909 344. Bain, R. H. & Quang Truong, N. Three new species of narrow-mouth frogs (Genus:
 1910 *Microhyla* from Indochina, with comments on *Microhyla annamensis* and *Microhyla*
 1911 *palmipes*. *Copeia* **2004**, 507-524 (2004).
- 1912 345. Su, M.-Y., Kam, Y.-C. & Fellers, G. M. Effectiveness of amphibian monitoring
 1913 techniques in a Taiwanese subtropical forest. *Herpetol. J.* **15**, 73-79 (2005).
- 1914 346. Matson, T. O. A morphometric comparison of gray treefrogs, *Hyla chrysoscelis* and
 1915 *H. versicolor*, from Ohio. *Ohio J. Sci.* **90**, 98-101 (1990).
- 1916 347. Ningombam, B. & Bordoloi, S. Amphibian fauna of Loktak Lake, Manipur, India
 1917 with ten new records for the state. *Zoos' Print J.* **22**, 2688-2690 (2007).
- 1918 348. Lance, S. L. & Wells, K. D. Are spring peeper satellite males physiologically inferior
 1919 to calling males? *Copeia* **1993**, 1162-1166 (1993).
- 1920 349. Da Silva, E. T., dos Reis, E. P., Feio, R. N. & Filho, O. P. R. Diet of the invasive frog
 1921 *Lithobates catesbeianus* (Shaw, 1802) (Anura: Ranidae) in Viçosa, Minas Gerais
 1922 State, Brazil. *South Am. J. Herpetol.* **4**, 286-294 (2009).

- 1923 350. Blomquist, S. M. & Hunter Jr., M. L. A multi-scale assessment of habitat selection
1924 and movement patterns by northern leopard frogs (*Lithobates [Rana] pipiens*) in a
1925 managed forest. *Herpetol. Conserv. Biol.* **4**, 142–160 (2009).
- 1926 351. Caramaschi, U. & da Cruz, C. A. G. Redescription of *Chiasmocleis albopunctata*
1927 (Boettger) and description of a new species of *Chiasmocleis* (Anura: Microhylidae).
1928 *Herpetologica* **53**, 259–268 (1997).
- 1929 352. Brasileiro, C. A., Sawaya, R. J., Kiefer, M. C. & Martins, M. Amphibians of an open
1930 cerrado fragment in southeastern Brazil. *Biota Neotrop.* **5**, BN00405022005 (2005).
- 1931 353. De Almeida Prado, C. P. Estratégias reprodutivas em uma comunidade de anuros no
1932 pantanal, estado de Mato Grosso do Sul, Brasil. (PhD Thesis, Universidade Estadual
1933 Paulista, 2003).
- 1934 354. De Almeida Prado, C. P., Uetanabaro, M. & Lopes, F. S. Reproductive strategies of
1935 *Leptodactylus chaquensis* and *L. podicipinus* in the Pantanal, Brazil. *J. Herpetol.* **34**,
1936 135–139 (2000).
- 1937 355. De Carvalho, T. R., Giaretta, A. A. & Facure, K. G. A new species of *Hypsiboas*
1938 Wagler (Anura: Hylidae) closely related to *H. multifasciatus* Günther from
1939 southeastern Brazil. *Zootaxa* **2521**, 37–52 (2010).
- 1940 356. Heyer, W. R. & Heyer, M. M. *Leptodactylus elenae* Heyer. *Cat. Am. Amphib. Reptil.*
1941 742.1–742.5 (2002).
- 1942 357. Heyer, W. R. Variation within the *Leptodactylus podicipinus-wagneri* complex of
1943 frogs (Amphibia: Leptodactylidae). *Smithson. Contrib. to Zool.* **546**, (1994).
- 1944 358. Jungfer, K.-H. & Hödl, W. A new species of *Osteocephalus* from Ecuador and a
1945 redescription of *O. lepreurii* (Dumeril & Bibron, 1841) (Anura: Hylidae). *Amphibia-*
1946 *Reptilia* **23**, 21–46 (2002).
- 1947 359. Fouquet, A., Gaucher, P., Blanc, M. & Velez-Rodriguez, C. M. Description of two
1948 new species of *Rhinella* (Anura: Bufonidae) from the lowlands of the Guiana shield.
1949 *Zootaxa* **1663**, 17–32 (2007).
- 1950 360. Lynch, J. D. A review of the leptodactylid frogs of the genus *Pseudopaludicola* in
1951 Northern South America. *Copeia* **1989**, 577–588 (1989).
- 1952 361. González, C. E. & Hamann, M. I. Nematode parasites of two anuran species *Rhinella*
1953 *schneideri* (Bufonidae) and *Scinax acuminatus* (Hylidae) from Corrientes, Argentina.
1954 *Rev. Biol. Trop.* **56**, 2147–2161 (2008).
- 1955 362. Pombal Jr., J. P., Bilate, M., Gambale, P. G., Signorelli, L. & Bastos, R. P. A new
1956 miniature treefrog of the *Scinax ruber* clade from the cerrado of central Brazil (Anura:
1957 Hylidae). *Herpetologica* **67**, 288–299 (2011).
- 1958 363. Ibáñez, R., Jaramillo, C. A. & Solis, F. A. Description of the advertisement call of a
1959 species without vocal sac: *Craugastor gollmeri* (Amphibia: Craugastoridae). *Zootaxa*
1960 **3184**, 67–68 (2012).
- 1961 364. Hertz, A., Hauenschild, F., Lotzkat, S. & Köhler, G. A new golden frog species of the
1962 genus *Diasporus* (Amphibia, Eleutherodactylidae) from the Cordillera Central,
1963 western Panama. *Zookeys* **196**, 23–46 (2012).
- 1964 365. Goldberg, S. R. & Bursey, C. R. Helminths from fifteen species of frogs (Anura,
1965 Hylidae) from Costa Rica. *Phyllomedusa* **7**, 25–33 (2008).
- 1966 366. Bennett, W. O., Summers, A. P. & Brainerd, E. L. Confirmation of the passive
1967 exhalation hypothesis for a terrestrial caecilian, *Dermophis mexicanus*. *Copeia* **1999**,
1968 206–209 (1999).
- 1969 367. Anderson, M. T. & Mathis, A. Diets of two sympatric Neotropical salamanders,
1970 *Bolitoglossa mexicana* and *B. rufescens*, with notes on reproduction for *B. rufescens*.
1971 *J. Herpetol.* **33**, 601–607 (1999).

- 1972 368. McCranie, J. R. & Wilson, L. D. Taxonomic changes associated with the names *Hyla*
1973 spinipollex Schmidt and *Ptychohyla merazi* Wilson and McCranie (Anura: Hylidae).
1974 *Southwest. Nat.* **38**, 100–104 (1993).
- 1975 369. Barrio-Amorós, C. L., Guayasamin, J. M. & Hedges, S. B. A new minute Andean
1976 *Pristimantis* (Anura: Strabomantidae) from Venezuela. *Phyllomedusa* **11**, 83–93
1977 (2012).
- 1978 370. Arroyo, S. B., Serrano-Cardozo, V. H. & Ramírez-Pinilla, M. P. Diet, microhabitat
1979 and time of activity in a *Pristimantis* (Anura, Strabomantidae) assemblage.
1980 *Phyllomedusa* **7**, 109–119 (2008).
- 1981 371. Savage, J. M. & Myers, C. Frogs of the *Eleutherodactylus biporcatus* group
1982 (Leptodactylidae) of Central America and northern South America, including
1983 rediscovered, resurrected, and new taxa. *Am. Museum Novit.* **3357**, 1–48 (2002).
- 1984 372. Simões, P. I. Diversificação do complexo *Allobates femoralis* (Anura, Dendrobatidae)
1985 em florestas da Amazônia brasileira: desvendando padrões atuais e históricos. (PhD
1986 Thesis, Instituto Nacional de Pesquisas da Amazônia, 2010).
- 1987 373. Guayasamin, J. M., Ron, S. R., Cisneros-Heredia, D. F., Lamar, W. & McCracken, S.
1988 F. A new species of frog of the *Eleutherodactylus lacrimosus* assemblage
1989 (Leptodactylidae) from the western Amazon Basin, with comments on the utility of
1990 canopy surveys in lowland rainforest. *Herpetologica* **62**, 191–202 (2006).
- 1991 374. Jared, C., Antoniazzi, M. M., Verdade, V. K. & Toledo, L. F. The Amazonian toad
1992 *Rhaebo guttatus* is able to voluntarily squirt poison from the paratoid macroglands.
1993 *Amphibia-Reptilia* **32**, 546–549 (2011).
- 1994 375. Wollenberg, K. C., Veith, M., Noonan, B. P. & Lötters, S. Polymorphism versus
1995 species richness—systematics of large *Dendrobates* from the eastern Guiana Shield
1996 (Amphibia: Dendrobatidae). *Copeia* **2006**, 623–629 (2006).
- 1997 376. Shepard, D. B. & Caldwell, J. P. From foam to free-living: ecology of larval
1998 *Leptodactylus labyrinthicus*. *Copeia* **2005**, 803–811 (2005).
- 1999 377. Heyer, W. R., García-Lopez, J. M. & Cardoso, A. J. Advertisement call variation in
2000 the *Leptodactylus mystaceus* species complex (Amphibia: Leptodactylidae) with a
2001 description of a new sibling species. *Amphibia-Reptilia* **17**, 7–31 (1996).
- 2002 378. Zimmermann, B. L. A comparison of structural features of calls of open and forest
2003 habitat frog species in the central Amazon. *Herpetologica* **39**, 235–246 (1983).
- 2004 379. Bernarde, P. S. & Kokubum, M. N. D. C. Seasonality, age structure and reproduction
2005 of *Leptodactylus (Lithodytes) lineatus* (Anura, Leptodactylidae) in Rondônia state,
2006 southwestern Amazon, Brazil. *Iheringia Série Zool.* **99**, 368–372 (2009).
- 2007 380. Campbell, J. A. & Clarke, B. T. A review of frogs of the genus *Otophryne*
2008 (Microhylidae) with the description of a new species. *Herpetologica* **54**, 301–317
2009 (1998).
- 2010 381. Kan, F. W. Population dynamics, diet and morphological variation of the Hong Kong
2011 newt (*Paramesotriton hongkongensis*). (MPhil Thesis, The University of Hong Kong,
2012 2010).
- 2013 382. Stuart, B. L., Chuaynkern, Y., Chan-ard, T. & Inger, R. F. Three new species of frogs
2014 and a new tadpole from eastern Thailand. *Fieldiana Zool. New Ser.* **111**, 1–19 (2006).
- 2015 383. Ao, J. M., Bordoloi, S. & Ohler, A. Amphibian fauna of Nagaland with nineteen new
2016 records from the state including five new records for India. *Zoos' Print J.* **18**, 1117–
2017 1125 (2003).
- 2018 384. Ohler, A. *et al.* Sorting out *Lalos*: description of new species and additional
2019 taxonomic data on megophryid frogs from northern Indochina (genus *Leptolalax*,
2020 Megophryidae, Anura). *Zootaxa* **3147**, 1–83 (2011).

- 2021 385. Meiri, S. Evolution and ecology of lizard body sizes. *Glob. Ecol. Biogeogr.* **17**, 724–
2022 734 (2008).
- 2023 386. Itescu, Y., Karraker, N. E., Raia, P., Pritchard, P. C. H. & Meiri, S. Is the island rule
2024 general? Turtles disagree. *Glob. Ecol. Biogeogr.* **23**, 689–700 (2014).
- 2025 387. Meiri, S. Length-weight allometries in lizards. *J. Zool.* **281**, 218–226 (2010).
- 2026 388. Feldman, A. & Meiri, S. Length-mass allometry in snakes. *Biol. J. Linn. Soc.* **108**,
2027 161–172 (2013).
- 2028 389. Edgar, M. What can we learn from body length? A study in Coleoptera. (MRes
2029 Thesis, Imperial College London, 2014).
- 2030 390. Gilbert, F., Rotheray, G. E., Zafar, R. & Emerson, P. in *Phylogenetics Ecol.* 324–343
2031 (Academic Press, 1994).
- 2032 391. ESRI. ArcGIS Desktop: Release 10. (2011).
- 2033 392. R Core Team. *R: A Language and Environment for Statistical Computing.* (R
2034 Foundation for Statistical Computing, 2013). <http://www.r-project.org>
- 2035 393. Zuur, A. F., Ieno, E. N., Walker, N. J., Saveliev, A. A. & Smith, G. M. *Mixed Effects*
2036 *Models and Extensions in Ecology with R.* (Springer, 2009).
- 2037 394. Rigby, R. A., Stasinopoulos, D. M. & Akantziliotou, C. A framework for modelling
2038 overdispersed count data, including the Poisson-shifted generalized inverse Gaussian
2039 distribution. *Comput. Stat. Data Anal.* **53**, 381–393 (2008).
- 2040 395. Bivand, R. spdep: Spatial dependence: weighting schemes, statistics and models. R
2041 Package Version 0.5-68. (2013). <http://cran.r-project.org/web/packages/spdep>
- 2042 396. Møller, A. P. & Jennions, M. D. Testing and adjusting for publication bias. *Trends*
2043 *Ecol. Evol.* **16**, 580–586 (2001).
- 2044 397. Van Vuuren, D. P. *et al.* The representative concentration pathways: an overview.
2045 *Clim. Change* **109**, 5–31 (2011).
- 2046 398. United Nations Population Division. World Population Prospects: The 2010 Revision
2047 Population Database. *World Popul. Prospect. 2010 Revis. Popul. Database* (2011).
2048 <http://www.un.org/esa/population/>
- 2049 399. Van Asselen, S. & Verburg, P. H. Land cover change or land-use intensification:
2050 simulating land system change with a global-scale land change model. *Glob. Chang.*
2051 *Biol.* **19**, 3648–67 (2013).
- 2052 400. Haberl, H. *et al.* Quantifying and mapping the human appropriation of net primary
2053 production in earth's terrestrial ecosystems. *Proc. Natl. Acad. Sci. U. S. A.* **104**,
2054 12942–12947 (2007).
- 2055 401. Hijmans, R. J. raster: Geographic data analysis and modeling. (2014). [http://cran.r-](http://cran.r-project.org/package=raster)
2056 [project.org/package=raster](http://cran.r-project.org/package=raster)
- 2057 402. Olson, D. M. *et al.* Terrestrial ecoregions of the world: a new map of life on Earth.
2058 *Bioscience* **51**, 933–938 (2001).

a **b** **c** **d**