Mean univariate- GARCH VaR portfolio optimization: actual portfolio approach

Article (Accepted Version)

Rankovic, Vladimir, Drenovak, Mikica, Urosevic, Branko and Jelic, Ranko (2016) Mean univariate- GARCH VaR portfolio optimization: actual portfolio approach. *Computers & Operations Research*, 72. pp. 83-92. ISSN 0305-0548

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/59655/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Mean Univariate- GARCH VaR portfolio
optimization: Actual portfolio approach

Vladimir Ranković, Mikica Drenovak, Branko
Urošević, Ranko Jelić

www.elsevier.com/locate/caor

PII: S0305-0548(16)30004-1
DOI: <http://dx.doi.org/10.1016/j.cor.2016.01.014>
Reference: CAOR3923

To appear in: *Computers and Operation Research*

Received date: 10 July 2015
Revised date: 3 December 2015
Accepted date: 24 January 2016

Cite this article as: Vladimir Ranković, Mikica Drenovak, Branko Urošević and Ranko Jelić, Mean Univariate- GARCH VaR portfolio optimization: Actual portfolio approach, *Computers and Operation Research* <http://dx.doi.org/10.1016/j.cor.2016.01.014>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Mean-Univariate GARCH VaR portfolio optimization: Actual portfolio approach

Vladimir Ranković^{a)}, Mikica Drenovak^{b)}, Branko Urošević^{c)}, and Ranko Jelić^{d)}*

^{a)} Faculty of Economics, University of Kragujevac, Djure Pucara 3, 34000 Kragujevac, Serbia;

E-mail: vladimir.rankovich@gmail.com.

^{b)} Faculty of Economics, University of Kragujevac, Djure Pucara 3, 34000 Kragujevac, Serbia;

E-mail: mikicadrenovak@gmail.com.

^{c)} Faculty of Economics, University of Belgrade, Kamenicka 6, 11000 Beograd, Serbia; National Bank of Serbia, Kralja Petra 12, 11000 Beograd, Serbia; CESifo Network, Poschinger Str. 5, 81679, Munich, Germany;

E-mail: urosevic@ekof.bg.ac.rs.

*Corresponding author:

^{d)} School of Business, Management and Economics, University of Sussex, Sussex House, Falmer, Brighton, BN1 9RH UK;

E-mail: r.jelic@sussex.ac.uk, Phone: +44 (0)121 4145990

Abstract

In accordance with Basel Capital Accords, the Capital Requirements (*CR*) for market risk exposure of banks is a nonlinear function of Value-at-Risk (*VaR*). Importantly, the *CR* is calculated based on a bank's actual portfolio, i.e. the portfolio represented by its current holdings. To tackle mean-*VaR* portfolio optimization within the actual portfolio framework (*APF*), we propose a novel mean-*VaR* optimization method where *VaR* is estimated using a univariate Generalized AutoRegressive Conditional Heteroscedasticity (*GARCH*) volatility model. The optimization was performed by employing a Nondominated Sorting Genetic Algorithm (*NSGA-II*). On a sample of 40 large US stocks, our procedure provided superior mean-*VaR* trade-offs compared to those obtained from applying more customary mean-multivariate *GARCH* and historical *VaR* models. The results hold true in both low and high volatility samples.

KEY WORDS: *Portfolio optimization, Actual portfolios, Value at Risk, GARCH, NSGA-II*

1. Introduction

Value-at-Risk (VaR) is defined as the loss associated with the low (typically first or fifth) percentile of the return distribution.¹ The Basel II Capital Accord codifies VaR as the de-facto industry standard for the banking and insurance industries alike (see, BIS [3, 4, 5]). In particular, for the market risk exposure of banks, a bank's internal VaR estimates corresponding to the actual portfolio, i.e. the portfolio represented by its current holdings, translate directly into the regulatory capital charge (see Hendricks and Hirtle [6]). Motivated by this regulatory feature, we utilized the actual portfolio framework (APF) to determine a set of portfolios characterized by the optimal trade-off between the expected return and VaR (i.e. Pareto-optimal frontier). We further proposed a mean-univariate Generalized AutoRegressive Conditional Heteroscedasticity ($GARCH$) VaR portfolio optimization model. We assumed that portfolio returns, standardized by time varying volatility, have a conditional Student's t distribution, while conditional variance follows a $GARCH(1, 1)$ process.² The Student's t distribution efficiently captures the fat tails of standardized asset returns (see Christoffersen [9], Huisman et al. [10]) whilst the $GARCH$ model addresses issues related to volatility clustering observed in the data.³ To the best of our knowledge, this is the first paper that studies mean- VaR portfolio optimization using the actual portfolio approach and also the first paper that uses the univariate $GARCH$ VaR model in this context.

Previous studies on mean- VaR optimization implicitly assumed fixed weights (i.e. fractions of assets) over the observed time period.⁴ Since prices change over time, maintaining the fixed portfolio weights (FWA) requires frequent trading and leads to changes in the number of shares of each asset in a portfolio. Regulatory capital charges, however, are determined by the VaR of an actual portfolio where the number of shares of an asset (rather than its weighting) is fixed over the observed period. It is, therefore, APF , rather than FWA , that is more relevant for asset managers facing regulatory VaR limits. To illustrate the effectiveness of our APF approach and univariate $GARCH$ model, we compared our results with two benchmarks. Our first benchmark is the mean-historical VaR approach developed in Rockafellar and Uryasev [20, 21] and Krokmal et al. [22]. These authors mapped conditional VaR ($CVaR$) optimization into a linear programming problem and argued that the mean- $CVaR$ efficient

¹ For classification and comparison of risk measures see [1] and [2], among others.

² $GARCH$ was introduced in Engle [7] and $GARCH(1, 1)$ in Bollerslev [8].

³ For more on superiority of $GARCH$ VaR compared to historical VaR see [11], [12], [13], [14], [15], among others.

⁴ For example, [16], [17], [18], [19], [20], [21].

frontier provided near-optimal solutions in the context of the mean-historical *VaR* trade-off.⁵ We referred to this benchmark as the Linear Programming (*LP*) model. The second point of comparison for our approach is the mean-multivariate *GARCH VaR* optimization that can be mapped into the Quadratic Programming (*QP*) problem (see Santos et al. [26]).

Use of the univariate *GARCH* approach for *VaR* modeling, however, makes the mean-*VaR* optimization problem rather complex. Previous literature documented that Multi-Objective Evolutionary Algorithms (*MOEA*) could reliably and efficiently be applied in complex portfolio optimization problems.⁶ For example, Anagnostopoulos and Mamanis [31] studied the effectiveness of different *MOEA* (e.g. Nondominated Sorting Genetic Algorithm (*NSGA-II*), Strength Pareto Evolutionary Algorithm (*SPEA2*), Pareto Envelope-based Selection Algorithm (*PESA*), etc.) in solving various complex mean-variance optimization problems. They reported the best *NSGA-II* and *SPEA2* average performance in terms of hypervolume indicators, while *PESA* performed best in terms of the proximity to the Pareto-optimal frontier. The same authors (see [32]) also examined the mean-variance, mean-*ES*, and mean-*VaR* optimization problem with quantity, cardinality and class constraints. They showed that *NSGA-II*, *SPEA2* and *PESA* performed efficiently and their performance was independent of the risk measure used.⁷ Deb et al. [34] reported *NSGA-II*'s advantages over the Pareto Archived Evolutionary Strategy (*PAES*) and *SPEA*. Deb et al. [35] developed a hybrid *NSGA-II* procedure for handling a mean-variance portfolio optimization problem with the cardinality constraint and lower and upper bounds as investment criteria. The authors provided evidence of *NSGA-II*'s superiority over classical quadratic programming approaches. Branke et al. [36] considered the mean-variance problem with the maximum exposure constraint. The authors generated mean-variance Pareto-optimal frontiers by using a hybrid algorithm that combined *NSGA-II* with the critical line algorithm.

In this study we used the *NSGA-II* algorithm. Our choice was motivated by the above studies, whose results highlighted the advantages of *NSGA-II* in tackling complex portfolio

⁵ *CVaR* (or Expected Shortfall, *ES*) is the expected loss, conditional that loss is higher than *VaR*. Thus, *CVaR* and *VaR* are closely related risk measures. The *LP* model is widely used in *CVaR* and *VaR* optimization literature (see [23], [24], [25], among others).

⁶ For comprehensive surveys of *MOEA* applications in portfolio optimization, see [27], [28], [29], [30].

⁷ Anagnostopoulos and Mamanis also showed that *NSGA-II*, *SPEA2*, and *PESA* provide a good approximation of risk-return trade-offs in a 3 objectives optimization problem (see [33]).

optimization problems.⁸ *NSGA-II* was first introduced by Deb et al. [34] and subsequently developed in Deb et al. [37]. Recently, a new version of *NSGA* (*NSGA-III*) was developed by Deb and Jain [38] and also Jain and Deb [39]. *NSGA-III* is better suited to many-objective optimization problems (three and more objectives) and offers the ability to define the desired part of a solution space defined by reference points.⁹ Here, we prefer *NSGA-II* to *NSGA-III*, since we consider only two objectives and are interested in the entire Pareto-optimal frontier. Knowledge of the entire frontier of the risk-return trade-offs is particularly useful for asset managers in banks and other financial institutions in order to comply with different in-house and regulatory requirements.

For our numerical tests we selected 40 of the largest US stocks in the Standard and Poor's stock market index (*S&P 100*) for which sufficient data was available. First, we determined the mean-historical *VaR* Pareto-optimal frontier, in the *APF* framework, using the *NSGA-II* algorithm and compared it to the *LP* Pareto optimal frontier. The comparison showed that *NSGA-II* produced better mean-historical *VaR* trade-offs compared to the *LP* optimization approach, in both Low and High volatility samples. Second, we compared our mean-univariate *GARCH VaR* Pareto-optimal frontier with the frontiers obtained by the two benchmarks (mean-historical *VaR* and mean-multivariate *GARCH VaR*). In comparison with the two benchmarks, the proposed univariate *GARCH VaR* procedure provided actual portfolios with better mean-univariate *GARCH VaR* trade-offs, in both Low and High volatility samples.

Previous portfolio optimization studies typically neglect the differences between using portfolios with fixed weights and portfolios with fixed holdings of assets. We contribute to portfolio optimization literature by addressing recent real world regulatory changes which impose *VaR* based on actual portfolio holdings. The rare *MOEA* portfolio optimization studies that measured risk by using *VaR* tend to use historical rather than analytical *VaR* (see Anagnostopoulos and Mamanis [33], Hochreiter et al. [41], Soler et al. [42], Rankovic et al.

⁸ *NSGA-II* algorithm is also the most widely used *MOEA* in portfolio optimization literature. According to the number of studies reviewed in [30], *NSGA-II* was used in twice as many studies compared to second most popular method (*SPEA2*).

⁹ The number of the Pareto optimal solutions could also be reduced by combination of *NSGA-II* with Data Envelopment Analysis (see [40]).

[43]).¹⁰ We therefore also contribute to *MOEA* literature by examining the mean-*VaR* optimization problem with analytical univariate *GARCH VaR* instead of historical *VAR*.

The remainder of this paper is organized as follows. In Section 2 we introduce historical and *GARCH VaR* models. The optimization problem is introduced in Section 3. *MOAE* and implementation details are discussed in Section 4. Section 5 contains sample descriptive statistics. Results of mean-historical *VaR* optimization are presented in Section 6. Results of our univariate and multivariate mean-*GARCH VaR* optimization are presented and discussed in Section 7. We conclude in Section 8. All acronyms and notations are listed in Appendix.

2. *VaR* Models

2.1. Traditional (historical) *VaR*

For a given portfolio, significance level α and time horizon h , portfolio *VaR* is a loss that will be exceeded, on average, only $\alpha \times 100$ percent of the time. If expressed in terms of portfolio value, *VaR* is the α -quantile of profit and loss distribution (cash *VaR*), while if expressed in terms of portfolio return r , it is the α -quantile of the return distribution (relative *VaR* or, simply, *VaR*). We focused on the α -quantile of the return distribution. Consider time horizon of $h=1$ day and return r_α such that probability $p(r < r_\alpha) = \alpha$. In that case, 1-day *VaR* with significance level α is:

$$VaR^\alpha = -r_\alpha \quad (1)$$

The minus sign is needed since *VaR* is defined as a loss. Given the cumulative distribution function of returns $F(\alpha)$, the α -quantile is calculated as $r_\alpha = F^{-1}(\alpha)$. The main reason for the popularity of this method is ease of its implementation and the fact that it makes no assumptions about the parametric form of return distributions. On the other hand, the historical *VaR* often slowly reacts to abrupt changes in market conditions (see Alexander [11]).

¹⁰ Only 4.27% of *MOEA* studies use *VaR* as one of the objectives in the portfolio optimization problem (see [27]).

2.2. Analytical VaR

A popular alternative to historical simulation is provided by analytical (or parametric) *VaR* models. These models take a stand on the shape of the return distribution and capture the following stylized facts about asset returns (see Christoffersen [9]). First, asset returns are difficult to predict based on their past realizations. Second, volatility of daily returns dominates their mean. Third, unlike returns, the variance of daily returns tends to exhibit clustering over time. Namely, days with highly volatile returns are likely followed by days with highly volatile returns and vice versa. Fourth, after periods of high or low asset volatility, volatility tends to return toward a stable long run level. Fifth, even though asset returns are often modeled using a normal distribution, an unconditional return distribution usually has much heavier tails than predicted by a normal distribution.¹¹ Using the first two conditional moments of distribution, the return can be presented in the form:

$$r_t = \mu_t + \sigma_t z_t \quad (2)$$

where r_t is the return at time t , μ_t is the conditional mean, σ_t is conditional volatility of the return, and z_t is the residual (innovation term) of the process. It is assumed that z_t are independently and identically distributed and follow a known theoretical distribution D with zero mean and unit variance $D(0, 1)$.

To specify the analytical *VaR* model we specified the volatility updating model, as well as the shape of the conditional return distribution $D(0, 1)$. A commonly applied class of conditional volatility models that captures all of the above stylized facts of returns is the *GARCH* model.¹² In estimating *GARCH* parameters on daily data, we have taken into account that the conditional mean is dominated by the standard deviation of returns (see Christoffersen [9], Pritsker [15], Alexander [44]).¹³ This implies:

$$r_t \approx \sigma_t z_t \quad (3)$$

We took into account non-normality of standardized returns by assuming they follow a standardized Student's t distribution (with zero mean and unit variance) with d degrees of

¹¹ Although this is partially rectified when returns are standardized by time-varying volatility, some residual non-normality still remains.

¹² *GARCH* process was first applied to *VaR* modeling in [14].

¹³ Alternatively, one can apply the same model to mean-adjusted returns.

freedom t (d). The degree of freedom parameter is an additional parameter estimated jointly with the other model parameters using Maximum Likelihood Estimation (*MLE*) method. The 1-day ahead analytical *VaR* estimate with significance level α , calculated at time t , is obtained as follows:

$$VaR_{t+1}^{\alpha} = -\sigma_{t+1} t_{\alpha}^{-1}(d) \quad (4)$$

where σ_{t+1} is the 1-day ahead conditional volatility estimate at time t obtained by applying the proposed model, d are degrees of freedom of the estimated Student's t distribution of standardized portfolio returns, and $t_{\alpha}^{-1}(d)$ is α -quantile of the standardized Student's t distribution with d degrees of freedom.

Conditional portfolio volatility can be modeled directly using the time series of portfolio returns (we referred to this method as the Univariate *GARCH* approach). Alternatively, one can estimate conditional portfolio volatility using the conditional variance-covariance matrix estimated via the multivariate *GARCH* model. We referred to this method as the multivariate *GARCH* approach.

2.2.1. Univariate *GARCH VaR*

Consider the (univariate) time series of portfolio returns and denote by r_t the portfolio return at time t . The simplest and by far the most popular *GARCH* model of conditional volatility, commonly referred to as *GARCH* (1, 1), has the following form (see Bollerslev [8]):

$$\sigma_{t+1}^2 = \omega + \theta r_t^2 + \beta \sigma_t^2 \quad (5)$$

Here, $\theta, \beta > 0$ and $\theta + \beta < 1$.¹⁴ Thus, the univariate *GARCH VaR* is:

$$VaR_{t+1}^{\alpha} = -\left(\omega + \theta r_t^2 + \beta \sigma_t^2\right)^{1/2} t_{\alpha}^{-1}(d) \quad (6)$$

¹⁴ The second condition assures stationarity of the conditional volatility process.

2.2.2. Multivariate *GARCH* *VaR*

In a multivariate *GARCH* modeling framework, the 1-day ahead portfolio conditional volatility can be estimated as a function of the returns on individual assets (from the opportunity set) as follows:¹⁵

$$\sigma_{t+1}^2 = \mathbf{w}_t' \mathbf{H}_{t+1} \mathbf{w}_t \quad (7)$$

where $\mathbf{H}_{t+1}$ is $N \times N$ conditional variance-covariance matrix of returns of individual assets, and $\mathbf{w}_t$ is the vector of portfolio weights at time t .¹⁶ Importantly, the multivariate *GARCH* implicitly assumes a fixed portfolio weights approach (*FWA*).¹⁷ The portfolio weights are thus fixed at $w_{i,t} = w_{i,T}$ for all, $t \leq T$, where T denotes optimization date. Thus, the multivariate *GARCH* *VaR* is:

$$VaR_{t+1}^\alpha = -(\mathbf{w}' \mathbf{H}_{t+1} \mathbf{w})^{1/2} t_\alpha^{-1}(d) \quad (8)$$

3. Optimization problem

3.1. Mean-univariate *GARCH* *VaR* optimization (*APF* approach)

The problem we are trying to solve has the following general form:

$$\min_{\mathbf{w}} VaR_{T+1}^\alpha = -(\omega + \theta r_T^2 + \beta \sigma_T^2)^{1/2} t_\alpha^{-1}(d) \quad (9)$$

$$\max_{\mathbf{w}} E(r) \quad (10)$$

$$\text{subject to} \quad \sum_{i=1}^N w_i = 1 \quad (11)$$

$$0 \leq w_i \leq 1, \quad i = 1, \dots, N \quad (12)$$

Here, $\mathbf{w}$ denotes the vector of portfolio weights at optimization date $t=T$, its components are w_i , r is the return on the portfolio, VaR_{T+1}^α denotes the portfolio risk measure that we try to

¹⁵ See, for example, [26] and [45].

¹⁶ Boldface denotes matrices and vectors. Matrices are in upper case, whereas vectors are in lower case.

¹⁷ As discussed in [46], the fixed weights assumption further implies continuous portfolio rebalancing.

minimize, and $E(r)$ is the expected return on the portfolio. $t_\alpha^{-1}(d)$ here refers to a quantile of a univariate Student's t distribution with d degrees of freedom. Equation (11) describes the standard budget constraint which requires that weights sum up to 1. Equation (12) states that no short sales are allowed.¹⁸

To calculate actual portfolio VaR and the corresponding expected return, a time series of portfolio returns with fixed asset holdings is needed. We determined asset holdings at the optimization date by adopting an arbitrary dollar portfolio value (set, without loss of generality, to 1). Thus, we assumed that, at optimization date $t=T$, dollar portfolio value $V_{p,t=T}=1$. Portfolio holdings n_i at time T were determined based on weights at $t=T$:

$$n_i = \frac{w_i V_{p,t=T}}{P_{i,t=T}} = \frac{w_i}{P_{i,t=T}} \quad (13)$$

Here $P_{i,t}$ denotes the price of shares in company i at time t . In the actual portfolio approach, holdings are held fixed over time. The actual portfolio return at time $t \leq T$ is, therefore, determined as follows:

$$r_t = \frac{V_{p,t}}{V_{p,t-1}} - 1 = \frac{\sum_{i=1}^N n_i P_{i,t}}{\sum_{i=1}^N n_i P_{i,t-1}} - 1 \quad (14)$$

$V_{p,t}$ denotes the dollar portfolio value at time t . The actual portfolio mean- VaR optimization problem simply means that we used equation (14) when determining input returns for the above optimization problem.

In the actual portfolio approach, portfolio weights at time $t \leq T$, are given by the expression:

$$w_{i,t} = \frac{n_i P_{i,t}}{\sum_{i=1}^N n_i P_{i,t}} \quad (15)$$

Equations (14) and (15) imply that:

¹⁸ Short sale prohibition is a common constraint imposed on large institutional investors such as mutual or pension funds.

$$r_t = \sum_{i=1}^N w_{i,t-1} r_{i,t} \quad (16)$$

where $r_{i,t}$ is simple return on asset i at time t .

3.2. Mean - multivariate *GARCH* *VaR* optimization (*FWA* approach)

The portfolio optimization problem (Eq. (9) – (12)) can be specified as follows:

$$\min_{\mathbf{w}} VaR_{T+1}^\alpha = -(\mathbf{w}' \mathbf{H}_{T+1} \mathbf{w})^{1/2} t_\alpha^{-1}(d) \quad (17)$$

$$\text{subject to } \mathbf{w}' \boldsymbol{\mu} = \bar{r} \quad (18)$$

$$\text{subject to } \sum_{i=1}^N w_i = 1 \quad (19)$$

$$0 \leq w_i \leq 1, \quad i = 1, \dots, N \quad (20)$$

where $\mathbf{H}_{T+1}$ denotes the conditional variance-covariance matrix of individual returns estimated at the time of optimization $t=T$, $\boldsymbol{\mu}$ denotes vector of expected returns of individual assets, and $\bar{r}$ denotes the expected portfolio return as a weighted average of constituents' expected returns.

A portfolio return for an arbitrary, $t \leq T$, is given by the expression:

$$r_t = \sum_{i=1}^N w_{i,T} r_{i,t} \quad (21)$$

Quantile $t_\alpha^{-1}(d)$ refers to a quantile of a multivariate Student's t distribution with d degrees of freedom. Here, d does not depend on a portfolio composition. Namely, d and, thus, quantile $t_\alpha^{-1}(d)$ depend solely on the choice of constituent return series and not on the way in which they are combined into a particular portfolio. $\mathbf{H}_{T+1}$ and d are obtained using the Dynamic Conditional Correlation (*DCC*) model of Engle [47].¹⁹

DCC decomposes conditional variance-covariance matrix $\mathbf{H}_{t+1}$ into conditional standard deviations and correlations, that is:

¹⁹ We used 'rmgarch' package [48] within software R [49]. For fitting *GARCH* model we used 'dccfit' method. For the 1-day ahead estimation of conditional covariance matrix we used 'dccforecast' method.

$$\mathbf{H}_{t+1} = \mathbf{D}_{t+1} \mathbf{R}_{t+1} \mathbf{D}_{t+1} \quad (22)$$

where $\mathbf{D}_{t+1} = \text{diag}(\sigma_{1,t+1}, \dots, \sigma_{N,t+1})$. Here $\text{diag}(\ast)$ is the operator transforming a $N \times 1$ vector into a $N \times N$ diagonal matrix. We assume that conditional variances $\sigma_{i,t+1}^2$, $i = 1, \dots, N$, follow a standard univariate *GARCH* (1,1) model.

Matrix $\mathbf{R}_{t+1}$ is a symmetric positive definite conditional correlation matrix defined as:

$$\mathbf{R}_{t+1} = \text{diag}(\mathbf{Q}_{t+1})^{-1/2} \mathbf{Q}_{t+1} \text{diag}(\mathbf{Q}_{t+1})^{-1/2} \quad (23)$$

where $\mathbf{Q}_{t+1}$ is a proxy process which is assumed to follow *GARCH*-type dynamics:

$$\mathbf{Q}_{t+1} = (1 - \theta - \beta) \bar{\mathbf{Q}} + \alpha \mathbf{z}_t \mathbf{z}_t^T + \beta \mathbf{Q}_t \quad (24)$$

Vector $\mathbf{z}_t = (z_{1,t}, \dots, z_{N,t})^T$ has elements $z_{i,t} = r_{i,t} / \sigma_{i,t}$ (standardized unexpected returns or innovations), $\bar{\mathbf{Q}}$ is the $N \times N$ covariance matrix of $\mathbf{z}_t$ and θ and β are non-negative scalar parameters satisfying $\theta + \beta < 1$.

Since $t_\alpha^{-1}(d)$ is constant, the optimization problem (Eq. 17) is equivalent to a problem represented by quadratic programming formulation (referred to as *QP* model):²⁰

$$\min_{\mathbf{w}} \mathbf{w}' \mathbf{H}_{T+1} \mathbf{w} \quad (25)$$

It should be emphasized that quantile $t_\alpha^{-1}(d)$ in this approach is constant and therefore not portfolio specific. Thus, the multivariate *GARCH* approach is more restrictive compared to the univariate *GARCH* approach where the degrees of freedom of the estimated standardized Student's *t* distribution (and therefore quantile $t_\alpha^{-1}(d)$) were portfolio specific.

3.3. Summary of assumptions

3.3.1. General

- i) Short sales are not allowed (Eqs. (12) and (21));
- ii) The standard budget constraint which requires that weights must sum up to 1 (Eqs. (11) and (20)).

²⁰ This problem is of the standard Markowitz type (see [50]) with added short sales constraints.

3.3.2. Mean-univariate *GARCH* VaR optimization

- i) Portfolio returns, standardized by time varying volatility, follow a conditional standardized Student's t distribution (with zero mean and unit variance);
- ii) Conditional variance follows *GARCH* (1, 1) process, $\sigma_{t+1}^2 = \omega + \theta r_t^2 + \beta \sigma_t^2$, (Eq. 5);
- iii) In estimating *GARCH* parameters on daily data, the mean value of daily returns is dominated by the standard deviation of returns and that $r_t \approx \sigma_t z_t$, (Eq. 3);
- iv) At optimization date $t=T$, without loss of generality, dollar portfolio value $V_{p,t=T}=1$.

3.3.3. Mean-multivariate *GARCH* VaR optimization

- i) Portfolio weights are fixed over the observed period;
- ii) Returns of assets from the opportunity set jointly follow a multivariate Student's t distribution;
- iii) $\mathbf{Q}_{t+1}$ (Eq. 23) follows *GARCH*-type dynamics: $\mathbf{Q}_{t+1} = (1 - \theta - \beta) \bar{\mathbf{Q}} + \alpha \mathbf{z}_t \mathbf{z}_t^T + \beta \mathbf{Q}_t$
- iv) In matrix $\mathbf{D}_{t+1}$ conditional variance $\sigma_{i,t+1}^2$ follows a univariate *GARCH* (1, 1) process

4. Portfolio optimization methodology

4.1. Evolutionary algorithms (EA)

EA are efficient stochastic search techniques for solving complex optimization and search problems (e.g., optimization problems with non-differentiable objective functions, large and non-convex solution spaces, complex constraints etc.) through an emulation of natural selection i.e. survival of the fittest (see Sastry et al. [51]). *EA* begin with a set of randomly generated candidate solutions, referred to as a population. In each of the iterations (i.e. generations) the following processes are performed: i) Good solutions from the current population are selected and transferred into a set of potential parents (mating pool); ii) Randomly selected parent solutions are combined (crossover) producing new solutions (offspring solutions); iii) Some randomly selected offspring solutions are slightly modified (mutation); iv) Generated offspring solutions constitute the population of the next generation. By performing these processes in each generation (until the termination condition is satisfied) the solutions "evolve" and become even better in fulfilling the stated objectives.

Over the past two decades, significant advances have been made on *EA* methods to solve multi-objective optimization problems.²¹ This has led to the creation of multi-objective evolutionary algorithms (*MOEA*). In general, *MOEA* address two important issues: i) How to generate a Pareto-optimal frontier; and ii) How to provide diversity in alternative solutions (i.e. how to avoid convergence to a single point on the Pareto-optimal frontier). To achieve the first goal, most *MOEA* implementations use ranking based on the concept of dominance, while different diversity-preserving techniques are employed to achieve the second goal.

4.2. *MOEA* implementation

We directly borrowed the *NSGA-II* algorithm from Deb et al. [37].²² Implementation of *NSGA-II* involves adopting settings for the solution representation, the population size, the crossover and mutation probabilities and the termination condition. Solution representation depends upon the specific optimization problem. For the model at hand, the solution was defined as a non-negative real-valued vector of portfolio weights at time $t=T$, that is, the vector of fractions of the total budget invested in individual securities (see Section 4). Population size (the number of candidate portfolios in each generation), was set to 100. The Pareto-optimal frontier was, therefore, approximated with 100 points. The step by step flow chart which presents a schematic view of the proposed portfolio optimization method is shown in Figure 1.

In the preparatory phase our software generated and printed a data file (data.csv) with a time series of occurrences of assets under consideration. Next, it generated and printed an R script file (Script.R) with commands for estimation of the univariate *GARCH VaR*. It then executed the *NSGA-II* algorithm in the execution phase.²³

²¹ For a detailed introduction to *MOEA*, see [52].

²² The algorithm was coded in C# and run on a personal computer with Intel i5 processor and 4GB of RAM.

²³ For more details on the algorithm steps, see [37].

Figure 1. Flow chart of the optimization algorithm

For the breeding of an offspring population we used the uniform crossover operator. Two solutions (portfolios) from the current population were randomly selected and were recombined with a predefined crossover probability. If the two solutions undergo recombination, every allele(individual asset weight) is exchanged between the pair of randomly selected solutions with a certain probability, known as the swapping probability, otherwise, the two offspring are simply copies of their parents (see Sastry et al. [51]).²⁴ In accordance with previous literature, we set the swapping probability to be 0.5.²⁵

We applied a uniform mutation operator (uniform replacement). When applying uniform mutation, each allele is selected with a predefined mutation probability and replaced with a realization of a random variable, uniformly distributed in the range defined by the lower and upper domain bounds. The selected crossover and mutation operator ensured that the constraint defined by Eq. (12) was satisfied for each offspring. However, these operators did not ensure satisfaction of the budget constraint (Eq.(11)). Hence, we had to normalize each of the offspring solutions.²⁶

In order to find the appropriate parameter values for the crossover and mutation probabilities, we performed a series of experiments. The performance was assessed using the ϵ –indicator,

²⁴ Candidate solutions are also referred to as chromosomes, decision variables are referred to as genes and the values of decision variables are called alleles.

²⁵ See [31], [32], [33], [51].

²⁶ We have done this by dividing each weight by sum of all weights.

and the hypervolume metric (see Zitzler et al. [53]). The ε -indicator is a binary performance metric which is used to measure how close the approximation set is to the reference set. As a reference set, the true or the best known Pareto-optimal frontier was used. The ε -indicator determines a minimum value the reference set must be multiplied by in order for every solution in the reference set to become weakly dominated by at least one solution in the approximation set. If the approximation set matches the reference set exactly then the ε -indicator takes the value of one. For this metric, values close to one indicate that the approximation set very closely matches the reference set. The hypervolume indicator was used to measure the diversity of the approximation set. The hypervolume measure quantifies the volume of the objective space dominated by an approximation set. For optimization problems with two objectives, it quantifies the area of the objective space dominated by the approximation set, bounded by a predefined reference point. Thus, for this metric higher values are preferable.

To determine the best performing mutation and crossover probability sets, we tested four crossover probabilities (0.7, 0.8, 0.9, 1.0) and five uniform mutation probabilities (0.001, 0.005, 0.01, 0.05, 0.1). The above tests resulted in 20 different configurations for each set of objectives (mean-historical *VaR*, mean-*GARCH VaR*) and for each data sample that we used. For each configuration, the algorithm was left to run until 100,000 solutions were generated.

The interactions between the *NSGA-II* algorithm and our statistical analysis were highlighted in Figure 2. For the evaluation of each individual solution portfolio, our software performed the following steps: i) It generated and printed a file with portfolio weights (weights.csv); ii) It called software R and executed the R script file (Script.R) created in the preparatory phase. During the execution of the script file, software R used the data file (data.csv) and the solution portfolio weights file (weights.csv) and generated a time series of actual portfolio returns (applying Eq. (13) –(14)). Then *VaR* was estimated using the univariate *GARCH* model (Eq. (4) and (5)).²⁷

²⁷ We used ‘rugarch’ package [54] within software R [49]. For fitting *GARCH* model we used ‘ugarchfit’ method; For the 1-day ahead estimation of conditional volatility we used ‘ugarchforecast’ method.

Figure 2. Interactions between the *NSGA-II* algorithm and statistical computing

In the case of the mean-historical *VaR* approach, algorithm execution requires calculating the α quantile of the empirical distribution of 100,000 candidate portfolios. However, in the case of the mean-univariate *GARCH VaR* approach, the optimization procedure implies calculating the α quantile of the parametric distribution of candidate portfolios. Additional complexity of the univariate *GARCH VaR* approach stems from the fact that the *GARCH* parameters for each of 100,000 portfolios were obtained by the *MLE* method. We ensure that all Pareto-optimal frontiers, obtained using different configurations, include solutions which provide the minimum risk and the maximum return.²⁸

5. Sample descriptive statistics

For the purposes of this study, we selected 40 constituents of the *S&P 100* with the highest market capitalization as of Sept 6th 2013.²⁹ We observed the constituents' time series of 1,421 returns in the period Jan 15th 2008-Sept 6th 2013. We used a rolling window of 1,000 returns for *VaR* estimation. For each day within the rolling estimation period of 421 days (Jan 4th 2012-Sept 6th 2013) we estimated 1-day ahead daily volatility of the *S&P 100* time series. The volatilities were estimated using the univariate *GARCH (1, 1)* model. Standardized returns were assumed to have a standardized *t*-distribution. *GARCH* volatility estimations of the *S&P 100* index were based on a rolling window of 1,000 daily returns. Maximum volatility of

²⁸ We obtained these solutions and included them into the initial population of each *MOEA* execution. Maximum return solution was obtained analytically, while minimum risk solution was obtained using single-objective genetic algorithm.

²⁹ Initially, 43 stocks were considered of which 3 were discarded (tickers: PM, V, KFT) due to incomplete data during the sample period.

0.01294 (20.5% in annual terms) was determined on June 29th 2012. Minimum volatility of 0.00583 (9.3% in annual terms) was determined on July 31st 2013.³⁰ From the chosen sample of 40 stocks we created two samples with 1,001 daily occurrences: i) First, ending on the maximum volatility date (referred to as the High volatility sample); and ii) Second, ending on the minimum volatility date (referred to as the Low volatility sample). All *VaR* values in this paper correspond to a significance level of 0.01, with a time horizon of one day. Table 1 presents a summary statistics of the time series comprising the Low and High volatility samples.

Table 1. Descriptive statistics for the Low and High volatility samples

Sample	Average expected return (annualized)	Average volatility (annualized)	Average historical <i>VaR</i>	Average <i>GARCH VaR</i>
Low volatility	16.61%	23.47%	3.96%	3.12%
High volatility	13.47%	37.94%	6.95%	4.12%

6. Mean-historical *VaR* optimization

Rockafellar and Uryasev [20, 21] and Krokmal et al. [22] argue that the mean-*CVaR* efficient frontier generated by the *LP* model also provides near-optimal solutions in the context of mean-historical *VaR* optimization. We tested this assertion in an actual portfolio framework. Namely, for each portfolio belonging to the mean-*CVaR* Pareto-optimal frontier generated by a *LP* (*FWA* approach), we created a time series of actual portfolio returns (applying Eq. (13)–(14)), calculated the corresponding 1% historical *VaR* and expected return, and represented the portfolios in the actual portfolio mean-*VaR* plane. In this way we generated an approximate Pareto-optimal frontier corresponding to the *LP* solution. Next, we determined the mean-historical *VaR* Pareto-optimal frontier (in the actual portfolio framework directly) using the *NSGA-II* optimization approach (see Figure 3). Figure 3 shows that the *NSGA-II* frontiers dominate the *LP* frontiers, especially in the High volatility sample.

³⁰ We annualize volatility using square root of time assuming 252 trading days per annum.

Figure 3.a. *NSGA-II* and *LP* approximate mean-historical *VaR* Pareto-optimal frontier in the Low volatility sample

Figure 3.b. *NSGA-II* and *LP* approximate mean-historical *VaR* Pareto-optimal frontier in the High volatility sample

Best performing *NSGA-II* mean-historical *VaR* approximate of the Pareto-optimal set was selected based on the hypervolume metric. To compare the *NSGA-II* mean-historical *VaR* Pareto-optimal frontier to the *LP* Pareto-optimal set, we used the ε -indicator. As a reference set, we adopted the *NSGA-II* mean-historical *VaR* Pareto-optimal frontier. The comparison is presented in Table 2. The values of the hypervolume parameter and reference points, together with the ε -indicator, were for the best performing *NSGA-II* approximates of the mean-*VaR* Pareto-optimal frontier.

Table 2. Comparison of *NSGA-II* mean-historical *VaR* and *LP* models

Sample	Mutation probability	Crossover rate	Hypervolume	Reference point	ε -indicator
Low volatility	0.005	0.8	4.2366E-05	(0.0521,0)	1.1300
High volatility	0.05	1	6.9160E-05	(0.0823,0)	1.1526

The values for the ε -indicator confirm that the *NSGA-II* optimization approach provides better mean-historical *VaR* trade-offs of actual portfolios compared to the *LP* solutions in both the High and Low volatility samples. The relatively higher value of the ε -indicator (1.1526) suggests that *NSGA-II* performed particularly well during the High volatility period.

7. Results for mean-GARCH VaR optimization

7.1. Pareto-optimal frontiers in the Low and High volatility samples

In this section we compared the Pareto-optimal frontiers for: i) The mean-univariate *GARCH VaR* (or univariate *GARCH* for short); ii) The mean-multivariate *GARCH VaR* (benchmark 1); and iii) The mean-historical *VaR* (benchmark 2).³¹ In order to obtain solutions in the actual portfolio framework for both benchmark portfolio solutions (represented by the corresponding vector of weights $\mathbf{w}$), we generated a times series of portfolio returns. The time series of portfolio returns was generated by employing Equations (13) and (14) within the *APF* approach. We then calculated *VaR* using the univariate *GARCH* model (Eq. (4) and (5)).³²

The results are shown in Figure 4. The Pareto-optimal frontiers, for actual portfolios, are for the two volatility regimes when 1%*VaR* was estimated using the univariate *GARCH* model. Triangle markers represent the univariate *GARCH* frontier (obtained via *NSGA-II*), while filled dots represent benchmark 1 solutions and empty dots represent benchmark 2 solutions.

³¹ As presented in Figure 3, the *NSGA-II* method provided superior actual portfolio mean-historical *VaR* trade-offs compared to the *LP* model, in both samples. Consequently, we adopted the *NSGA-II* mean-historical *VaR* portfolios as benchmark 2.

³² During this process some of benchmark 1 and benchmark 2 portfolio solutions became dominated and thus have been discarded from the approximation set.

Figure 4a. Pareto-optimal frontiers in the Low volatility sample

Figure 4b. Pareto-optimal frontiers in the High volatility sample

The differences are particularly prominent in the area of low returns (and low risk). In this segment of the Pareto-optimal set, thus, the opportunities for improving the mean-*VaR* trade-off through optimization are the greatest. It is worth noting that this segment of the Pareto-optimal set is associated with most diversified portfolios. In contrast, with higher expected return values, the cardinality of efficient portfolios is reduced (the highest return portfolio, by construction, corresponds to a single asset).

To further compare the univariate *GARCH* Pareto-optimal frontier to the respective benchmarks, we used the ϵ -indicator. As a reference set, we adopted the univariate *GARCH*

frontier. Table 3 presents the values of the hypervolume parameter and reference points for the best performing *NSGA-II* mean-univariate *GARCH VaR* approximate Pareto-optimal sets. The table also shows ε -indicators corresponding to the respective benchmarks in the Low and High volatility samples.

Table 3. Parameters of the best performing mean-univariate *GARCH VaR*

	Mutation probability	Crossover rate	Hypervolume	Reference Point	ε -indicator	
					Benchmark 1	Benchmark 2
Low volatility	0.05	0.7	5.0973E-05	(0.0504, 0)	1.2608	1.4618
High volatility	0.01	0.9	5.3451E-05	(0.0581, 0)	1.0866	1.3243

The above presented ε -indicators confirm superiority of the univariate *GARCH VaR* optimization approach. The advantages of the univariate *GARCH VaR* optimization approach are particularly pronounced in the Low volatility sample. Measured by the ε -indicators, the mean-historical *VaR* approach provides the worst approximations out of the three approaches. The above result is particularly important given that approximately 75% of banks tend to use the historical *VaR* models for portfolio optimization (see Pérignon and Smith [55]).

7.2. Process time (CPU time)

As expected, our *CPU* time is longer compared to the *CPU* time of the traditional methods (e.g. *QP*). For example, the execution of one generation of *NSGA-II*, when solving the mean-univariate *GARCH VaR* problem, lasted 41 seconds. Hence, to execute 1,000 generations of *NSGA-II* we needed 41,000 seconds.³³ The total *CPU* time for mean-univariate *GARCH VaR* optimization with the *QP* solver was 424 seconds.³⁴ Our additional analysis, however, revealed that the *NSGA-II* *CPU* time could be shorter since the execution of 1,000 generations was not always necessary. For example, for the mean-historical *VaR* optimization, 99% of final hypervolume was achieved after 50 generations (12 seconds) in the Low volatility sample, and after 52 generations (12.5 seconds) in the High volatility sample.³⁵ The same

³³ In comparison, it took only 1 second for the *LP* model to generate mean-historical *VaR* Pareto-optimal frontier consisting of 100 solutions.

³⁴ The mean-univariate *GARCH VaR* optimization (using multivariate *GARCH VaR* approach) consisted of three steps: i) Determination of conditional variance-covariance matrix of individual returns (H_{T+1}); ii) Execution of *QP* solver; and iii) Estimation of univariate *GARCH VaR* for each solution obtained by using *QP* solver. To determine conditional variance covariance matrix 382 seconds was needed. The execution of *QP* solver for Pareto optimal front of 100 solutions lasted 1 second. Finally the estimation of univariate *GARCH VaR*, for 100 solutions obtained by using *QP* solver, lasted 41 second.

³⁵ We define final hypervolume as a hypervolume of Pareto-optimal set generated after 1,000 generations.

analysis applied on the mean-univariate *GARCH VaR* optimization revealed that 99% of final hypervolume was achieved after 40 generations (1,640 seconds) in the Low volatility sample. The corresponding values in the High volatility sample were 20 generations and 820 seconds. Given its longer *CPU* time, the use of *NSGA-II* could be justified by its greater flexibility and ability to deal with more complex real-life portfolio optimization problems compared to the *QP* method (see Deb et al. [35]).

7.3. Out-of-sample estimates

Bank managers and regulators are interested in the out-of-sample performance of different optimization models. Thus, we compared out-of-sample performance of the different optimization models used in our study. The optimization dates for our High and Low volatility samples were dates with the highest (29th June 2012) and lowest (31st July 2013) volatilities. Based on the estimated portfolios on the optimization dates, we calculated respective mean and *VaR* values for dates which fall exactly 1 month later (30th July 2012 and 3rd September 2013 respectively).

Figure 5a. Out-of-sample test in the High volatility sample

Figure 5b. Out-of-sample test in the Low volatility sample

The results of the out-of-sample analysis were consistent with the results reported in Figure 4, thus confirming the superiority of estimates based on the mean-univariate *GARCH* model in both volatility samples. As expected, the differences between the results for the mean-univariate and mean-multivariate *GARCH* models are less pronounced in the Low volatility sample.

8. Conclusions

According to Basel regulation, the riskiness of a portfolio and the implied capital charge are calculated using *VaR* estimations of an actual portfolio of a financial institution (i.e. the portfolio that corresponds to current portfolio holdings). This paper proposes a novel approach to mean-*VaR* portfolio optimization within the actual portfolio framework when *VaR* is estimated by the analytical univariate *GARCH VaR* model with the assumption of a conditional *t* distribution of standardized portfolio returns. Due to the complexity of the proposed optimization problem, we applied metaheuristics. Specifically, we developed software which combined a *NSGA-II* multi-objective evolutionary algorithm with software for statistical computing R. In the empirical section, we examined the opportunity set consisting of 40 large US stocks belonging to the *S&P 100* index in two volatility regimes. We found that the *NSGA-II* method resulted in better actual portfolio mean-historical *VaR* trade-offs in both Low and High volatility regimes.

Next, we compared the mean-univariate *GARCH VaR* Pareto-optimal frontier to the mean-multivariate *GARCH VaR* and to the mean-historical *VaR* Pareto-optimal frontiers. In comparison to the two benchmarks, the proposed univariate *GARCH VaR* procedure, again, provided actual portfolios with a superior mean-univariate *VaR* frontier for both volatility samples. The results suggest that the multivariate *GARCH* modeling framework lacks flexibility to conform to the actual portfolio framework which is inherent in regulation. At the same time, the mean-historical *VaR* frontier provided the worst mean-univariate *GARCH VaR* trade-offs. Overall, our results bear two important implications for financial institutions and their regulators. First, the results highlighted differences between the actual portfolio approach and the approach based on the fixed weights. Second, the results show the importance of carefully selecting amongst different *VaR* methodologies used in portfolio optimization.

Recently, some multiple criteria decision making (*MCDM*) tools were combined with *NSGA-III* to either rank or reduce the Pareto optimal frontier (see Tavana et al. [56]). Further enhancement of our optimization approach with some of the *MCDM* reduction techniques could be an interesting area for future research.

Acknowledgements: We would like to thank two anonymous referees for their helpful suggestions. Research presented in this paper was supported by Serbian Ministry of Education, Science and Technology, Grants OH 179005 and III-44010.

References

- [1] Artzner P, Delbaen F, Eber JM, Heath D. Coherent measures of risk. *Mathematical Finance* 1999; 9(3): 203–228.
- [2] Szego G. Measures of risk. *Journal of Banking & Finance* 2002; 26: 1253–1272.
- [3] Basel Committee on Banking Supervision. International convergence of capital measurement and capital standards. Bank for International Settlements (BIS); 1988.
- [4] Basel Committee on Banking Supervision. Amendment to the capital accord to incorporate market risks. Bank for International Settlements (BIS); 1996.
- [5] Basel Committee on Banking Supervision. Revisions to the Basel II market risk framework. Bank for International Settlements (BIS); 2009.
- [6] Hendricks D, Hirtle B. Bank capital requirements for market risk: the internal models approach. *Economic Policy Review* 1997; 3: 1–12.
- [7] Engle R. Autoregressive conditional heteroscedasticity with estimates of the variance of UK inflation. *Econometrica* 1982; 50(4): 987–1007.
- [8] Bollerslev T. Generalized autoregressive conditional heteroscedasticity. *Journal of Econometrics* 1986; 31: 307–327.
- [9] Christoffersen PF. *Elements of Financial Risk Management*. 2nd ed. San Diego CA: Academic Press; 2012.
- [10] Huisman R, Koedijk K, Pownall R. VaR-x: Fat tails in financial risk management. *Journal of Risk* 1998; 1: 47–61.
- [11] Alexander C. *Value-at-Risk Models*. 1th ed. England: John Wiley & Sons; 2008b.
- [12] Alexander C, Sheedy E. Developing a stress testing framework based on market risk models. *Journal of Banking & Finance* 2008; 32: 2220–2236.
- [13] Berkowitz J, O'Brien J. How accurate are value-at-risk models at commercial banks? *Journal of Finance* 2002; 57: 1093–1111.
- [14] Hull J, White A. Incorporating volatility updating into the historical simulation method for value at risk. *Journal of Risk* 1998; 1: 5–19.
- [15] Pritsker M. The hidden dangers of historical simulation, *Journal of Banking & Finance* 2006; 30: 561–582.
- [16] Gaivoronski AA, Pflug G. Value-at-risk in portfolio optimization: properties and computational approach. *Journal of Risk* 2005; 7(2): 1–31.

- [17] Goh JW, Guan LK, Sim, M, Zhang W. Portfolio value-at-risk optimization for asymmetrically distributed asset returns. *European Journal of Operational Research* 2012; 221: 397–406.
- [18] Krejić N, Kumaresan M, Rožnjik A. VaR optimal portfolio with transaction costs. *Applied Mathematics and Computation* 2011; 218: 4626-4637.
- [19] Natarajan K, Pachamanova D, Sim M. Incorporating asymmetric distributional information in robust value-at-risk optimization. *Management Science* 2008; 54(3): 573-585.
- [20] Rockafellar RT, Uryasev S. Optimization of conditional value-at-risk. *Journal of Risk* 2000; 2(3): 21–41.
- [21] Rockafellar RT, Uryasev S. Conditional value-at-risk for general loss distributions. *Journal of Banking & Finance* 2002; 26: 1443–1471.
- [22] Krokmal P, Palmquist J, Uryasev S. Portfolio optimization with conditional value-at-risk objective and constraints. *Journal of Risk* 2002; 4(2): 11-27.
- [23] Iscoe I, Kreinin A, Mausser H, Romanko O. Portfolio credit-risk optimization, *Journal of Banking & Finance* 2012; 36: 1604–1615.
- [24] Pfaff B. *Financial risk modelling and portfolio optimization with R*. UK: John Wiley & Sons; 2013.
- [25] Würtz D, Chalabi Y, Chen W, Ellis A. *Portfolio optimization with R/Rmetrics*. Rmetrics; 2009.
- [26] Santos A, Francisco J, Nogales ER, Dick VD. Optimal portfolios with minimum capital requirements. *Journal of Banking & Finance* 2012; 36: 1928–1942.
- [27] Metaxiotis K, Liagkouras K. Multi-objective evolutionary algorithms for portfolio management: a comprehensive literature review. *Journal of Expert Systems with Applications: An International Review* 2012; 39 (14): 11685-11698.
- [28] Schlottmann F, Seese D. Financial applications of multi-objective evolutionary algorithms: Recent development and future research directions. In: Coello-Coello C, Lamont G. editors. *Applications of Multi-objective Evolutionary Algorithms*. World Scientific; 2004, p. 627-652.
- [29] Tapia MGC, Coello CAC. Applications of multi-objective evolutionary algorithms in economics and finance. In: *IEEE Congress on Evolutionary Computation*; 2007, p. 532–539.
- [30] Ponsich A, L'opez J A, Coello CAC. A Survey on multiobjective evolutionary algorithms for the solution of the portfolio optimization problem and other finance

- and economics applications, *IEEE Transactions on Evolutionary Computation* 2013; 17 (3): 321-344.
- [31] Anagnostopoulos KP, Mamanis G. The mean-variance cardinality portfolio optimization problem: An experimental evaluation of five multi-objective evolutionary algorithms, *Expert Systems with Applications* 2011; 38: 14208–14217.
 - [32] Anagnostopoulos KP, Mamanis G. Multiobjective evolutionary algorithms for complex portfolio optimization problems, *Computational Management Science* 2011; 8: 259–279.
 - [33] Anagnostopoulos KP, Mamanis G. A portfolio optimization model with three objectives and discrete variables. *Computers & Operations Research* 2010; 37: 1285–1297.
 - [34] Deb K, Agrawal S, Pratap A, Meyarivan TA. A fast elitist non-dominated sorting genetic algorithm for multi-objective optimization: NSGA-II. *Lecture Notes in Computer Science* 2000; 1917: 849-858.
 - [35] Deb K, Steuer RE, Tewari R, Tewari R. Bi-objective portfolio optimization using a customized hybrid NSGA-II procedure. In: Takahashi RHC, Deb K, Wanner EF, Greco S. editors. *EMO 2011. Lecture Notes in Computer Science* 6576. Berlin: Springer-Verlag; 2011, p. 358–373.
 - [36] Branke J, Scheckenbach B, Stein M, Deb K, Schmeck H. Portfolio optimization with an envelope-based multi-objective evolutionary algorithm. *European Journal of Operational Research* 2009; 199: 684–693.
 - [37] Deb K, Pratap A, Agarwal S, Meyarivan, TA. Fast and elitist multi-objective genetic algorithm: NSGA-II. *IEEE Transactions on Evolutionary Computation* 2002; 6(2):182–197.
 - [38] Deb K, Jain H. An evolutionary many-objective optimization algorithm using reference-point-based nondominated sorting approach, part 1: solving problems with box constraints. *IEEE Transactions on Evolutionary Computation* 2014; 18(4): 577-601.
 - [39] Jain H, Deb K. An evolutionary many-objective optimization algorithm using reference-point based non-dominated sorting approach, part 2: handling constraints and extending to an adaptive approach. *IEEE Transactions on Evolutionary Computation* 2014; 18(4): 602-622.

- [40] Mobin M, Li Z, Khoraskani M. Multi-objective X-bar control chart design by integrating NSGA-II and data envelopment analysis. In: Proceedings of the Industrial and Systems Engineering Research Conference (ISERC), TN, USA; 2015.
- [41] Hochreiter R. An evolutionary computation approach to scenario risk-return portfolio optimization for general risk measures. Department of Statistics and Decision Support Systems-University of Vienna, 2007.
- [42] Soler JS, Cid EA, Blanco MO. Mean-VaR portfolio selection under real constraints. *Comput. Econ* 2011; 37:113-131.
- [43] Ranković V, Drenovak M, Stojanović B, Kalinić Z, Arsovski Z. The mean-value-at-risk static portfolio optimization using genetic algorithm. *Computer Science and Information Systems* 2014; 11 (1): 89–109.
- [44] Alexander C. *Practical Financial Econometrics*. England: John Wiley & Sons; 2008.
- [45] Engle RF, Kroner KF. Multivariate simultaneous generalized ARCH. *Econometric Theory* 1995; 11:122-150.
- [46] Alexander C. *Quantitative methods in finance*. England: John Wiley & Sons; 2008.
- [47] Engle R. Dynamic conditional correlation: a simple class of multivariate generalized autoregressive conditional heteroskedasticity models. *Journal of Business & Economic Statistics* 2002; 20: 339–350.
- [48] Ghalanos A. rmgarch: Multivariate GARCH models. R package version 1.2-8; 2014.
- [49] R Core Team. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing 2014. Vienna: URL <http://www.R-project.org/>.
- [50] Markowitz HM. Portfolio selection. *Journal of Finance* 1952; 7: 77–91.
- [51] Sastry K, Goldberg D, Kendall G. Genetic algorithms. In: Edmund KB, Graham K. editors. *Search methodologies: Introductory tutorials in optimization and decision support techniques*. New York: Springer Science+Business Media; 2005, p. 97-127.
- [52] Coello CAC, Lamont GB, Van Veldhuizen DA. *Evolutionary algorithms for solving multi-objective problems*. 2nd ed. Berlin: Springer; 2007.

- [53] Zitzler E, Thiele L, Laumanns M, Fonseca CM, Grunert da Fonseca V. Performance assessment of multi-objective optimizers: an analysis and review. *IEEE Transactions on Evolutionary Computation* 2003; 7: 117–32.
- [54] Ghalanos A. rugarch: Univariate GARCH models. R package version 1.3-3; 2014.
- [55] Pérignon C, Smith D. The Level and quality of value-at-risk disclosure by commercial banks, Working paper - Simon Fraser University, 2007.
- [56] Tavana, M., Li, Z., Mobin, M., Komaki, M., Teymourian, E. Multi-objective control chart design optimization using NSGA-III and MOPSO enhanced with DEA and TOPSIS. *Expert Systems with Applications* 2016; 50; 17-39.

Accepted manuscript

Appendix

Table A1: Acronyms

<i>APF</i>	Actual Portfolio Framework
<i>CPU Time</i>	Process Time
<i>CR</i>	Capital Requirements
<i>CVaR</i>	Conditional Value-at-Risk
<i>DCC</i>	Dynamic Conditional Correlation Model
<i>EA</i>	Evolutionary Algorithm
<i>ES</i>	Expected Shortfall
<i>FWA</i>	Fixed Weights Approach
<i>GARCH</i>	Generalized AutoRegressive Conditional Heteroscedasticity
<i>LP</i>	Linear Programming
<i>MCDM</i>	Multiple Criteria Decision Making
<i>MLE</i>	Maximum Likelihood Estimation
<i>MOEA</i>	Multi-objective Evolutionary Algorithm
<i>NSGA-II (NSGA-III)</i>	Nondominated Sorting Genetic Algorithm
<i>PESA</i>	Pareto Envelope-based Selection Algorithm
<i>PAES</i>	Pareto Archived Evolutionary Strategy
<i>QP</i>	Quadratic Programming
<i>S&P 100</i>	Standard and Poor's 100 stock market index
<i>SPEA</i>	Strength Pareto Evolutionary Algorithm
<i>VaR</i>	Value-at-Risk

Table A2: Notations

r_t	Portfolio return
μ_t	Conditional expected return
σ_t^2	Conditional variance
z_t	Residual (innovation) term
p	Portfolio
ρ	Probability
$V_{p,t}$	Dollar portfolio value
σ_{t+1}	1-day ahead conditional volatility
$F(\alpha)$	Cumulative distribution function of returns
α	Significance level
h	Time horizon
d	Degrees of freedom
$\mathbf{H}_{t+1}$	1-day ahead conditional variance-covariance matrix of returns
$\mathbf{w}_t$	Vector of portfolio weights
VaR_{t+1}^α	<i>GARCH</i> <i>VaR</i> of a portfolio
$E(r)$	Portfolio expected return
$P_{i,t}$	Company i share price
n_i	Portfolio holdings