

## The dual standard model and the 750 GeV events at the LHC

Article (Accepted Version)

Calmet, Xavier (2016) The dual standard model and the 750 GeV events at the LHC. Europhysics Letters (EPL), 115 (2). p. 21001. ISSN 0295-5075

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/62231/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

### **Copyright and reuse:**

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

# The Dual Standard Model and the 750 GeV Events at the LHC

Xavier Calmet<sup>1</sup>

*Physics and Astronomy, University of Sussex, Falmer, Brighton, BN1 9QH, UK*

## Abstract

The aim of this short paper is to discuss the recently observed excess at 750 GeV by both CMS and ATLAS in the light of the dual standard model. Within this framework it is natural to introduce neutral spin 0 and/or spin 2  $SU(2)$  glue mesons which could easily account for this observation if it is confirmed. The model predicts that these glue mesons would be part of  $SU(2)$  triplets and that there must thus be charged counterparts of these glue mesons carrying a QED charge of  $\pm 1$  with a spin 0 and/or 2 as well.

---

<sup>1</sup>x.calmet@sussex.ac.uk

The aim of this short paper is to discuss the recently observed excess at 750 GeV by both CMS [1] and ATLAS [2] in the light of the dual standard model [3,4]. This excess can be interpreted as a spin 0 or spin 2 resonance of mass 750 GeV decaying into two photons. After briefly reviewing the dual standard model, we show that if the recent excess observed at CERN is a spin 0 or spin 2 resonance described within this framework, then this resonance must be the first sign of a new series of particles with similar masses. Indeed the spin 0 or 2 particle would be part of a  $SU(2)$  triplet. Besides the neutral spin 0 or 2 particles, one expects two states with the same spin and QED charges  $\pm 1$ . As we shall see, up to QED effects, these charged particles would have masses similar to that of their neutral counterpart.

The standard model provides a very successful description of the electroweak and strong interactions. It is based on the concept of local  $SU(3)_c \times SU(2)_L \times U(1)_Y$  gauge invariance where  $SU(3)_c$  is the gauge group that describes the strong interaction and  $SU(2)_L$  and  $U(1)_Y$  describe the weak and electromagnetic interactions. Besides the local gauge symmetries, there is another important but approximative global symmetry  $SU(2)$  due to the structure of the Higgs potential of the standard model. This symmetry would be exact in the limit where the gauge coupling of the  $U(1)_Y$  goes to zero. This custodial symmetry is particularly important when studying the standard model in its dual picture as we shall explain shortly. Indeed, it is well known that the standard model of particle physics can be defined in terms of gauge invariant fields. This formulation, while equivalent to the original presentation due to Glashow, Weinberg and Salam, is useful as it makes clear that particles are never isolated but always surrounded by virtual particles. Indeed a physical electron is never isolated from the photon and virtual pairs of particles constantly pop out of the vacuum. We can define fields which are gauge invariant under the local  $SU(2)_L$  gauge symmetry. These “physical” fields are given by [3–7]

$$\begin{aligned}
\Phi &= \Omega^\dagger \phi, \\
\Psi_L^a &= \Omega^\dagger \psi_L^a \\
W_\mu^i &= \frac{i}{2g} \text{Tr} \left[ \Omega^\dagger \overleftrightarrow{D}_\mu \Omega \tau^i \right] \\
(\mathcal{D}_\mu)_{SU(2)_L} &= \Omega^\dagger (D_\mu)_{SU(2)_L} \Omega = \partial_\mu - igW_\mu \\
\mathcal{F}_{\mu\nu} &= \frac{i}{g} [(\mathcal{D}_\mu)_{SU(2)_L}, (\mathcal{D}_\nu)_{SU(2)_L}],
\end{aligned} \tag{1}$$

with  $\phi^\dagger \overleftrightarrow{D}_\mu \phi = \phi^\dagger D_\mu \phi - (D_\mu \phi)^\dagger \phi$  and where  $\Omega$  is given by

$$\Omega = \frac{1}{\sqrt{\phi^\dagger \phi}} \begin{pmatrix} \phi_2^* & \phi_1 \\ -\phi_1^* & \phi_2 \end{pmatrix}. \tag{2}$$

Here  $\Phi$  corresponds to the Higgs doublet,  $\Psi_L^a$  to the left-handed fermions of the standard model (the index  $a$  runs over all the lepton and quark flavors) and  $W_\mu^i$  to the three electroweak

bosons. Note that while these objects are gauge invariant under local  $SU(2)_L$ , they still transform under the global  $SU(2)$  transformations:  $\Phi$  and  $\Psi_L$  are global  $SU(2)$  doublets while  $W_\mu^i$  are triplets under the global  $SU(2)$ . Using the unitarity gauge, which corresponds to the choice  $\Omega = 1$ , one finds  $\Phi \rightarrow \phi$ ,  $\Psi \rightarrow \psi$  and  $W_\mu^i \rightarrow B_\mu^i$ . Note that we could generalize this to locally invariant fields under the full gauge group of the standard model [8,9].

It is easy to reformulate the standard model in terms of these new fields. Starting from the standard model Lagrangian

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2}\text{Tr}F_{\mu\nu}F^{\mu\nu} - \frac{1}{4}f_{\mu\nu}f^{\mu\nu} + i\bar{\psi}_L^a \left( (\mathcal{D}_\mu)_{SU(2)} - i\frac{1}{2}g'Y\mathcal{A} \right) \psi_L^a \\ & + i\bar{\psi}_R^a (\not{\partial} - i\frac{1}{2}g'Y\mathcal{A})\psi_R^a \\ & + \left( \left( (\mathcal{D}_\mu)_{SU(2)} - i\frac{1}{2}g'Y\mathcal{A}_\mu \right) \phi \right)^\dagger \left( \left( (D^\mu)_{SU(2)} - i\frac{1}{2}g'Y\mathcal{A}^\mu \right) \phi \right) \\ & + V(\phi^\dagger\phi) - G_u(\bar{\psi}_L^a\phi\psi_R^a + \bar{\psi}_R^a\phi^\dagger\psi_L^a) - G_d(\bar{\psi}_L^a\bar{\phi}\psi_R^a + \bar{\psi}_R^a\bar{\phi}^\dagger\psi_L^a) \end{aligned} \quad (3)$$

where  $\bar{\phi} = i\tau_2\phi^*$ , the fermion  $\psi^a$  is a generic fermion field and the covariant derivative is given by  $(\mathcal{D}_\mu)_{SU(2)} = \partial_\mu - igB_\mu$ , with  $B_\mu = 1/2\tau^a B_\mu^a$ . We denote the Yukawa couplings by  $G_u$  and  $G_d$ . The gauge dependent fields can be replaced by their  $SU(2)_L$  gauge invariant counterparts. One obtains the dual standard model

$$\begin{aligned} \mathcal{L} = & -\frac{1}{2}\text{Tr}\mathcal{F}_{\mu\nu}\mathcal{F}^{\mu\nu} - \frac{1}{4}f_{\mu\nu}f^{\mu\nu} + i\bar{\Psi}_L^a \left( (\mathcal{D}_\mu)_{SU(2)_L} - i\frac{1}{2}g'Y\mathcal{A} \right) \Psi_L^a \\ & + i\bar{\psi}_R^a (\not{\partial} - i\frac{1}{2}g'Y\mathcal{A})\psi_R^a \\ & + \left( \left( (\mathcal{D}_\mu)_{SU(2)_L} - i\frac{1}{2}g'Y\mathcal{A}_\mu \right) \Phi \right)^\dagger \left( \left( (D^\mu)_{SU(2)_L} - i\frac{1}{2}g'Y\mathcal{A}^\mu \right) \Phi \right) \\ & + V(\Phi^\dagger\Phi) - G_u(\bar{\Psi}_L^a\Phi\psi_R^a + \bar{\psi}_R^a\Phi^\dagger\Psi_L^a) - G_d(\bar{\Psi}_L^a\bar{\Phi}\psi_R^a + \bar{\psi}_R^a\bar{\Phi}^\dagger\Psi_L^a). \end{aligned} \quad (4)$$

The scalar field potential is as usual

$$V(\phi^\dagger\phi) = \frac{1}{2}\lambda \left( \phi^\dagger\phi - \frac{1}{2}v^2 \right)^2, \quad (5)$$

its counterpart in term of  $SU(2)_L$  gauge invariant fields is given by

$$V(\Phi^\dagger\Phi) = \frac{1}{2}\lambda \left( \Phi^\dagger\Phi - \frac{1}{2}v^2 \right)^2. \quad (6)$$

This potential can be minimized if the field  $\Phi$  is forced to form the gauge invariant condensate

$$\langle \Phi^\dagger\Phi \rangle = \langle \phi^\dagger\phi \rangle = \frac{1}{2}v^2. \quad (7)$$

As in the standard model, the gauge invariant charged vector bosons receive a mass term of the form  $m_W = gv/2$ , the fermions receive masses of the type  $m_u = G_u v/\sqrt{2}$  for the up-type fermions and  $m_d = G_d v/\sqrt{2}$  for the down-type fermions. We also see that a term

$$\frac{1}{2}g'g\mathcal{A}_\mu W_\mu^3\Phi^\dagger\Phi \quad (8)$$

appears, which gives rise to a mixing between the  $U(1)$  generator and the  $W^3$  gauge invariant field. After diagonalization according to

$$\begin{aligned} A_\mu &= \sin\theta_W W_\mu^3 + \cos\theta_W \mathcal{A}_\mu \\ Z_\mu &= \cos\theta_W W_\mu^3 - \sin\theta_W \mathcal{A}_\mu, \end{aligned} \quad (9)$$

we can identify the photon  $A_\mu$  and the  $Z_\mu$  boson which are identical to the photon and  $Z$  boson of the standard model.

The dual standard model [3,4] is identical to the usual standard model and should not be confused with composite models of the electroweak interactions such as that of Abbott and Farhi [10–12]. While the dual formulation is physically equivalent to the standard model, it is a useful starting point to discuss potential physics effects beyond the standard model. In the sequel we depart from the dual formulation of the standard model and use it only as a guiding principle for physics beyond the standard model that is compatible with the  $SU(2) \times U(1)$  structure of the standard model. Note that if one takes the gauge invariant picture seriously there could be a relation between the Higgs boson’s mass and that of the three electroweak bosons, using this duality we were able to estimate the Higgs boson’s mass and obtained a prediction of 129.6 GeV [13] which is not far off from the measured Higgs boson’s mass at 125 GeV.

Using the dual picture of the standard model as a guiding principle, it is possible to construct more gauge invariant fields than those corresponding to the standard model field content. Obviously, it is not because once can write down a gauge invariant operator that there is a particle corresponding to that object, however this is a useful guiding principle to study potential candidates for the 750 GeV object. If the observation is confirmed, one of the operators we shall be discussing below could correspond to a particle of mass 750 GeV. For example, we can construct objects which are a superposition of gauge bosons which we shall call electroweak glue mesons<sup>2</sup> despite the fact that this is not a composite model. For example, since there are three gauge bosons in  $SU(2)_L$ , we can construct “glue mesons”

---

<sup>2</sup>A justification for this name is that the operators given in Eqs. (10) and (11) correspond in the unitary gauge respectively to  $F^{\mu\nu}F^{\alpha\beta}\epsilon_{\mu\nu\alpha\beta}$  and  $F_\mu^\alpha F_{\alpha\nu} - 1/4g_{\mu\nu}F^{\alpha\beta}F_{\alpha\beta}$  which have been used to represent true glue mesons in QCD [14]. Note that ’t Hooft introduced the terms weak mesons and weak baryons for the particles of the standard model within this dual picture [5].

containing two gauge bosons

$$S^{ij} = \frac{-1}{4g^2} \text{Tr} \left[ \Omega^\dagger \overset{\leftrightarrow}{D}_\mu \Omega \tau^i \Omega^\dagger \overset{\leftrightarrow}{D}^{\mu} \Omega \tau^j \right] \rightarrow B_\mu^i B^{j\mu} \quad (10)$$

$$T_{\mu\nu}^{ij} = \frac{-1}{4g^2} \text{Tr} \left[ \Omega^\dagger \overset{\leftrightarrow}{D}_{\{\mu} \Omega \tau^i \Omega^\dagger \overset{\leftrightarrow}{D}_{\nu\}} \Omega \tau^j \right] \rightarrow B_\mu^i B_\nu^j + B_\nu^i B_\mu^j \quad (11)$$

or in matrix form (using the standard notation  $W_\mu^\pm = (B_\mu^1 \mp iB_\mu^2)/\sqrt{2}$ ):

$$S = \tau^i S^{ij} \tau^j = \begin{pmatrix} B_\mu^3 B^{3\mu} + 2W_\mu^+ W^{-\mu} & \sqrt{2} B_\mu^3 W^{+\mu} - \sqrt{2} W_\mu^+ B^{3\mu} \\ \sqrt{2} W_\mu^- B^{3\mu} - \sqrt{2} B_\mu^3 W^{-\mu} & B_\mu^3 B^{3\mu} + 2W_\mu^+ W^{-\mu} \end{pmatrix}. \quad (12)$$

$$T_{\mu\nu} = \tau^i T_{\mu\nu}^{ij} \tau^j = \begin{pmatrix} B_{\{\mu}^3 B_{\nu\}}^3 + 2W_{\{\mu}^+ W_{\nu\}}^- & \sqrt{2} B_{\{\mu}^3 W_{\nu\}}^+ - \sqrt{2} W_{\{\mu}^+ B_{\nu\}}^3 \\ \sqrt{2} W_{\{\mu}^- B_{\nu\}}^3 - \sqrt{2} B_{\{\mu}^3 W_{\nu\}}^- & B_{\{\mu}^3 B_{\nu\}}^3 + 2W_{\{\mu}^+ W_{\nu\}}^- \end{pmatrix}. \quad (13)$$

There are thus three spin 0 states with QED charges 0 and  $\pm 1$  and three spin 2 states (d-waves [15]) with QED charges 0 and  $\pm 1$ . The charge 0 states are a superposition two electroweak boson  $ZZ$  or  $W^+W^-$  and the states carrying charges  $\pm 1$  are of the form  $ZW^+$  or  $ZW^-$ . It is obviously possible to define states built from more than two gauge bosons as well as states involving left-handed fermions

$$\Psi_L^{a*} = \frac{i}{2g} \Omega^\dagger \not{D} \psi_L^a \rightarrow \mathcal{B}^i \tau^i \psi_L^a. \quad (14)$$

Note that due to the chiral nature of the interactions of the standard model, it is not straightforward to write a mass term for the excited fermions in the effective Lagrangian. We could simply introduce a right-handed singlet  $\psi_R^{a*}$  in analogy to the fermions of the standard model. We will not investigate this question further here, but shall simply assume that if these fermions exist, they are very massive.

Because there are no states corresponding to  $S^{ij}$  and  $T_{\mu\nu}^{ij}$  in the standard model, we do not know how to couple these states to the particles of the standard model. The only guiding principle at our disposal is that we must preserve the approximate global  $SU(2)$  which is only broken by the  $U(1)_Y$  interaction. However, in the limit in which  $U(1)_Y$  decouples from the  $SU(2)_L$  sector, we must recover this symmetry. The new fermions  $\Psi_L^{a*}$  that are charged under  $SU(3)$  will behave as heavy quarks, they can also couple to the usual weak bosons of the standard model. They carry the hypercharge of their standard model counterparts and will thus couple to the hyperphoton. These new fermions could generate couplings between the gluons, the photon and electroweak bosons with the new scalar fields and the tensor modes which can be parametrized using effective field theory techniques. The coupling of the neutral scalar field to the spin 1 bosons of the standard model is given by

$$L_{spin0} = S^0 \sum_{a=1}^3 \frac{c_a^{(0)}}{\Lambda} \left( \frac{1}{4} {}^a F_{\alpha\beta} {}^a F^{\alpha\beta} \right) \quad (15)$$

where  $\Lambda$  is an energy scale probably of the order of the weak scale,  $c_a^{(0)}$  are Wilson coefficients,  $S^0$  is the neutral spin 0 state and  ${}^a F_{\alpha\beta}$  are the field strength tensors of the  $U(1)_Y$ ,  $SU(2)_L$  and  $SU(3)_c$  gauge fields. In the basis of the gauge boson mass eigenstates one finds:

$$L_{spin0} = \frac{S^0}{\Lambda} \left( \frac{c_{\gamma\gamma}^{(0)}}{4} A_{\alpha\beta} A^{\alpha\beta} + \frac{c_{\gamma Z}^{(0)}}{4} A_{\alpha\beta} Z^{\alpha\beta} + \frac{c_{ZZ}^{(0)}}{4} Z_{\alpha\beta} Z^{\alpha\beta} + \frac{c_{WW}^{(0)}}{4} \eta^{\mu\nu} W_{\alpha\beta} W^{\alpha\beta} + \frac{c_{gg}^{(0)}}{4} G_{\alpha\beta} G^{\alpha\beta} \right) \quad (16)$$

with

$$c_{\gamma\gamma}^{(0)} = c_1^{(0)} \cos^2 \theta_W + c_2^{(0)} \sin^2 \theta_W \quad (17)$$

$$c_{\gamma Z}^{(0)} = (c_1^{(0)} - c_2^{(0)}) \sin 2\theta_W \quad (18)$$

$$c_{ZZ}^{(0)} = c_1^{(0)} \sin^2 \theta_W + c_2^{(0)} \cos^2 \theta_W \quad (19)$$

$$c_{WW}^{(0)} = 2c_2^{(0)} \quad (20)$$

$$c_{gg}^{(0)} = c_3^{(0)}. \quad (21)$$

The coefficients  $c_i^{(0)}$  are free parameters.

For the spin 2 tensor, we have the following Lagrangian

$$L_{spin2} = T_{\mu\nu}^0 \sum_{a=1}^3 \frac{c_a^{(2)}}{\Lambda} \left( \frac{1}{4} \eta^{\mu\nu} {}^a F_{\alpha\beta} {}^a F^{\alpha\beta} - {}^a F_{\alpha}^{\mu} {}^a F^{\alpha\nu} \right) \quad (22)$$

where  $T_{\mu\nu}^0$  is the neutral spin 2 d-wave and as before  ${}^a F_{\alpha\beta}$  are the field strength tensors of the  $U(1)_Y$ ,  $SU(2)_L$  and  $SU(3)_c$  gauge fields. As before as well,  $\Lambda$  is an energy scale probably of the order of the weak scale,  $c_a^{(2)}$  are Wilson coefficients. In the basis of the gauge boson mass eigenstates one finds:

$$L_{spin2} = \frac{T_{\mu\nu}^0}{\Lambda} \left( c_{\gamma\gamma}^{(2)} \left( \frac{1}{4} \eta^{\mu\nu} A_{\alpha\beta} A^{\alpha\beta} - A_{\alpha}^{\mu} A^{\alpha\nu} \right) + c_{\gamma Z}^{(2)} \left( \frac{1}{4} \eta^{\mu\nu} A_{\alpha\beta} Z^{\alpha\beta} - A_{\alpha}^{\mu} Z^{\alpha\nu} \right) + c_{ZZ}^{(2)} \left( \frac{1}{4} \eta^{\mu\nu} Z_{\alpha\beta} Z^{\alpha\beta} - Z_{\alpha}^{\mu} Z^{\alpha\nu} \right) + c_{WW}^{(2)} \left( \frac{1}{4} \eta^{\mu\nu} W_{\alpha\beta} Z^{\alpha\beta} - A_{\alpha}^{\mu} Z^{\alpha\nu} \right) + c_{gg}^{(2)} \left( \frac{1}{4} \eta^{\mu\nu} G_{\alpha\beta} G^{\alpha\beta} - G_{\alpha}^{\mu} G^{\alpha\nu} \right) \right) \quad (23)$$

with

$$c_{\gamma\gamma}^{(2)} = c_1^{(2)} \cos^2 \theta_W + c_2^{(2)} \sin^2 \theta_W \quad (24)$$

$$c_{\gamma Z}^{(2)} = (c_1^{(2)} - c_2^{(2)}) \sin 2\theta_W \quad (25)$$

$$c_{ZZ}^{(2)} = c_1^{(2)} \sin^2 \theta_W + c_2^{(2)} \cos^2 \theta_W \quad (26)$$

$$c_{WW}^{(2)} = 2c_2^{(2)} \quad (27)$$

$$c_{gg}^{(2)} = c_3^{(2)}. \quad (28)$$

The coefficients  $c_i^{(2)}$  are free parameters.

The neutral spin 0 or spin 2 glue mesons could easily account for the recent events observed at CERN corresponding to a 750 GeV particle decaying to two photons. Let us first consider the neutral scalar field  $S^0$ , using the narrow width approximation one finds (see e.g. [16]):

$$\sigma(\text{proton proton} \rightarrow S^0 \rightarrow \gamma\gamma) = \sigma(\text{proton proton} \rightarrow S^0) \text{Br}(S^0 \rightarrow \gamma\gamma). \quad (29)$$

The production cross section at the parton level is given by

$$\hat{\sigma}(gg \rightarrow S^0) = \frac{(c_{gg}^{(0)})^2 \pi m_{S^0}^2}{4 \Lambda^2} \delta(\hat{s} - m_{S^0}^2). \quad (30)$$

Using the results of [17], it is straightforward to calculate the decay width of the neutral glue scalar meson to the electroweak bosons and gluons of the standard model:

$$\Gamma(S^0 \rightarrow \gamma\gamma) = \frac{(c_{\gamma\gamma}^{(0)})^2 m_{S^0}^3}{4\pi \Lambda^2} \quad (31)$$

$$\Gamma(S^0 \rightarrow gg) = 2 \frac{(c_{gg}^{(0)})^2 m_{S^0}^3}{\pi \Lambda^2} \quad (32)$$

$$\Gamma(S^0 \rightarrow \gamma Z) = \frac{1}{8} \frac{(c_{\gamma Z}^{(0)})^2 m_{S^0}^3}{\pi \Lambda^2} \left(1 - \frac{m_Z^2}{m_{S^0}^2}\right)^3 \quad (33)$$

$$\Gamma(S^0 \rightarrow WW) = 64\pi^3 (c_{WW}^{(0)})^2 \frac{m_{S^0}^5}{\Lambda^2 m_W^2} \sqrt{1 - 4 \frac{m_W^2}{m_{S^0}^2}} \left(1 - 4 \frac{m_W^2}{m_{S^0}^2} + 12 \frac{m_W^4}{m_{S^0}^4}\right) \quad (34)$$

$$\Gamma(S^0 \rightarrow WW) = 32\pi^3 (c_{ZZ}^{(0)})^2 \frac{m_{S^0}^5}{\Lambda^2 m_Z^2} \sqrt{1 - 4 \frac{m_Z^2}{m_{S^0}^2}} \left(1 - 4 \frac{m_Z^2}{m_{S^0}^2} + 12 \frac{m_Z^4}{m_{S^0}^4}\right). \quad (35)$$

Similarly, for the neutral d-wave could be produced as a resonance and the narrow width approximation can be used to calculate the cross section  $\sigma(\text{proton proton} \rightarrow T^0 \rightarrow \gamma\gamma)$ . The widths can easily be calculated. One has [18]:

$$\Gamma(T_{\mu\nu}^0 \rightarrow \gamma\gamma) = \frac{(c_{\gamma\gamma}^{(2)})^2 m_{T_{\mu\nu}^0}^3}{80\pi \Lambda^2} \quad (36)$$

$$\Gamma(T_{\mu\nu}^0 \rightarrow Z\gamma) = \frac{(c_{\gamma Z}^{(2)})^2 m_{T_{\mu\nu}^0}^3}{160\pi \Lambda^2} \left(1 - \frac{m_Z^2}{m_{T_{\mu\nu}^0}^2}\right)^3 \left(1 + \frac{m_Z^2}{2m_{T_{\mu\nu}^0}^2} + \frac{m_Z^4}{6m_{T_{\mu\nu}^0}^4}\right) \quad (37)$$

$$\Gamma(T_{\mu\nu}^0 \rightarrow ZZ) = \frac{(c_{ZZ}^{(2)})^2 m_{T_{\mu\nu}^0}^3}{80\pi \Lambda^2} \sqrt{1 - 4 \frac{m_Z^2}{m_{T_{\mu\nu}^0}^2}} \left(1 - 3 \frac{m_Z^2}{m_{T_{\mu\nu}^0}^2} + 6 \frac{m_Z^4}{m_{T_{\mu\nu}^0}^4}\right) \quad (38)$$

$$\Gamma(T_{\mu\nu}^0 \rightarrow WW) = \frac{(c_{WW}^{(2)})^2 m_{T_{\mu\nu}^0}^3}{160\pi \Lambda^2} \sqrt{1 - 4 \frac{m_W^2}{m_{T_{\mu\nu}^0}^2}} \left(1 - 3 \frac{m_W^2}{m_{T_{\mu\nu}^0}^2} + 6 \frac{m_W^4}{m_{T_{\mu\nu}^0}^4}\right) \quad (39)$$

$$\Gamma(T_{\mu\nu}^0 \rightarrow gg) = \frac{(c_{gg}^{(2)})^2 m_{T_{\mu\nu}^0}^3}{10\pi \Lambda^2} \quad (40)$$



Clearly, the 750 GeV excess could easily be explained by either the production of the spin 0 or spin 2 glue mesons discussed above. The analysis performed in e.g. [19–22] of the 750 GeV resonance as a spin 0 or spin 2 particle applies to our model. In particular the absence of the observation of the final state  $\gamma Z$  so far implies that the Wilson coefficient  $c_{\gamma Z}^{(0/2)}$  is smaller than  $c_{\gamma\gamma}^{(0/2)}$ , see e.g. [23] for a recent review which discusses the most recent bounds on the Wilson coefficients. Because of the global  $SU(2)$  symmetry, we know that besides the neutral spin 0 or spin 2 particles, we are expecting a charged counterpart with a mass close to 750 GeV modulo some small quantum electrodynamics effects in analogy to the mass difference between the charge electroweak bosons and the Z-bosons. The charged glue mesons are however more difficult to produce than their neutral counterparts as this would have happen from the fusion of electroweak bosons. Also their decays principally in charged fermions via  $S/T \rightarrow W + Z \rightarrow 4$  fermions are less straightforward than the clear  $\gamma\gamma$  signature of decay of a neutral glue meson.

If the 750 GeV resonance is confirmed by future measurements at the LHC, the dual standard model and its extension as described above is a very natural framework to account for this effect. Within that model one expects for three spin 0 and spin 2 glue mesons, there are charged  $SU(2)$  partners corresponding to the neutral resonance which may have been recently observed. Clearly if this picture is correct, higher spin resonances in  $SU(2)$  multiplets are also expected as well as more exotic particles corresponding to "excitations" of the known Higgs boson and fermions. While the Lagrangian proposed above is not renormalizable in the usual sense, it could be the first sign that asymptotic safety [24] is truly relevant to the electroweak interactions.

*Acknowledgments:* This work is supported in part by the Science and Technology Facilities Council (grant number ST/L000504/1).

## References

- [1] CMS Collaboration [CMS Collaboration], "Search for new physics in high mass diphoton events in proton-proton collisions at 13TeV," CMS-PAS-EXO-15-004.
- [2] The ATLAS collaboration, "Search for resonances decaying to photon pairs in  $3.2 \text{ fb}^{-1}$  of  $pp$  collisions at  $\sqrt{s} = 13 \text{ TeV}$  with the ATLAS detector," ATLAS-CONF-2015-081.
- [3] X. Calmet and H. Fritzsch, Phys. Lett. B **496**, 161 (2000) [arXiv:hep-ph/0008243].
- [4] X. Calmet, "A duality as a theory for the electroweak interactions," arXiv:hep-ph/0206251, 2002, Ph.D. Thesis, Shaker Verlag Aachen, ISBN 3-8322-0324-9.
- [5] G. 't Hooft, "Topological aspects of quantum chromodynamics", Lectures given at International School of Nuclear Physics: 20th Course: Heavy Ion Collisions from Nuclear

- to Quark Matter (Erice 98), Erice, Italy, 17-25 Sep 1998, “Topological aspects of quantum chromodynamics,” hep-th/9812204, published in “Erice 1998, From the Planck length to the Hubble radius,” 216-236.
- [6] G. 't Hooft, in “Recent Developments In Gauge Theories,” Cargèse 1979, ed. G. 't Hooft et al. Plenum Press, New York, 1980, Lecture II, p.117, “Recent Developments In Gauge Theories. Proceedings, Nato Advanced Study Institute, Cargèse, France, August 26 - September 8, 1979.”
- [7] V. Visnjic, *Nuovo Cim. A* **101**, 385 (1989).
- [8] M. Lavelle and D. McMullan, *Phys. Rept.* **279**, 1 (1997) [arXiv:hep-ph/9509344].
- [9] X. Calmet, *Int. J. Mod. Phys. A* **26**, 2855 (2011) doi:10.1142/S0217751X11053699 [arXiv:1008.3780 [hep-ph]].
- [10] L. F. Abbott and E. Farhi, *Phys. Lett.* **B101** 69 (1981).
- [11] L. F. Abbott and E. Farhi, *Nucl. Phys.* **B189** 547 (1981).
- [12] M. Claudson, E. Farhi and R. L. Jaffe, *Phys. Rev.* **D34** 873 (1986).
- [13] X. Calmet and H. Fritzsche, *Phys. Lett. B* **525**, 297 (2002) doi:10.1016/S0370-2693(01)01449-6 [hep-ph/0107085].
- [14] H. Fritzsche and P. Minkowski, *Nuovo Cim. A* **30**, 393 (1975). doi:10.1007/BF02730295
- [15] X. Calmet and H. Fritzsche, *Phys. Lett. B* **526**, 90 (2002) doi:10.1016/S0370-2693(01)01471-X [hep-ph/0103333].
- [16] A. Djouadi, *Phys. Rept.* **457**, 1 (2008) doi:10.1016/j.physrep.2007.10.004 [hep-ph/0503172].
- [17] I. Low, J. Lykken and G. Shaughnessy, *Phys. Rev. D* **84**, 035027 (2011) doi:10.1103/PhysRevD.84.035027 [arXiv:1105.4587 [hep-ph]].
- [18] T. Han, J. D. Lykken and R. J. Zhang, *Phys. Rev. D* **59**, 105006 (1999) doi:10.1103/PhysRevD.59.105006 [hep-ph/9811350].
- [19] J. Ellis, S. A. R. Ellis, J. Quevillon, V. Sanz and T. You, *JHEP* **1603**, 176 (2016) doi:10.1007/JHEP03(2016)176 [arXiv:1512.05327 [hep-ph]].
- [20] R. S. Gupta, S. Jger, Y. Kats, G. Perez and E. Stamou, arXiv:1512.05332 [hep-ph].
- [21] C. Han, H. M. Lee, M. Park and V. Sanz, *Phys. Lett. B* **755**, 371 (2016) doi:10.1016/j.physletb.2016.02.040 [arXiv:1512.06376 [hep-ph]].
- [22] V. Sanz, arXiv:1603.05574 [hep-ph].
- [23] A. Strumia, arXiv:1605.09401 [hep-ph].
- [24] X. Calmet, *Mod. Phys. Lett. A* **26**, 1571 (2011) doi:10.1142/S0217732311035900 [arXiv:1012.5529 [hep-ph]].