

## Stable isotope geochemistry of massive ice

Article (Accepted Version)

Vasil'chuk, Yuriy K and Murton, Julian (2016) Stable isotope geochemistry of massive ice. *Geography, Environment, Sustainability*, 9 (3). pp. 4-24. ISSN 2071-9388

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/63781/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

### **Copyright and reuse:**

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

## Monograph Synopsis

**Stable Isotope Geochemistry of Massive Ice. 2 Vols. by Yuriy K. Vasil'chuk, Moscow University Press, Moscow, Vol. I. 2012. 472 pp. ISBN 978-5-211-06338-4 & Vol. II. 2014. 244 pp. ISBN 978-5-19-010897-2**

*Yuriy K. Vasil'chuk<sup>1\*</sup> and Julian B. Murton<sup>2</sup>*

<sup>1\*</sup>Faculty of Geography and Faculty of Geology, Lomonosov Moscow State University, Moscow, Russia

<sup>2</sup>Permafrost Laboratory, Department of Geography, University of Sussex, Brighton, UK.

\*Correspondence to: Yu.Vasil'chuk, Professor of Geocryology and Isotope Geochemistry, Faculty of Geography, Lomonosov Moscow State University (MSU), Russia 119991, Moscow, Leninskie Gory, 1, vasilch@geol.msu.ru & vasilch\_geo@mail.ru

### ABSTRACT

The monograph summarises stable-isotope research on massive ice in the Russian and North American Arctic, and includes the latest understanding of massive-ice formation. A new classification of massive-ice complexes is proposed, encompassing the range and variability of massive ice. It distinguishes two new categories of massive-ice complexes: *homogeneous* massive-ice complexes have a similar structure, properties and genesis throughout, whereas *heterogeneous* massive-ice complexes vary spatially (in their structure and properties) and genetically within a locality and consist of two or more homogeneous massive-ice bodies. Analysis of pollen and spores in massive ice from Subarctic regions and from ice and snow cover of Arctic ice caps assists with interpretation of the origin of massive ice. Radiocarbon ages of massive ice and host sediments are considered together with isotope values of heavy oxygen and deuterium from massive ice plotted at a uniform scale in order to assist interpretation and correlation of the ice. The monograph is intended for both undergraduates and graduate students, and will assist researchers in geocryology, glaciology, geomorphology, Quaternary geology and palaeoclimatology in understanding of the origin and palaeoenvironmental significance of massive ice.

**KEY WORDS:** massive ice, stable isotopes, radiocarbon dating, homogeneous and heterogeneous

### INTRODUCTION

The research for the monograph began in July 1977 during the boating expedition of the Geological Faculty of Moscow State University in the Yuribey River valley (68°26'N, 72°08'E) on central Yamal Peninsula, west Siberia. Here, the senior author encountered a complex exposure of massive-ice bodies that probably represent an assemblage of buried river ice and injection (intrusive) ice. This exposure

has contributed significantly to the development of a new massive-ice classification by Y. K. Vasil'chuk, the basic principles of which are set out in Chapter 1 (Figure 1).

## **STUDY AREA**

The monograph is based primarily on the senior author's experience, involvement and field studies over more than 35 years (from 1977 to 2014) in numerous expeditions concerning massive ice in Russian permafrost. It also summarizes the experience of many international researchers, and describes exposures with large bodies of massive ice in west and east Siberia, Chukotka, Alaska and Yukon, the Tuktoyaktuk Coastlands, the Canadian Arctic Archipelago and Russian Arctic islands, China and the Antarctic.

## **CHAPTER 1: MASSIVE-ICE CLASSIFICATION**

Chapter 1 presents a new classification of massive-ice bodies and includes two new categories: homogeneous and heterogeneous (Figure 1). *Homogeneous* massive-ice bodies have a similar genesis, composition and properties in all parts of a massive-ice complex, whereas *heterogeneous* massive-ice bodies have a variable genesis, composition and properties across a massive-ice complex, and consist of two or more homogeneous ice bodies. The distinction between homogeneous and heterogeneous massive-ice bodies elucidates the wider complex structure of massive-ice bodies and encourages the search for different mechanisms of ice formation.

### **Homogeneous Massive-ice Complexes**

Homogeneous massive-ice complexes are usually no more than a few meters high,  $\leq 20\text{--}30$  m wide and occur as single layers of ice or, less commonly, as multiple layers of the same genesis. Typical examples of segregated (or infiltrated and segregated) ice were studied by Y. K. Vasilchuk (1992) in the first terrace of the Gyda River estuary ( $70^{\circ}53'N$ ,  $78^{\circ}30'E$ ). Four similar lens-shaped bodies of massive ice (up to 0.3–0.4 m thick and 6–8 m wide) were composed of clear ice and, as a rule, associated with peat. The structure of the ice bodies and their bedding parallel to the sedimentary stratification suggest that they formed synchronously with the accumulation and freezing of the ground mass, consistent with the occurrence of a 4.5 m high syngenetic ice wedge in the sedimentary sequence. The ice is interpreted as infiltrated or segregated in origin. The wide range of  $\delta^{18}O$  values ( $-33.8$  to  $-16.2\text{‰}$ ) in these ice bodies indicates closed-system freezing with a small inflow of water from outside the system. Such a wide range of the heavy oxygen values indicates significant cryogenic fractionation during freezing of waters whose initial average composition was close to  $-20\text{‰}$ . This

could occur only under conditions of closed-system freezing, when isotopically heavier ice was the first to form ( $\delta^{18}\text{O}$  values of about  $-16$  and  $-18\text{‰}$ ), while the  $\delta^{18}\text{O}$  values of the remaining water were about  $-22\text{‰}$ . The partial freezing of the water led to the formation of ice with  $\delta^{18}\text{O}$  values of about  $-20\text{‰}$ , while the rest of the water had  $\delta^{18}\text{O}$  values of about  $-24$  to  $-25\text{‰}$ . Repeated freezing of this water provided extremely low  $\delta^{18}\text{O}$  values (about  $-34\text{‰}$ ) in the last portions of the water to freeze. The low average  $\delta^{18}\text{O}$  values in this massive ice (about  $-20\text{‰}$ ) indicate that the ice formed under conditions more severe than those at present. In view of the radiocarbon age of the ground mass (more than  $15^{14}\text{C}$  ages from 10,260 to 15,890 ka BP obtained on allochthonous peat), the accumulation of the ground mass and the formation of massive ice and the syngenetic ice wedge occurred no earlier than 14,000–11,000 years ago (Y. K. Vasilchuk, 1992).

### **Heterogeneous massive-ice complexes**

Heterogeneous massive-ice complexes comprise two or more ice layers (tiers) of different genesis. The layers may be in contact or adjacent to each other, influencing the shape of ice complex and the conditions of its occurrence. Examples include massive ice of the Yuribey River valley (Figure 2a; Y. K. Vasil'chuk *et al.*, 2012), massive ice in the Erkutayaha River valley in the south of Yamal Peninsula (Figure 2b), and in many places in the Bovanenkovo gas condensate field (Y. K. Vasil'chuk *et al.*, 2009, 2014; Kritsuk, 2012).

The second division of the classification distinguishes between *autochthonous* (i.e., *intrasedimental* in terms of Mackay and Dallimore, 1992) and *allochthonous* (i.e. buried) ice (Figure 1). Heterogeneous massive-ice complexes can include a combination of autochthonous and allochthonous deposits. The third division classifies the massive ice according to its specific genetic process (e.g., injection, segregation, infiltration, burial) and shows the wide diversity of genetic ice types.

The proposed classification is illustrated through examples in chapter 1, and more widely in chapter 2, which critically examines the main hypotheses of massive-ice formation.

## **CHAPTER 2. MODERN AND HOLOCENE ANALOGUES OF PLEISTOCENE MASSIVE ICE, AND THE MAIN HYPOTHESES OF MASSIVE-ICE FORMATION**

Chapter 2 discusses modern and Holocene analogues of Pleistocene massive ice, beginning with massive ice in marine sediments. One of the most convincing analogues of Pleistocene massive ice is Holocene massive segregated ice 8–9 m thick on the Fosheim Peninsula, Ellesmere Island, in the Canadian Arctic ( $79^{\circ}59'\text{N}$ ,  $85^{\circ}56'\text{W}$ ; Pollard and Bell, 1998). The  $\delta^{18}\text{O}$  values of the ice range

between  $-28.9$  and  $-34.8\text{‰}$  for reticulate ice,  $-33.1$  and  $-36.8\text{‰}$  for massive ice, and  $-36.1\text{‰}$  for an ice vein in adjacent Tertiary sandstone. The massive ice is interpreted as intrasedimental ice because: a) it is conformably overlain by marine sediments and contains internal structures parallel the upper ice contact; the ice lacks evidence for primary thaw or erosional contacts, which would indicate buried ice; b) sediment within inclusions in the massive ice is similar to the overlying marine sediments, indicating that the overlying sediment was deposited before the ice formed. Radiocarbon ages of shells from raised marine deposits in the Slidre River valley suggest that the Holocene marine limit was likely established by 10.6 ka BP, with sea level remaining high (within 10 m of marine limit) until 8.7 ka BP or later. The massive-ice deposits formed time transgressively as permafrost aggraded into marine sediments during the Holocene (Pollard and Bell, 1998). Ice segregation occurred as permafrost aggraded downward through a fine-grained layer of silt that overlay saturated sands with an ample water source. The injection of water under high pressure into the overlying frozen bedrock at Eureka Sound formed intrusive massive ice (Robinson and Pollard, 1998). As a result, a heterogeneous massive-ice complex (according to Y. K. Vasil'chuk's classification) developed.

Massive-ice bodies are extraordinarily widespread in Holocene sediments of the first lagoon-marine terrace and the modern lagoon-marine floodplain of Ob Bay, at the mouth of the Sabettayaha River, in west Siberia ( $71^{\circ}15'N$ ,  $72^{\circ}06'E$ ). More than 1000 boreholes through Holocene marine lagoon and floodplain deposits have been analyzed (Y. K. Vasilchuk *et al.*, 2015) and show massive-ice bodies up to 5.7 m thick in the upper 5–10 m of Holocene silty sand. Their  $\delta D$  values vary from  $-107$  to  $-199.7\text{‰}$ , and their  $\delta^{18}O$  values vary from  $-15.7$  to  $-26.5\text{‰}$ . The massive ice is mostly autochthonous and of segregated origin (Y. K. Vasil'chuk *et al.*, 2015). It formed syngenetically during freezing of water-saturated soils under intensive cryogenic fractionation in the late Holocene. Thick massive-ice bodies also occur in saline sediments, for example a 14 m thick massive ice layer in beach sediments near the mouth of the Khatanga River (Ponomarev, 1960), fresh (salinity is 0.62 g / l) massive ice at a depth of 19–29 m beneath the sea bottom (Melnikov and Spesivtsev, 2000), and a 1.5 m thick layer of salty ice at the depth about 10 m in sediments beneath the shallow waters of the Mechigmen Gulf (Chukotka). Y.K. Vasil'chuk proposed the original mechanism of the formation of icy bodies in saline lake (meromictic lakes) sediments.

Other possible modern and Holocene analogues of Pleistocene massive ice are freshwater ice deposits in shallow, saline (up to 103 g/l) lakes at 4117–4730 m above sea level (asl) in Bolivia. The ice deposits (consisting of several ice lenses, each up to 1 m thick) are up to several hundred meters wide and elevated up to 7 m above the lake or playa surface. They are located near the lake or salar

(salt-crust desert) margins; some are completely surrounded by water, others by playa deposits or salt crusts (Hurlbert and Chang, 1984). The  $\delta^{18}\text{O}$  values for the ice lenses ( $-10.6$  to  $-11.5\text{‰}$ ) are much lower than those of lakewater ( $+13.5\text{‰}$ ) but similar to those of precipitation ( $\delta^{18}\text{O}$  value in fresh snow  $-13.2\text{‰}$ ).

Ice in the cores of modern pingos provides another analogue of Pleistocene massive-ice deposits. The distribution of stable isotopes in pingo ice cores indicates that the ice formed under conditions of open and closed systems. Ice in the core of the Holocene pingo in the Pestsovoe gas field area ( $66^{\circ}10' \text{ N } 76^{\circ}30' \text{ E}$ ) formed in closed-system conditions; its  $\delta\text{D}$  values vary from  $-93.2$  to  $-123\text{‰}$ , and  $\delta^{18}\text{O}$  values vary from  $-11.6$  to  $-15.8\text{‰}$  (Y. K. Vasil'chuk *et al.*, 2014).

The main hypotheses of massive-ice formation identify the ice as segregated, intrusive, repeated intrusive and buried glacier ice (Vtyurin, 1975; Mackay, 1973; Rampton, 1988; Zhestkova and Shur, 1978; Y. K. Vasil'chuk, 1992, 2012, 2014; Dubikov, 2002; Murton, 2005, 2009; French, 2007; Solomatin, 2013; Streletskaya *et al.*, 2013; Belova, 2014). Holmsen (1914) proposed the infiltrated and segregated hypothesis of massive-ice genesis. He hypothesized that the formation of thick (up to 15 m) massive ground-ice deposits is linked to the infiltration of surface water through seasonally thawed ground and the freezing of this water on the top (see also Zhestkova and Shur, 1978). Vtyurin (1975) favoured ice segregation, with the most favorable conditions for the formation of massive segregated ice being near the contact between clayey sediments and water-bearing coarse-grained sediments. Gasanov (1969) hypothesized that the main factor of ice formation is water intrusion, and distinguished different types of intrusive ice: seasonal intrusive ice, multi-seasonal intrusive ice (short-term permafrost), intrusive ice, repeated intrusive ice and hydrolaccoliths. Mackay (1971, 1973) proposed that the mechanism of water injection and segregation in closed-system conditions that is widely applied to explain pingo growth can also apply to the mechanism of massive-ice formation.

### **CHAPTER 3. COMPARISON OF POLLEN SPECTRA OF MASSIVE ICE AND GLACIERS FOR CRYOGENIC INDICATION**

Palynological analysis of deposit-forming ground ice has identified several distinctive characteristics of the pollen spectra. First, pollen and spores are present in almost in all types of deposit-forming ground ice, at concentrations from 50 to 1500 units per 1 kg ice or 1 litre of ice meltwater. Second, pollen spectra with characteristics similar to those of subfossil tundra, including the predominance of dwarf birch and ericaceous pollen and green moss spores, occur in most massive-ice deposits. Third, the massive-ice deposits frequently contain redeposited pre-Quaternary palynomorphs of Cenozoic, Mesozoic and Palaeozoic age. Fourth, pollen of hydrophilous plants (e.g., pondweed, bur

reed and reed mace), as well as horsetail spores, limnetic diatoms and green algae remains also occur in most of the studied massive-ice deposits, indicating a non-glacial genesis of the ice. Pollen spectra that are typical of non-glacial deposit-forming ice show: 1) a lack of exotic thermophilic species such as *Acer*, *Fraxinus*, *Quercus*, *Ulmus*, *Populus*, *Tilia* and *Abies* in the initial occurrence; 2) the presence of cloudberry, aquiferbosa species, as well as green moss and horsetail; and 3) the presence of redeposited pollen and spores. This palynological approach indicates that the massive ice on the Yamal Peninsula of west Siberia is mainly non-glacial in origin (A. C. Vasil'chuk and Y. K. Vasul'chuk, 2010, 2012).

#### **CHAPTER 4. ISOTOPIC COMPOSITION OF MASSIVE ICE IN RUSSIAN PERMAFROST**

In chapter 4 the senior author considers his own isotopic studies of massive ice in Siberia and interprets the isotope records obtained by others. He considers the vertical variations in more than 35 isotope plots published in the last 30 years (Vaikmäe and Karpov, 1986; Y. K. Vasil'chuk and Trofimov, 1988; Vaikmäe and Y. K. Vasil'chuk, 1991; Korolyov, 1993; Kotov, 1998; Dubikov, 2002; Leibman *et al.*, 2003; Ingólfsson and Lokrantz, 2003; Y. K. Vasil'chuk *et al.*, 2009, 2011, 2012, 2014; Kritsuk, 2010; Slagoda *et al.*, 2012; Ivanova, 2012; Belova, 2014). The isotope curves are plotted at a single vertical and horizontal scale to facilitate their comparison.

The isotope composition of the massive-ice deposits in northern European Russia is considered for a number of localities, including the More-Yu River valley, Cape Shpindler, and the Oyuyakha River valley (68°51'N, 66°44'E). For the massive-ice body at Cape Shpindler in the Yugorski Peninsula (69°43'N; 62°48'E), Ingólfsson and Lokrantz (2003) concluded that it is buried glacier ice and suggested that it is older than 190–200 kyr, whereas Leibman *et al.* (2003), who also examined the internal structures, stratigraphy and isotopic composition of the massive ice, attributed it to syngenetic or epigenetic freezing after marine regression. The senior author assumes that this massive ice is intrasedimental, heterogeneous and autochthonous because the  $\delta^{18}\text{O}$  values in different types of massive ice at Cape Spindler vary from –13.1 to –25.6‰. Such variability may be explained by fractionation during ice segregation in a closed system.

The isotope composition of massive-ice deposits is also considered for northwest Siberia. Localities there include: 1) the Erkutayaha River valley (68°11'N, 68°51'E), 2) the Bovanenkovo gas field area (70°21'N, 68°26'E), 3) the Mordyyakha River valley (69°33'N, 68°59'E), 4) the Kharasavey settlement (71°10'N, 66°51'E), 5) near Marre-Sale meteorostation (69°45'N; 66°50'E), 6) at Tyurinto Lake (70°44'N, 67°57'E), 7) at Voivareto Lake (68°43'N; 72°25'E), 8) near Gyda settlement (70°53'N, 78°30'E), 9) the Yuribey River valley (68°26'N, 72°08'E), 10) near Tab-Salya settlement

(71°45'N; 82°45'E), 11) near Dorofeevskaya settlement (71°23'N; 82°58'E), 12) at the Sopochnaya Karga Cape (71°50'N, 82°40'E) and 13) Ledyanaya Gora in the Yenisey River valley (66°35'N, 86°34'E).

On the southernmost Yamal Peninsula heterogeneous autochthonous massive ice is located on the left bank of the Erkutayaha River (68°11'N, 68°51'E). A massive-ice body approximately 100 m long is embedded predominantly in stratified sand. The ice layers sharply drop on both sides of the central part, and just 15 m from the centre the cover of the massive ice appears at a depth of 8 m. The central part of the massive-ice body takes a form similar to a stock (a type of igneous intrusion), with vertical and subvertical ice layers, whereas the peripheral parts comprise horizontally layered ice. The range of  $\delta^{18}\text{O}$  (~4‰) and  $\delta\text{D}$  (~20‰) values indicates comparatively small fluctuations of the isotopic composition for ice with different characteristics: *pure white ice* has  $\delta^{18}\text{O}$  values from -19.6 to -20.5‰, and  $\delta\text{D}$  varies from -152.4 to -156.9‰; *clear transparent ice* has  $\delta^{18}\text{O}$  values from -19.2 to -20.3‰, and  $\delta\text{D}$  varies from -149.6 to -160.7‰; *transparent gray ice* with a steely shimmer has  $\delta^{18}\text{O}$  values from -19.4 to -21.3‰, and  $\delta\text{D}$  varies from -150.3 to -163.8‰; and *gray ice and in the dirty gray ice* has  $\delta^{18}\text{O}$  values from -22.1 to -23.4‰, and  $\delta\text{D}$  varies from -165.5 to -172.7‰. Hence, the difference between the isotope signal of the initial water and ice is comparatively small. The isotopic differences almost did not exceed the usual isotopic difference that results from fractionation when free water freezes. Significantly, the pollen spectra in the vertically layered ice from the central stock (most likely of injection genesis) lack reworked pollen and spores, whereas those from the horizontally layered peripheral ice (most likely of segregation or infiltration-segregation genesis) contain a high concentration of reworked pollen and spores (35%).

On the central part of the Yamal Peninsula (69°33'N, 68°59'E) beside the upper Mordyyakha River is a heterogeneous autochthonous massive-ice body at 66–70 m asl. The ice body is more than 4 m thick and has horizontal bedding that passes laterally into vertical bedding and is crossed by syngenetic ice wedges 4–5 m in high. The isotopic composition of ice from obliquely oriented ice lenses (lenticular-layered cryostructure) in the north part of the outcrop shows insignificant isotopic variations (from -22.4 to -23.3‰  $\delta^{18}\text{O}$ ) and thus suggests that ice segregation occurred under open-system conditions. The pollen spectra are characterized by high contents of pollen and spores of hygrophilous plants, with sedge pollen and horsetail spores dominant. Such specific features were noted for sediments of small floodplain lakes. It can be assumed that the ice deposits formed during freezing of a talik after a former lake drained.

In the Bovanenkovo gas field area (70°21'N, 68°26'E) heterogeneous autochthonous and allochthonous massive-ice bodies occur as layers, laccoliths, rods and lenses. The maximum thickness

of the tabular ice is 28.5 m, and the mean thickness is about 8 m.  $\delta^{18}\text{O}$  values of massive ice range from  $-12.4$  to  $-22.9\text{‰}$  (Michel, 1998; Y. K. Vasil'chuk *et al.*, 2009), and deuterium ( $\delta\text{D}$ ) values vary from  $-91.7$  to  $-177.1\text{‰}$ . The contrasting distribution of  $\delta^{18}\text{O}$  and ( $\delta\text{D}$ ) vs. depth (like a vertical bilateral rake with very long teeth of different length) in massive ice bodies suggest a segregated and/or infiltrated-segregated origin of the ice. Pollen, spores and algal spectra from the massive ice are similar to pollen characteristics of modern lacustrine and coastal floodplain sediments in the area. The senior author inferred that the ingress of cold seawaters on a coastal flood plain caused freezing and ice segregation, with the formation of extensive ice layers under the large but shallow lakes.

The material presented shows that the methodological use of isotopic data from massive ice is still far from ideal. Even within the same article, the authors have expressed different hypotheses about the origin of massive ice in Ledyanaya Gora (Ice Mountain) on the Yenisei River valley ( $66^{\circ}35'\text{N}$ ,  $86^{\circ}34'\text{E}$ ) because intrasedimental ice may be difficult to distinguish from basal glacier ice, as both ice types can form by the same freezing processes (Vtyurin and Glazovskiy, 1986). Stratigraphic and isotopic study of this massive ice and its host sediments led Vaikmäe and Karpov (1986) and Astakhov and Isayeva (1988) to interpret the ice is relict glacier ice, probably emplaced during the Early Weichselian. However, interpretation of the Ledyanaya Gora massive ice is problematic. Most of the  $\delta^{18}\text{O}$  values from the ice range from  $-20$  to  $-21.5\text{‰}$  (Vaikmäe and Y. K. Vasil'chuk, 1990; Vaikmäe *et al.*, 1993), pointing to uniformity of the oxygen isotope profile. The isotopic composition of the massive ice is similar to that of vertical ice schlieren in the overlying diamicton ( $\delta^{18}\text{O} = -20.7\text{‰}$ ). This similarity, coupled with a significant change from a salinity of  $10\text{--}80\text{ mg / l}$  in the upper part of the massive ice to one of  $200\text{--}340\text{ mg / l}$  in the middle and lower parts of it, according Y. K. Vasil'chuk, indicates an intrasedimental origin. In conclusion, the massive ice at Ledyanaya Gora can be regarded as heterogeneous autochthonous massive ice.

Isotope data summarized from Chukotka include homogeneous autochthonous massive ice at the Koolen' Lake, near the town of Anadyr' and at Onemen Bay (Vasil'chuk, 1992), Tanyurer River valley (Kotov, 1998), and heterogeneous allochthonous and autochthonous massive ice in the Anguema River valley (Korolyov, 1993).

Finally, heterogeneous allochthonous and autochthonous massive-ice bodies are found on Novaya Siberia Island ( $75^{\circ}05'\text{N}$ ,  $148^{\circ}27'\text{E}$ ). Ivanova (2012) showed that  $\delta^{18}\text{O}$  values of massive ice there vary from  $-8.9$  to  $-29\text{‰}$ , and  $\delta\text{D}$  values vary from  $-66.8$  to  $-228.4\text{‰}$ . The  $\delta^{18}\text{O}$  and  $\delta\text{D}$  values in the upper and lower horizons of massive ice are very different. These data can be interpreted to indicate

an injected origin for the lower horizon of massive ice, and a segregation origin for the upper horizon (Ivanova, 2012).

## **CHAPTER 5. ISOTOPIC COMPOSITION OF MASSIVE ICE IN PERMAFROST OF THE CANADIAN ARCTIC**

Chapter 5 examines the vertical variations in more than 30 isotope plots of massive ice from the Canadian Arctic (Mackay, 1983; Fujino *et al.*, 1983; Lorrain and Demeur, 1985; Michel, 1983, 2011; Dallimore and Wolfe, 1988; French and Harry, 1990; Mackay and Dallimore, 1992; Moorman *et al.*, 1996, 1998; Robinson and Pollard, 1998; Pollard and Bell, 1998; Hyatt, 1998; Murton *et al.*, 2004, 2005; Lacelle *et al.*, 2004, 2007, 2009, 2011; Murton, 2005, 2009; Cardyn *et al.*, 2007; French, 2007; Lacelle, 2011; Fritz *et al.*, 2011). Here we consider a few of them.

Massive ice adjacent to Tuktoyaktuk (69°27'N, 133°02'W), near the mouth of the Mackenzie River, has been isotopically studied by Mackay (1983), Fujino *et al.* (1983, 1988), Mackay and Dallimore (1992) and others. Fujino *et al.* (1983) concluded that most of the massive-ice body at Peninsula Point originated from superimposed ice formed by congelation of water in which it was submerged. Mackay and Dallimore (1992), however, rejected this interpretation. They reported that the geochemical and stable isotope values of the massive ice were similar to those of ice dikes that extended from the ice into the overlying diamicton, which indicates a common water source and suggests that the massive ice is intrasedimental in origin. Y. K. Vasil'chuk concluded that massive ice at this site is heterogeneous: autochthonous and allochthonous.

A massive-ice complex is often well exposed by coastal erosion at North Head, on Richards Island (69°20'N, 134°30'W), in the Tuktoyaktuk Coastlands. According to Murton (2005), some of the massive ice is buried and some is intrasedimental. Y. K. Vasil'chuk agreed and identified the ice as heterogeneous: autochthonous and allochthonous.

Massive ground-ice bodies in the Sandhills Moraine of southern Banks Island, and the southern Eskimo Lakes region of the Tuktoyaktuk Coastlands, according to French and Harry (1990), are interpreted as basal glacier ice. Other massive ground-ice bodies in the Western Canadian Arctic, however, are explained better in terms of segregation-injection (French and Harry, 1990). Assessing the isotopic variations in ice around the Eskimo Lakes, Y. K. Vasil'chuk noted that the bottom of the massive ice described by French and Harry (1990) is characterized by a relatively stable distribution and a small range (about 1.5 ‰) of  $\delta^{18}\text{O}$  values (from -33.5 to -35 ‰) and by abrupt changes in the upper 2 m of ice: upward, the value increases to -31 ‰, and then sharply decreases to -36‰. Y. K. Vasil'chuk suggested that the isotopic changes in the upper 2 m of ice are consistent with ice

segregation commencing in a semi-closed or open system, and ending in a closed system, leading to an appreciable isotope differentiation in the top of the ice deposits.

Massive ice also occurs within glacially deformed permafrost in the coastal lowlands near Tuktoyaktuk (e.g. at Liverpool Bay - 70°N, 129°W, southwest of Nicholson Island; and on northern 'Crane Island', in the central Eskimo Lakes; Murton *et al.*, 2004). The massive ice is at least 2.5–8 m thick, and either white and bubble-rich, or grey, debris-rich and banded. Two types of ground ice are identified in the deformed permafrost: (1) massive ice and ice clasts, both of which have been glacially deformed, eroded or moved; and (2) segregated ice and ice-wedge ice that have not been glacially disturbed because their postdate deformation. Y. K. Vasilchuk agreed with these interpretations and determined the massive ice as heterogeneous: autochthonous and allochthonous.

Within the limits of the Willow River drainage basin, on the Aklavik Plateau (69°N, 124°W), Lacelle *et al.* (2004) examined four exposures of debris-rich massive ice. The  $\delta^{18}\text{O}$  values of the ice change abruptly from an average of about  $-30\text{‰}$  (for Late Pleistocene ice) to  $-22.6\text{‰}$  (for Holocene ice). Physical and isotopic properties of the former suggest that it is segregated-intrusive ice, because the  $\text{CO}_2$  content and  $\delta^{13}\text{C}$  value in the debris-rich ice was acquired during movement of  $\text{CO}_2$  through the sediments, giving  $\text{CO}_2$  concentrations that are 3–9 times higher than those for air trapped in glacier ice and  $\delta^{13}\text{C}$  values in the range of  $\text{CO}_2$  produced by the decay of  $\text{C}_3$  plant material (Lacelle *et al.*, 2004). Y. K. Vasilchuk agreed with these interpretations and determined the massive ice as heterogeneous: autochthonous and allochthonous.

Massive ice within but close to the glacial limit of the Laurentide Ice Sheet, on Herschel Island (69°N, 139°W), in the southern Beaufort Sea, is highly depleted in heavy isotopes (mean  $\delta^{18}\text{O}$  value:  $-33\text{‰}$ ;  $\delta\text{D}$ :  $-258\text{‰}$ ; Fritz *et al.*, 2011, 2012). These authors noted that such stable isotope signatures indicate a full-glacial water source for the massive ice on Herschel Island. However, an origin as glacially deformed segregated or segregated-intrusive ice cannot be excluded. Pollard (1990) concluded that segregated ice is the most common massive-ice type in this area and in places constitutes up to 70% of the upper 10–15 m of permafrost. Y. K. Vasil'chuk suggested that the origin of massive ice on Herschel Island varies, and the ice includes both heterogeneous and homogeneous types.

## **CHAPTER 6. DATING OF MASSIVE ICE OF RUSSIA AND NORTH AMERICA, AND CORRELATION OF STABLE ISOTOPE CURVES**

Chapter 6 focuses on radiocarbon dating of organic material in the sediments that *surround* massive ice in Russia and Canada, and on direct accelerator mass spectrometry (AMS) radiocarbon dating of

organic microinclusions and trapped gases *within* massive ice in Canada. The radiocarbon ages suggest that most of the massive ice accumulated between the Holocene and 20–40 ka BP (Figure 3). Comparison of isotopic plots of massive ice in Russia and Canada (Figures 4–6) shows that they have more similarities than differences in the regional isotopic record of ice formation. An initial isotopic signature indicates the nature of the water and the conditions of ice formation. Ice segregation in a closed system leads to a contrasting distribution of  $\delta^{18}\text{O}$  and  $\delta\text{D}$  values, both vertically and laterally (Figure 4). The shape of isotopic plots of massive ice relates to the type of ice formation. Homogeneous, undifferentiated isotope plots with a narrow range of isotopic changes ( $\delta^{18}\text{O}$  range  $<4\text{‰}$ ,  $\Delta\delta\text{D}$  range  $<32\text{‰}$ ) characterize massive ice formed in an open system, with freezing of inflowing water under homogeneous conditions. By contrast, isotope plots with a wide range of isotopic changes ( $\Delta\delta^{18}\text{O}$  range  $> 8\text{‰}$ ,  $\Delta\delta\text{D}$  range  $> 64\text{‰}$ ) characterize massive ice formed in a closed system, where there is no inflowing water. Closed-system freezing is typical of segregation ice formation, or rarely for the final phase of injection ice formation when there is no inflow of water (Figs. 5 and 6).

## CONCLUSIONS

- The monograph considers practically all the selected ages and mechanisms of massive-ice formation. The senior author proposed the original mechanism of the formation of massive ice in saline lake sediments. New mechanisms will undoubtedly be proposed in the future.
- A new genetic classification of massive-ice deposits introduces two new categories at its highest level: *homogeneous* and *heterogeneous* massive-ice deposits.
- Assemblages of different massive-ice types are common in permafrost exposures in the Yamal Peninsula of west Siberia and the Tuktoyaktuk Coastlands and Yukon of northwest Canada.
- Plotting of isotopic data from massive ice on graphs with a single vertical and horizontal size facilitates objective assessment of the isotopic characteristics of massive ice.

## Acknowledgments

Researches of Yu.Vasil'chuk are financially supported by Russian Scientific Foundation (grant RSF № 14-27-00083).

## REFERENCES

- Astakhov VI, Isayeva LL. 1988. The “Ice Hill”: an example of ‘redarded deglaciation’ in Siberia. *Quaternary Science Reviews* 7: 29-40. DOI:10.1016/0277-3791(88)90091-1.
- Belova NG. 2014. *Massive ice beds of south-western coast of the Kara Sea*. MAKS Press: Moscow. (in Russian).
- Cardyn R, Clark ID, Lacelle D, Lauriol B, Zdanowicz C, Calmels F. 2007. Molar gas ratios of air entrapped in ice: A new tool to determine the origin of relict massive ground ice bodies in permafrost. *Quaternary Research* 68: 239–248. DOI: 10.1016/j.yqres.2007.05.003.
- Dallimore SR, Wolfe SA. 1988. Massive ground ice associated with glacial sediments, Richards Island, N.W.T., Canada. In *Proceedings of the Fifth International Conference on Permafrost*, Senneset K. (ed.). Tapir Publishers: Trondheim, Norway; 132 – 138.
- Danilov ID. 1989. Massive ice and subaqueous cryolithogenesis. In *Geocryological Study*, Yershov ED (ed). Moscow University Press; Moscow; 16–29. (in Russian).
- Dubikov GI. 2002. *Composition and Cryogenic Structure of the Permafrost Strata of Western Siberia*. Geos Publishing House: Moscow. (in Russian).
- French HM. 2007. *The Periglacial Environment*. Third Edition. John Wiley and Sons Ltd: Chichester.
- French HM, Harry DG. 1990. Observations on buried glacier ice and massive segregated ice, western arctic coast, Canada. *Permafrost and Periglacial Processes* 1: 31–43. DOI: 10.1002/ppp.3430010105.
- Fritz M, Wetterich S, Meyer H, Schirmermeister L, Lantuit H, Pollard WH. 2011. Origin and characteristics of massive ground ice on Herschel Island (Western Canadian Arctic) as revealed by stable water Isotope and hydrochemical signatures. *Permafrost and Periglacial Processes* 22: 26–38. DOI: 10.1002/ppp.714.
- Fritz M, Wetterich S, Schirmermeister L, Meyer H, Lantuit H, Preusser F, Pollard WH. 2012. Eastern Beringia and beyond: Late Wisconsinan and Holocene landscape dynamics along the Yukon Coastal Plain, Canada. *Palaeogeography, Palaeoclimatology, Palaeoecology* 319–320: 28–45. doi:10.1016/j.palaeo.2011.12.015.
- Fujino K, Horiguchi K, Shinbori M, Kato K. 1983. Analysis and characteristics of cores from a massive ice body in Mackenzie delta, N.W.T., Canada. In *Proceedings of the Fourth International Conference on Permafrost*. National Academy Press: Washington, D.C.; 316 – 321.
- Fujino K, Sato S, Matsuda K, Sasa G, Shimizu O, Kato K. 1988. Characteristics of the massive ground ice body in the western Canadian Arctic. In *Proceedings of the Fifth International Conference on Permafrost*, Senneset K. (ed). Tapir Publishers: Trondheim, Norway; 143–146.

- Gasanov SS. 1969. *The Structure and the Formation History of the Frozen Grounds in Eastern Chukotka*. Nauka: Moscow. (in Russian).
- Holmsen G. 1914. Spitsbergens jordbunds is og de bidrag dens undersøkelse har kunnet gi til forstaaelsen a de i arktiske land optrædende varige isleier i jorden. (Ground ice in Spitsbergen and contributions toward the understanding of perennial ground ice found in Arctic environments). *Det Norske Geografiske Selskabs Årbok* **B XXIV**, 1912–1913, Kristiania: 1–150.
- Hurlbert SH, Chang CCY. 1984. Ancient Ice Islands in Salt Lakes of the Central Andes. *Science* **224**: 299–302. DOI:10.1126/science.224.4646.299.
- Hyatt JA. 1998. The origin of lake-bed ground ice at Water Supply Lake, Pond Inlet, Nunavut, Canada. In *Proceedings of the 7th International Conference on Permafrost*, Yellowknife, Canada, Lewkowicz AG, Allard M (eds.). Universite Laval, Canada. Collection Nordicana **57**; 757–762.
- Ingólfsson Ó, Lokrantz H. 2003. Massive ground ice body of glacial origin at Yugorski Peninsula, Arctic Russia. *Permafrost and Periglacial Processes* **14**: 199 – 215. DOI: 10.1002/ppp.455.
- Ivanova VV. 2012. Geochemical features of formation of massive ground ice bodies (New Siberia Island, Siberian Arctic). *Earth Cryosphere* **1**: 56–70 (in Russian).
- Kato K., Fujino K. 1981. Oxygen isotopic composition of massive ice at Tuktoyaktuk, North Canada. In *Joint Studies on physical and biological environments in the permafrost, North Canada, July to August 1980, and February to March, 1981*, S.Kinosita S. (ed). Institute of Low Temperature Science: Hokkaido University, Sapporo; 13–20.
- Kato K, Sato S, Fujino K. 1988. Radiocarbon dating by accelerator mass spectrometry on sediment in a core from a massive ice body in Mackenzie Delta, N.W.T., Canada: preliminary report. In *Characteristics of the massive ground ice body in the western Canadian Arctic related to paleoclimatology 1986–1987*, Fujino K (ed). Institute of Low Temperature Science: Hokkaido University, Sapporo; 58–69.
- Korolev SY. 1993. The discovery of the Late Pleistocene glacier ice in the valley of the Amguema River (Northern Chukotka). *Doklady of the Russian Academy of Sciences*. **329**: 195–198 (in Russian).
- Kotov AN. 1998. The cryolithogenic ridges in the valley of the Tanurer River (Chukotka). *Earth Cryosphere* **4**: 62–71 (in Russian).
- Kritsuk LN. 2010. *Ground Ice of Western Siberia*. Nauchny Mir Publishing House: Moscow. (in Russian).

- Lacelle D. 2011. On the  $\delta^{18}\text{O}$ ,  $\delta\text{D}$  and d-excess relations in meteoric precipitation and during equilibrium freezing: theoretical approach and field examples. *Permafrost and Periglacial Processes* **22**: 13–25. DOI: 10.1002/ppp.712.
- Lacelle D, Bjornson J, Lauriol B, Clark ID, Troutet Y. 2004. Segregated-intrusive ice of subglacial meltwater origin in retrogressive thaw flow headwalls, Richardson Mountains, N.W.T., Canada. *Quaternary Science Reviews* **23**: 681 – 696. DOI:10.1016/j.quascirev.2003.09.005.
- Lacelle D, Lauriol B, Clark ID, Cardyn R, Zdanowicz CH. 2007. Nature and origin of a Pleistocene-age massive ground-ice body exposed in the Chapman Lake moraine complex, central Yukon Territory. *Quaternary Research* **68**: 249–260. DOI:10.1016/j.yqres.2007.05.002.
- Lacelle D, St-Jean M, Lauriol B, Clark ID, Lewkowicz A, Froese DG, Kuehn SC, Zazula G. 2009. Burial and preservation of a 30,000 year old perennial snowbank in Red Creek valley, Ogilvie Mountains, central Yukon, Canada. *Quaternary Science Reviews* **28**: 3401–3413. DOI:10.1016/j.quascirev.2009.09.013
- Lacelle D, Davila AF, Pollard WH, Andersen D, Heldmann J, Marinova M, McKay CP. 2011. Stability of massive ground ice bodies in University Valley, McMurdo Dry Valleys of Antarctica: Using stable O–H isotope as tracers of sublimation in hyper-arid regions. *Earth and Planetary Science Letters* **301**: 403–411.
- Leibman MO, Hubberten H.-W, Lein AY, Streletskaya ID, Vanshtein BG. 2003. Tabular ground ice origin: cryolithological and isotope-geochemical study. In *Proceedings of the 8th International Conference on Permafrost*, M.Phillips M, Springman SM, Arenson L. (eds). A.A. Balkema Publishers: Lisse, Netherlands; 645–650.
- Lorrain RD, Demeur P. 1985. Isotopic evidence for relic Pleistocene glacier ice on Victoria Island, Canadian Arctic Archipelago. *Arctic and Alpine Research* **17**: 89–98. DOI: 10.2307/1550964.
- Mackay JR. 1971. The Origin of Massive Icy Beds in Permafrost, Western Arctic Coast, Canada. *Canadian Journal of Earth Sciences* **8/4**: 397–422. DOI:10.1139/e71-043.
- Mackay JR. 1973. Problems in the origin of massive ice beds, western Arctic Canada. In *Proceedings of the 2nd International Conference on Permafrost, Yakutsk, USSR, 13–28 July 1973*, Péwé TL, Mackay JR (eds). National Academy of Sciences: Washington D.C., North American Contributions; 223–238.
- Mackay JR. 1983. Oxygen isotope variations in permafrost, Tuktoyaktuk Peninsula area, Northwest Territories. In *Current Research, Part B*. Geological Survey of Canada, Paper **83–1B**: 67–74.
- Mackay JR, Dallimore SR. 1992. Massive ice of the Tuktoyaktuk area, western Arctic coast, Canada. *Canadian Journal of Earth Sciences* **29**: 1235–1249. DOI:10.1139/e92-099.

- Melnikov VP, Spesivtsev VI. 2000. Cryogenic formations in the earth's lithosphere (graphic version). Konishchev VN (ed.). Novosibirsk: Scientific Publishing Center of the UIGGM, Siberian Branch of the RAS, Publishing House, Siberian Branch of the RAS, 2000.
- Michel FA. 1983. Isotope variations in permafrost waters along the Dempster Highway pipeline corridor In *Proceedings of the 4th International Conference on Permafrost*. Fairbanks. Alaska. National Academy Press: Washington; 843–848.
- Michel FA. 1998. The relationship of massive ground ice and the Late Pleistocene history of Northwest Siberia. *Quaternary International* **45/46**: 43–48. DOI:10.1016/S1040-6182(97)00005-0.
- Michel FA. 2011. Isotope characterisation of ground ice in northern Canada. *Permafrost and Periglacial Processes* **22**: 3–12. DOI: 10.1002/ppp.721.
- Moorman BJ, Michel FA, Wilson A. 1996. <sup>14</sup>C dating of trapped gases in massive ground ice, Western Canadian Arctic. *Permafrost and Periglacial Processes* **7**: 257–266. DOI: 10.1002/(SICI)1099-1530(199609)7:3<257::AID-PPP220>3.0.CO;2-P.
- Moorman BJ, Michel FA, Wilson AT. 1998. The development of tabular massive ground ice at Peninsula Point, N.W.T., Canada. In *Proceedings of the 7th International Conference on Permafrost*, Lewkowicz AG, Allard M. (eds). Universite Laval, Canada. Collection Nordicana **57**; 757–762.
- Murton JB. 2005. Ground-ice stratigraphy and formation at North Head, Tuktoyaktuk Coastlands, western Arctic Canada: a product of glacier–permafrost interactions. *Permafrost and Periglacial Processes* **16**: 31–50. DOI: 10.1002/ppp.513.
- Murton JB. 2009. Stratigraphy and palaeoenvironments of Richards Island and the eastern Beaufort Continental Shelf during the last glacial–interglacial cycle. *Permafrost and Periglacial Processes* **20**: 107–125. DOI: 10.1002/ppp.647.
- Murton JB, Waller RI, Hart JK, Whiteman CA, Pollard WH, Clark ID. 2004. Stratigraphy and glaciotectonic structures of permafrost deformed beneath the northwest margin of the Laurentide Ice Sheet, Tuktoyaktuk Coastlands, Canada. *Journal of Glaciology* **49**: 399–412. DOI: 10.3189/172756504781829927.
- Murton JB, Whiteman CA, Waller RI, Pollard WH, Clark ID, Dallimore SR. 2005. Basal ice facies and supraglacial melt–out till of the Laurentide Ice Sheet, Tuktoyaktuk Coastlands, western Arctic Canada. *Quaternary Science Reviews* **24**: 681–708. DOI:10.1016/j.quascirev.2004.06.008.
- Pollard W. 1990. The nature and origin of ground ice in the Herschel Island area, Yukon Territory. In *Permafrost – Canada. Proceedings of the Fifth Canadian Permafrost Conference*. Collection

- Nordicana, N54. Centre d'etudes Nordiques, Universite Laval. Quebec: National Research Council of Canada; 23–30.
- Pollard W, Bell T. 1998. Massive ice formation in the Eureka Sound Lowlands: A landscape model. In *Proceedings of the 7th International Conference on Permafrost*, Yellowknife. Canada, Lewkowicz AG, Allard M (eds). Universite Laval, Canada. Collection Nordicana **57**; 903–908.
- Ponomarev VM. 1960. *Underground waters formation in the permafrost along the coast of the northern seas*. Moscow: Publishing House of the USSR Academy of Sciences. 1960. (in Russian).
- Rampton VN. 1988. Origin of massive ground ice on Tuktoyaktuk Peninsula, Northwest Territories, Canada. In *Proceedings of the Fifth International Conference on Permafrost*, Senneset K. (ed). Tapir Publishers: Trondheim, Norway; 850–855.
- Robinson S, Pollard W. 1998. Massive ground ice within Eureka Sound bedrock, Fosheim Peninsula, Ellesmere Island, N.W.T. In *Proceedings of the 7th International Conference on Permafrost*, Yellowknife. Canada, Lewkowicz AG, Allard M (eds). Universite Laval, Canada. Collection Nordicana **57**; 949–954.
- Slagoda EA, Opokina OL, Kurchatova AN, Rogov VV. 2012. Structure and composition of complex massive ice bodies in Late Pleistocene-Holocene sediments of the Marre-Sale Cape, West Yamal. In *Proceedings of the Tenth International Conference on Permafrost*, Melnikov VP, Drozdov DD, Romanovsky VE (eds). Salekhard, 25–29 June 2012. Vol. 2. The Northern Publisher: Salekhard, Russia; 415–420.
- Shpolyanskaya NA. 2014. Permafrost (Cryolithozone) of the Russian Arctic Shelf. *Universal Journal of Geoscience* **2**: 7–17. DOI: 10.13189/ujg.2014.020102.
- Solomatina VI. 2013. *Physics and Geography of Ground Glaciation*. Academic Publishing House "Geo": Novosibirsk. 246 pp. (in Russian).
- Streletskaya ID, Gusev EA, Vasiliev AA, Oblogov GE, Molodkov AN. 2013. Pleistocene - Holocene paleoenvironmental records from permafrost sequences at the Kara Sea coasts (NW Siberia, Russia). *Geography, Environment, Sustainability* **3**: 60–76.
- Vaikmäe RA, Karpov YG. 1986. Study of tabular ground ice bodies from the Ledyanaya Gora section in the Yenisey valley using the Oxygen isotope method. *Polar Geography and Geology* **10**: 32–38. DOI:10.1080/10889378609377270.
- Vaikmäe RA, Vasil'chuk YK. 1991. *Oxygen isotope analyses of ground ice north-west Siberia, Yakutia and Chukotka*. Tallinn. Geology Institute of the Estonia Academy of Science. (in Russian).

- Vaikmäe R, Michel, FA, Solomatin VI. 1993. Morphology, stratigraphy and oxygen isotope composition of fossil glacier ice at Ledyanaya Gora, Northwest Siberia, Russia. *Boreas* **22**: 205–213. DOI: 10.1111/j.1502-3885.1993.tb00180.x.
- Vasil'chuk AC, Vasil'chuk YK. 2010. Local pollen spectra as a new criterion for nonglacial origin of massive ice. *Transactions (Doclady) of Russian Academy of Sciences, Earth Sciences* **433**: 985–990. DOI: 10.1134/S1028334X10070305.
- Vasil'chuk AC, Vasil'chuk YK. 2012. Pollen and Spores as Massive Ice Origin Indicators. In *Proceedings of the Tenth International Conference on Permafrost*, Melnikov VP, Drozdov DD, Romanovsky VE (eds). Salekhard, 25–29 June 2012. Vol. 2. The Northern Publisher: Salekhard, Russia; 487 – 491.
- Vasil'chuk YK. 2011. Homogeneous and Heterogeneous Massive Ice in Permafrost. *Earth Cryosphere* **1**: 40–51 (in Russian).
- Vasil'chuk YK. 1992. Oxygen Isotope composition of Ground Ice (Application to Paleogeocryological Reconstruction). Department of theoretical problems the Russian Academy of Sciences. Moscow State University, Industrial and Research Institute of Engineering Survey for Construction. In 2 volumes. Vols. 1 and 2. (in Russian).
- Vasil'chuk YK. 2012. Classification of Massive Ice. In *Proceedings of the Tenth International Conference on Permafrost*, Melnikov VP, Drozdov DD, Romanovsky VE (eds.). Salekhard, 25–29 June 2012. Vol. 2. Salekhard: The Northern Publisher, Russia; 493–497.
- Vasil'chuk YK. 2012. *Stable Isotope Geochemistry of Massive Ice. Vol. I*. Moscow University Press: Moscow (in Russian).
- Vasil'chuk YK. 2013. Syngenetic ice wedges: cyclical formation, radiocarbon age and stable-isotope records. *Permafrost and Periglacial Processes* **24**: 82–93. DOI: 10.1002/ppp.1764.
- Vasil'chuk YK. 2014. *Stable Isotope Geochemistry of Massive Ice. Vol. II*. Moscow University Press: Moscow (in Russian).
- Vasil'chuk YK, Trofimov VT. 1988. Oxygen isotope variations in ice-wedge and massive ice. In *Proceedings of the Fifth International Conference on Permafrost*, Senneset K. (ed.). Trondheim. Norway. Tapir Publishers; 489 – 492.
- Vasil'chuk YK, Vasil'chuk AC, Budantseva NA, Chizhova JN, Papesch W, Podborny YY, Sulerzhitsky LD. 2009. Oxygen Isotope and Deuterium Indication of the Origin and <sup>14</sup>C Age of the Massive Ice, Bovanenkovo, Central Yamal Peninsula. *Doklady Earth Sciences* **429**: 1326–1332. DOI: 10.1134/S1028334X09080194.

- Vasil'chuk YK, Budantseva NA., Vasil'chuk AC. 2011. Variations in  $\delta^{18}\text{O}$ ,  $\delta\text{D}$ , and the concentration of pollen and spores in autochthonic heterogeneous massive ice on the Erkutayaha River in the southern part of the Yamal Peninsula. *Doklady Earth Sciences* **438**: 721–726.  
DOI: 10.1134/S1028334X11050382.
- Vasil'chuk YK, Vasil'chuk AC, Budantseva NA. 2012. Isotopic and palynological compositions of a massive ice in the Mordyyakha River, Central Yamal Peninsula. *Doklady Earth Sciences* **446**: 1105–1109. DOI: 0.1134/S1028334X12090164.
- Vasil'chuk YK, Vasil'chuk AC, Budantseva NA. 2012. Isotopic and palynological compositions of a massive ice in the Mordyyakha River, Central Yamal Peninsula. *Doklady Earth Sciences* **446**: 1105–1109. DOI: 0.1134/S1028334X12090164.
- Vasil'chuk Y, Vasil'chuk A. 2014. Spatial distribution of mean winter air temperatures in Siberian permafrost at 20–18 ka BP using oxygen isotope data. *Boreas* **43**: 43–52. DOI: 10.1111/bor.12033.
- Vasil'chuk YK, Vasil'chuk AC, Budantseva NA, Chizhova JN, Papesch W, Podborny YY. 2014.  $^{14}\text{C}$  age, stable isotope composition and pollen analysis of massive ice, Bovanenkovo gas field, Central Yamal Peninsula. *Geography, Environment, Sustainability*. **2**: 49–70.
- Vasil'chuk YK, Budantseva NA, Vasil'chuk AC, Yoshikawa K, Podborny YY, Chizhova JN. 2014. Isotope composition of pingo ice core at Evoyakha River valley, north-west Siberia. *Earth Cryosphere* **4**: 47–58 (in Russian).
- Vasil'chuk YK, Budantseva NA, Vasil'chuk AC, Podborny YY, Sullina AN, Chizhova JN. 2015. Isotope composition of the Holocene heterogeneous massive ice, the north-eastern part of the Yamal Peninsula. *Earth Cryosphere* **3**: 35–54 (in Russian).
- Vtyurin BI. 1975. *Ground Ice of the USSR*. Nauka: Moscow (in Russian).
- Vtyurin BI, Glazovskiy AF. 1986. Composition and structure of the Ledyanaya Gora tabular ground ice body on the Yenisey. *Polar Geography and Geology* **10**: 273–285.  
DOI:10.1080/10889378609377297.
- Wolfe SA. 1998. Massive ice associated with glaciolacustrine delta sediments Slave geological province, N.W.T., Canada // *Permafrost. Seventh International Conference. Proceedings*. Yellowknife. Canada. Lewkowicz AG, Allard M (eds.). Universite Laval, Collection Nordicana, N57. Canada. 1133 – 1139.
- Zhestkova TN, Shur YL. 1978. On the genesis of tabular massive ice. *The Bulletin of the Moscow State University. Geology Series* **3**: 35–42 (in Russian).