

Disrupted neural activity patterns to novelty and effort in young adult APOE-e4 carriers performing a subsequent memory task

Article (Accepted Version)

Evans, Simon, Dowell, Nicolas G, Tabet, Naji, King, Sarah L, Hutton, Samuel B and Rusted, Jennifer M (2017) Disrupted neural activity patterns to novelty and effort in young adult APOE-e4 carriers performing a subsequent memory task. *Brain and Behavior*, 7 (2). e00612. ISSN 2162-3279

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/66194/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Disrupted neural activity patterns to novelty and effort in young adult *APOE*-e4 carriers performing a subsequent memory task

Simon Evans^{1,2}, Nicholas G Dowell³, Naji Tabet³, Sarah L King¹, Samuel B Hutton¹, Jennifer M Rusted^{1,4}

¹School of Psychology, University of Sussex, Brighton, East Sussex BN1 9QG, United Kingdom

²School of Psychology, University of Surrey, Guildford, Surrey GU2 7XH, United Kingdom

³Brighton and Sussex Medical School (BSMS), Brighton, East Sussex BN1 9RR, United Kingdom

⁴Corresponding author: School of Psychology, University of Sussex, Brighton, East Sussex BN1 9QG, United Kingdom. Tel: (+44) 1273 878325

Short title: Subsequent memory in young adult *APOE* e4 carriers

Keywords: *APOE*; Memory; fMRI; Reflex, Pupillary; Hippocampus

Abstract

Introduction: The APOE e4 allele has been linked to poorer cognitive ageing and enhanced dementia risk. Previous imaging studies have used subsequent memory paradigms to probe hippocampal function in e4 carriers across the age range, and evidence suggests a pattern of hippocampal overactivation in young adult e4 carriers.

Methods: In this study, we employed a word-based subsequent memory task, under fMRI; pupillometry data was also acquired as an index of cognitive effort.

Participants (26 non-e4 carriers and 28 e4 carriers) performed an incidental encoding task (presented as word categorization) followed by a surprise old/new recognition task after a 40 minute delay.

Results: In e4 carriers only, subsequently remembered words were linked to increased hippocampal activity. Across all participants, increased pupil diameter differentiated subsequently remembered from forgotten words, and neural activity covaried with pupil diameter in cuneus and precuneus.

These effects were weaker in e4 carriers, and e4 carriers did not show greater pupil diameter to remembered words. In the recognition phase, genotype status also modulated hippocampal activity: here, however, e4 carriers failed to show the conventional pattern of greater hippocampal activity to novel words.

Conclusions:

Overall, neural activity changes were unstable in e4 carriers, failed to respond to novelty, and did not link strongly to cognitive effort, as indexed by pupil diameter.

This provides further evidence of abnormal hippocampal recruitment in young adult e4 carriers, manifesting as both up- and down-regulation of neural activity, in the absence of behavioural performance differences.

Introduction

In humans, 3 variants of the *APOE* gene exist (e2, e3, e4). The e4 allelic variant has been the focus of considerable recent research activity due to it being a well-established risk factor for Alzheimer's disease (AD) (Rocchi, et al. 2003). It also impacts healthy ageing: carriers of the e4 variant (from this point referred to as e4+) have been shown (in the absence of AD) to be cognitively disadvantaged in later life relative to non-e4 carriers (e4-) on measures of episodic memory, executive functioning and overall global cognitive ability (Wisdom, et al. 2011), and longitudinal studies suggest that healthy age-related cognitive decline begins earlier in e4+ and progresses quicker (Caselli, et al. 2009; Davies, et al. 2012). These effects occur in the context of brain structural differences. Healthy older e4+ show grey matter (GM) reductions in hippocampal and frontotemporal regions (Wishart, et al. 2006); this is noteworthy since these regions are amongst the first to atrophy in AD (Thompson, et al. 2003). Neural activation differences are also evident, with greater BOLD activity observed in various regions including precuneus, frontal and right hippocampal regions during picture encoding in healthy e4+ aged 70-80 (Bondi, et al. 2005). Retrieval of memorised word pairs has also been shown to induce greater activity in parietal, and prefrontal and hippocampal regions in e4+ (aged 47-82), with degree of overactivity correlating with degree of memory decline measured 2 years later (Bookheimer, et al. 2000). Overactivity has also been demonstrated during working memory tasks, with e4+ aged 50-75 showing greater recruitment of medial frontal and parahippocampal areas (Filbey, et al. 2010). Another study has reported increased activity in prefrontal, temporal and parietal regions during memory encoding, but coupled with frontal decreases during retrieval, in e4+ aged 55-65 (Kukolja, et al. 2010). These findings have been interpreted as representing

compensatory mechanisms: e4+ recruit additional neural resources to maintain cognitive performance (Tuminello and Han 2011), thus requiring additional cognitive effort to achieve comparable performance levels to their non-e4 peers (Bondi, et al. 2005)

There is some evidence that e4+ might show neural differences even in young adulthood. Most work has focused on hippocampal activity patterns to try and characterize differences that might anticipate later-life pathology, and various studies point to a pattern of hippocampal overactivity in e4+. Dennis et al. (2009) employed a subsequent memory task: this paradigm begins with an acquisition phase containing a set of stimuli to be remembered, followed after some fixed interval by a recognition phase where those same stimuli are presented again, interleaved with some novel stimuli, and participants respond to indicate whether they think each item was previously studied or novel. Dennis et al. employed pictorial stimuli and a 24-hour retention period, and investigated activation in the medial temporal lobe during the acquisition phase, comparing activity to items that were subsequently remembered and items that were subsequently forgotten. In adults aged 20-25, hippocampal activity in e4- did not differentiate remembered from forgotten, but significantly greater bilateral hippocampal recruitment to subsequently remembered items was seen in e4+. Task performance was equal across genotypes. Similarly, a study by Filippini et al. (2009) used a variant of the subsequent memory paradigm, again using pictorial stimuli but focussing on the recognition phase, comparing effects of novel versus familiar stimuli. It was found that young adult e4+ (mean age 28) showed a pattern of hippocampal over-recruitment to novel stimuli when presented amongst well-learned 'familiar' stimuli. This was replicated in a follow-up study in a slightly older age

range (32-55), which also reported hippocampal overactivity during a Stroop task, where hippocampal activation was not to be expected (Trachtenberg, et al. 2012a). Similarly, we have also reported hippocampal recruitment in e4+ (aged 18-28) during a covert attention task which does not usually elicit such activity (Rusted, et al. 2013). It has been argued that such neural overrecruitment, seemingly evident across the lifespan in e4+ and possibly compensatory in nature, could drive cognitive performance advantages in young adulthood (Tuminello and Han 2011). Some studies have reported that young adult e4+ can manifest cognitive advantages in certain domains, with e4+ outperforming e4- on measures of verbal fluency and prospective memory (Marchant, et al. 2010), and sustained and covert attention (Rusted, et al. 2013), but larger studies using more general cognitive test batteries report no evidence for advantages (Bunce, et al. 2011). Further work is required to resolve this issue, and interpret the significance of hippocampal overactivity in young adult e4+. Some MRI studies in young adult e4+ point to reduced volume in medial temporal lobe (MTL) (O'Dwyer, et al. 2012; Wishart, et al. 2006), and resting state studies have shown enhanced co-activation within hippocampal (Trachtenberg, et al. 2012b) and default mode (Filippini, et al. 2009; Su, et al. 2015) networks, supporting a compensatory recruitment hypothesis.

Not all data is consistent with this, however. Mondadori et al., using an associative learning task, found that e4+ aged 20-25 actually showed diminishing hippocampal recruitment as the task progressed and this was linked to better performance. In contrast, e4- showed activity increases, leading the authors to suggest that e4+ might actually under-recruit neural resources under certain circumstances (Mondadori, et al.

2007) and thus be more efficient in terms of neural recruitment. In young adulthood, therefore, a straightforward compensatory model might be overly simplistic.

In the current study, we reverted to a classic subsequent memory paradigm, and extending the work outlined above, imaged both the acquisition and recognition phases so as to fully characterise hippocampal activation patterns in young adult e4+ during the task. Pupillometry data was acquired during the acquisition phase as an index of cognitive effort. Since compensatory neural recruitment likely reflects increased cognitive effort in older e4+ (Bondi, et al. 2005) measuring cognitive effort could provide insight into whether differences in neural recruitment serve a similar compensatory role in younger e4+. Word stimuli were employed, to minimise luminance changes and eye movements. Evidence that pupil diameter can serve as an index of cognitive effort has been demonstrated across a variety of cognitive domains: for example, pupil size increases with task complexity during sentence comprehension (Just and Carpenter 1993), and pitch discrimination (Schlemmer, et al. 2005). Pupil diameter has been shown to correlate with neural activity in dorsal attentional networks during a divided attention task (Alnaes, et al. 2014), suggesting that pupil diameter could indicate the level of cognitive resources being directed towards a stimulus. In subsequent memory tasks, pupil diameter is enlarged to words that are subsequently remembered, versus forgotten (Papesh, et al. 2012). If neural recruitment differences reflect enhanced cognitive effort being deployed in e4+ as a means of achieving the same level of cognitive performance as in e4-, this should be detectable in the pupillometry measures. As such, we predicted genotype-specific effects in pupil diameter (specifically, greater pupil diameter in e4+), and these effects were tested in two ways. Firstly we examined average pupil diameter in each

condition (remembered/forgotten), by genotype. We then included pupil diameter as a covariate in the fMRI analyses to link pupillometry and neural activity measures. We did not anticipate any genotype differences in memory performance: a recent study using a word-based subsequent memory task found that APOE status did not affect performance in young adults (Stening, et al. 2016), as did the majority of studies using pictorial stimuli (outlined above), although it should be noted that these imaging studies have typically employed relatively small numbers and therefore might not have sufficient power to detect subtle memory impairment.

In terms of neural activation patterns, we predicted genotype-specific differences in hippocampal activation, and a small volume correction was employed using a mask that incorporated both hippocampal and parahippocampal regions, bilaterally. This was used to determine whether levels of hippocampal activity showed any interactions between genotype and task condition, and specifically to test whether e4+ show greater hippocampal activity to trials that are subsequently remembered relative to those subsequently forgotten (as demonstrated by Dennis et al. (2009)).

Materials and methods

Participants 328 healthy participants (aged 18-28 years) were recruited from the University of Sussex. Protocols specified by the Human Tissue Act were followed throughout, participants consented to not being informed of their genotyping result, and volunteer call-back was performed by a third party so that the researcher remained blind. *APOE* genotype was determined by buccal swab. Genotype analyses were performed by a third party (LGC Genomics, Hoddesdon, UK) using fluorescence-based competitive allele-specific polymerase chain reaction (KASPar)

targeting two APOE single-nucleotide polymorphisms (SNPs): rs429358 and rs7412. Invitation to the study was based on a random sampling so genotype status could not be inferred from an invitation to take part. Of these 328, 61 volunteers carried at least one e2 allele and were excluded. 69 volunteers carried at least one e4 allele: 40 of these individuals were randomly invited to the study, of which 28 consented to take part. 197 volunteers were homozygous e3 carriers and of these 50 were also randomly invited to the study, of which 26 consented to take part. Amongst the e4+ group, 6 participants were homozygous e4 carriers. Inclusion criteria were: age 18-28, right handed, and fluent English speaker. Participants were excluded if they reported having high blood pressure, current treatment for a psychiatric condition, or failed the MRI safety screening.

The two groups were matched in age, but there was a trend towards an unequal gender balance, with more females than males overall (1 tailed proportion test, $z = 1.631$, $p=0.052$). For participants included in the fMRI analyses (whose recognition performance exceeded 50%), there was no significant difference in gender balance (1 tailed proportion test, $z = 0.316$, $p=0.376$), see Table 1. Nevertheless, gender was entered as a covariate in the behavioural, imaging and pupillometry analyses.

All participants			Participants included in fMRI analyses		
Group	Age (years)	Gender	Group	Age (years)	Gender
e4- (n=26)	20.91 ± 1.90	14F/12M	e4- (n=19)	20.93 ± 2.14	8F/11M
e4+ (n=28)	20.92 ± 2.59	19F/9M	e4+ (n=21)	21.04 ± 2.77	13F/8M
t-statistic	0.444, ns		t-statistic	0.481, ns	

Table 1 Volunteer characteristics for all participants, and those included in the fMRI analyses (recognition performance >50%).

Experimental design All participants volunteered under a written informed consent procedure approved by the Sussex University Schools of Psychology and Life Sciences Research Ethics Committee. Experimental procedures complied with the Code of Ethics of the World Medical Association (Declaration of Helsinki). The task was run as a component of a one hour scanner session. The acquisition phase of the task was presented as a semantic categorisation task, and consisted of 100 words (all of which were 6 letters long) presented sequentially. Each word was presented at a central point on-screen for 1 second. There was a variable ISI of 2.5 to 4.5 seconds. A mask (#####) was presented between each stimulus. Participants were simply instructed to make a button press response to any word that described a profession, of which there were 8, quasi-randomly distributed throughout the set, such that there were two profession words in each quarter. The acquisition phase duration was approximately 7.5 minutes. The surprise recognition phase began approximately 35 minutes after the acquisition phase. In the intervening period, participants completed some structural imaging and a vigilance task in the scanner (outcomes reported elsewhere). In the recognition phase, 180 words (the 100 words seen previously, plus 80 new words) were presented in random order using the same timings as in the acquisition phase. This time, participants were instructed to respond to each word, to indicate whether they thought it was previously studied in the acquisition (categorisation task) phase ('old') or a novel word ('new'). The recognition phase lasted approximately 13.5 minutes. The words used in both the acquisition and recognition phases were drawn from the MRC psycholinguistic database (RRID:SCR_014646) (http://www.psych.rl.ac.uk/MRC_Psych_Db.html) and matched for lexico-semantic features of length (all words employed were 6 letters

long), frequency, familiarity and imageability, according to Kucera-Francis norms, as this can impact recognition performance (Bauer, et al. 2009).

fMRI recording and analysis fMRI datasets sensitive to BOLD (Blood Oxygen Level Dependent) contrast were acquired at 1.5 T (Siemens Avanto). To minimise signal artefacts originating from the sinuses, axial slices were tilted 30° from inter-commissural plane. Thirty-six 3 mm slices (0.75 mm interslice gap) were acquired with an in-plane resolution of 3 mm × 3 mm (TR = 3300 ms per volume, TE = 50 ms). Images were pre-processed using SPM8 (RRID:SCR_007037) (<http://www.fil.ion.ucl.ac.uk/spm/>). Raw T2 volumes were spatially realigned and unwarped, spatially normalised to standard space and smoothed (8 mm kernel). fMRI data were analysed with the standard hierarchal model approach employed in SPM. Design matrices were constructed for each participant's acquisition phase, which modelled subsequently remembered, subsequently forgotten, and profession sort trials as separate regressors. Design matrices were also constructed for each recognition phase, which modelled profession sort, 'Old' correct, 'Old' incorrect, 'New' correct and 'New' incorrect trials as separate regressors. Movement parameters were also entered. Processing of fMRI data was performed blind to group membership. For the acquisition phase, contrasts for subsequently remembered and forgotten trials were entered at the first level, and effects of condition (remembered/forgotten) and genotype (e4-/e4+) were analysed at the second level in a full factorial design. For the recognition phase, contrasts for 'Old' correct, 'Old' incorrect, 'New' correct and 'New' incorrect were entered at the first level. At the second level, effects of condition for correct responses (ie. 'New' correct/'Old' correct) and genotype (e4-/e4+) were analysed using a flexible factorial to test for the effects of condition and

condition by genotype interaction, followed by a 2 sample t-test (with the 2 conditions averaged) to test for main effects of genotype. In addition, a separate model examined effect of condition when participants made a ‘new’ judgement (ie. ‘New’ correct, ‘Old’ incorrect). Again, a flexible factorial followed by t-test was employed.¹

Recognition performance (proportion of studied words correctly identified) and gender were entered as covariates. The recognition performance covariate was entered to control for between-subject variance in performance; furthermore, to ensure we were only analysing data from participants who performed the task correctly (and ensure sufficient trials in the subsequently remembered condition), we excluded participants whose percentage of subsequently remembered words was <50%.

The small volume correction for the MTL was performed using a mask generated by the Wake Forest University PickAtlas (RRID:SCR_007378) (Maldjian, et al. 2003), incorporating hippocampal and parahippocampal regions. The significance threshold was set at $p < 0.05$ FWE-corrected (cluster level). When the small volume correction was applied, the significance threshold was set at $p < 0.05$ FWE-corrected (peak level). Images (Figure 1, Figure 2, Figure 3) were thresholded at $p < 0.001$ uncorrected.

Parameter estimates and 90% confidence intervals (Figure 1, Figure 3(b), (c)) were extracted using the corresponding coordinates from Tables 3 and 6 respectively.

¹ We thank an anonymous reviewer for suggesting this procedure (the original approach was to utilise SPM’s full factorial design, but for mixed within- and between- subject analyses this can be problematic as only one error term is used.) We would also like to thank the same reviewer for suggesting the multiple regression analysis (p.19).

Pupillometry recording and analysis Pupil diameter was recorded throughout the fMRI acquisition using an ASL Eyetrac 6 system with a 120Hz sampling rate. Data was converted using ASL's EyeNal software package (RRID:SCR_005997).. Data was quality checked and deemed usable for 40 participants (20 e4+ and 20 e4-). The criteria for including a participant was that >75% of data samples had to be available for all word stimuli. Intermittent tracking of the pupil, resulting in insufficient data samples, was due to use of the MRI-safe goggles, light-coloured irises, or head position in the coil. For each participant, average pupil diameter was calculated for each word (incorporating the time period when the word was on-screen, and the mask that followed it), averages were then calculated for words subsequently remembered/forgotten. Data was analysed using a within-subjects ANOVA, with gender as a covariate. Furthermore, to investigate the neural correlates of pupil diameter, average pupil diameter for each participant was added as a covariate to the full factorial model for the acquisition phase (described above). For each participant, two values were entered: average pupil diameter to remembered trials and average pupil diameter to forgotten trials. These values were entered against each participant's corresponding first-level contrast image. The effect of this covariate was then examined using a 2nd-level contrast, allowing us to determine where neural activity during forgotten and remembered trials correlated with pupil diameter in each participant.

Results

Behavioural Data

Acquisition phase

Participants were accurate in identifying the 8 profession words (see Table 2). The number of false alarms was low: Mean = 0.87, sd = 1.12. There were no effects of genotype.

Recognition phase

See Table 2. There were no effects of genotype or interactions with stimulus type ('Old'/'New'). Performance was poor, however, with a number of participants failing to recognise over half of the words presented in the acquisition phase. Participants who failed to identify at least 50% of studied words in the recognition phase were excluded from the fMRI analysis. This criterion meant that 7 e4- and 7 e4+ were excluded (Table 2a). Therefore fMRI datasets from 19 e4- and 21 e4+ were analysed, recognition performance in this group (including the discriminability index d') is shown in Table 2b, ²

² To explore the recognition performance patterns further, we investigated whether position in the word list at acquisition had an effect on likelihood of recognition. Evidence for a primacy effect was found: words presented earlier in the list were significantly more likely to be successfully classified as 'Old' in the recognition phase.

Group	Sort	'Old'	'New'
All (n=54)	0.95 ± 0.07	0.56 ± 0.14	0.78 ± 0.12
Acc>0.5 (n=40)	0.96± 0.03	0.61 ± 0.10	0.75 ± 0.12

Table 2a Proportion correct for Sort trials at acquisition (n=8), 'Old' words presented at recognition (n=92), 'New' words at recognition (n=80). Data presented for all participants, and the group included in the fMRI analyses, whose recognition performance exceeded 50%. There were no genotype effects.

	'Old'	'New'	d'
e4- (n=19)	0.60 ± 0.11	0.72 ± 0.14	0.90 ± 0.43
e4+ (n=21)	0.57± 0.10	0.77 ± 0.10	0.97 ± 0.38

Table 2b Proportion correct for 'Old' and 'New' words presented at recognition, and the discriminability index d', for the group included in the fMRI analyses (by genotype).

Neuroimaging data

Acquisition phase

Effects of genotype

The contrast e4+>e4- over both conditions (remembered/forgotten) showed no genotype effects. The contrast e4->e4+ revealed effects in left BA6 (Table 3, figure 1a).

Remembered>Forgotten

Across all subjects, significantly greater activation was seen in a left middle temporal region to subsequently remembered over forgotten trials (Table 3, figure 1b).

Interaction with genotype

No significant interaction was observed between condition (Remembered/Forgotten) and genotype.

Remembered>Forgotten in e4+

In accordance with our specific predictions, we examined activity related to Remembered>Forgotten in e4+ using a SVC incorporating bilateral parahippocampus and hippocampus. Activity was observed in left hippocampus (Table 3, figure 1c). A similar contrast in e4- showed no such activity.

Contrast	Region	Vox	x, y, z	P value
e4->e4+	Left BA4/BA6	353	-46, -17, 48	P<0.001 FWE-corrected
Remembered > Forgotten (all subjects)	Left middle temporal	172	-62, -40, -14	P=0.020 FWE corrected
Remembered > Forgotten (e4+)	Left hippocampus	11	-34, -8, -24	P=0.045 FWE-corrected after S.V.C.
Effect of pupil diameter as 2 nd -level covariate	Right BA18	120	32, -88, -6	P<0.001 FWE-corrected
	Left anterior cuneus	775	-14, -74, 28	P=0.035 FWE-corrected
	Right precuneus	129	10, -52, 36	P=0.026 FWE corrected

Table 3 Acquisition phase: fMRI results by contrast.




Figure 1 Activation maps (at $p < 0.001$ unc) and associated parameter estimates with 90% CI (F = Forgotten, R = Remembered) showing (a) Greater overall activity in left BA4/BA6 in e4-. (b) Activity in left middle temporal lobe differentiates remembered and forgotten trials. (c) Only e4+ show greater activity in left hippocampus to remembered trials.

Pupillometry data

Average pupil diameter during acquisition for subsequently remembered and forgotten words is shown in table 4. Data met all assumptions for use of parametric tests. Analysed using ANOVA, there was a main (within-subjects) effect of condition, with significantly greater pupil diameter for subsequently remembered words ($F=13.611$, $p=0.001$). There was no main effect of genotype ($F=0.003$, $p=0.953$) and no genotype by condition interaction ($F=1.623$, $p=0.210$).

		Mean (arbitrary units)	s.d.	F	p value
All subjects	Remembered	36.63	8.62	13.611	0.001
	Forgotten	35.72	9.23		
e4-	Remembered	36.72	8.64	12.913	0.002
	Forgotten	35.44	9.47		
e4+	Remembered	36.51	8.82	2.794	0.111
	Forgotten	35.90	9.21		

Table 4 Acquisition phase: Average pupil diameter to subsequently remembered and forgotten words; F and p values (2-tailed) from a repeated measures ANOVA testing for a within-subjects main effect of condition (remembered/forgotten).

Adding pupil diameter as 2nd-level covariate

To investigate the neural correlates of pupil diameter, pupil diameter was added as a covariate to the 2nd-level model. Two values were entered per participant, corresponding to the average over remembered and forgotten trials. This covariate was seen to explain variance in a posterior midline region (anterior cuneus extending into superior parietal regions), extrastriate visual cortex (BA18) and precuneus (Table 5, figure 2 (a), (b), (c)). Beta estimates for each participant by condition (forgotten/remembered) were extracted for the peak voxel in each cluster (i.e. 2 values were extracted per participant, corresponding to mean over forgotten and mean over remembered). To test for genotype effects, these were correlated against mean pupil

diameter by condition for each participant. As 6 correlations were assessed, a Bonferroni-adjusted significance threshold of $p < 0.00833$ was employed. In anterior cuneus, betas showed a negative correlation with pupil diameter for both e4- and e4+. In BA18 a significant positive correlation was seen in e4- only, while in precuneus significant negative correlation was seen in e4- only (Table 5). Plots of pupil diameter against beta estimates are included in the supplementary materials, for each of these regions. Multiple regression confirmed a main effect of genotype on the interaction with pupil diameter, in BA18 (78 voxels, 30, -84, -5, $p = 0.039$ FWE – corrected) and in precuneus (83 voxels, 8, -50, 34, $p = 0.042$ FWE – corrected).

As these correlations indicated genotype-specific effects, we then conducted ANOVA on the pupil diameter data, separately for each genotype group. A main (within-subject) effect of condition was significant only in e4- ($F = 12.91$, $p = 0.002$, Table 4).

Contrast	Region	Vox	X, y, z	P value
Old>New	Left insula	448	-32, 25, -6	$P < 0.001$ FWE-corrected
	Right insula	251	34, -26, -8	$P = 0.001$ FWE-corrected
	Left inf parietal	430	-32, -52, 32	$P < 0.001$ FWE-corrected
	Left orbitofrontal	431	-22, 64, -2	$P = 0.005$ FWE-corrected
	ACC/middle cingulate	368	-4, 35, 34	$P < 0.001$ FWE-corrected)
New>Old	Left BA18	401	-20, -86, 21	$P < 0.001$
	Right BA18	255	20, -87, 26	$P = 0.005$ FWE-corrected)
	Right Hippocampus	47	20 -9 -24	$P = 0.009$
	Left Hippocampus	12	-22 -14 -22	$P = 0.044$ FWE-corrected after bilateral S.V.C.
Interaction with genotype	Right Hippocampus	6	20 -8 -25	$P = 0.048$ FWE-corrected after bilateral S.V.C.

Table 6 Recognition phase: fMRI results by contrast.




Figure 2 Activation maps (at $p < 0.001$ unc.) showing variance explained by pupil diameter as a 2nd-level covariate in (a) BA18 (b) anterior cuneus/SPL and (c) precuneus.

Recognition phase

Correctly identified 'Old' > Correctly identified 'New' words

Significant effects were seen in bilateral insula, left inferior parietal and left orbitofrontal (see table 6). There were no main effects of genotype group.

Correctly identified 'New' > Correctly identified 'Old' words

Significant effects were seen in bilateral BA18 and bilateral hippocampus (see table 6, Figure 3). There were no main effects of genotype group.

Interaction with genotype

A significant interaction between condition (Correctly identified 'New' / Correctly identified 'Old') and genotype was seen in right hippocampus (see table 6, Figure 3(b)). A follow up t-test ('New' > 'Old') was significant in e4- (Right hippocampus, 38 vox, $p=0.043$ FWE-corrected, cluster level) but not in e4+ (3 vox, $p=0.241$ FWE-corrected, cluster level).

Recognition phase – 'New' responses

In a separate 2nd-level model, we investigated effect of condition ('New'/'Old') when participants responded 'New' (ie. contrasting correctly identified 'New' with incorrectly identified 'Old'). There was no effect of condition and no interaction with genotype.




Figure 3 Bilateral hippocampal activity to ‘New’>’Old’ contrast in recognition phase.

Parameter estimates and 90% C.I. for cluster in (b) right hippocampus (c) left hippocampus.

Discussion

In this study we set out to explore *APOE* effects on subsequent memory performance in young adults, specifically with reference to previous findings suggesting a pattern of hippocampal overactivity in e4+. In line with previous studies using subsequent memory paradigms (Dennis, et al. 2009; Filippini, et al. 2009), we found no genotype differences on recognition performance. Participants returned near-perfect scores on the sorting of profession words during the acquisition phase, indicating that they paid attention to the word stimuli. Recognition performance in the retrieval phase was necessarily reduced by the use of word, as opposed to picture, stimuli, by the employment of an incidental memory procedure, and by the forty minute filled delay between acquisition and recognition phases. Although recognition rates were low,

they followed the anticipated pattern: serial position effects were evident, with words presented nearer the beginning of the acquisition phase more likely to be recognised when re-presented forty minutes later.

For the neuroimaging data analyses, we contrasted activity to subsequently remembered against subsequently forgotten words in the acquisition phase. In the recognition phase only correct responses were considered, contrasting 'Old' against 'New' words. To ensure reliable data, we excluded participants from the neuroimaging analyses if they failed to identify at least 50% of previously studied words in the recognition phase. This meant that 7 participants from each genotype group were excluded. The poor levels of performance necessitating such exclusions should be noted as a shortcoming of this study.

At acquisition, e4+ showed less activity in BA4/BA6 relative to e4-, across both subsequently remembered and forgotten words. We have previously demonstrated genotype effects in BA6 on a covert attention task (Rusted, et al. 2013), in which young adult e4+ were faster at attentional switching. In that study, e4+ showed greater activity in BA6 and precuneus, which previous studies have linked to better performance on sustained attention tasks (Lawrence, et al. 2003); indeed, we also found young adult e4+ to show enhanced sustained attention performance (Rusted, et al. 2013). Here, e4+ were seen to consistently underactivate BA4/BA6. The attentional demands of the acquisition task used here are likely considerably less than those of the covert attention task employed previously, suggesting that activity in this region in e4+ might be more labile and sensitive to task demand than in e4-. APOE effects in BA6 have been identified at mid-age, with e4+ showing diminished left

BA6 recruitment during an object-naming task, alongside decreased activity in occipital and medial temporal lobes (Tomaszewski Farias, et al. 2005).

Based on previous findings (Dennis, et al. 2009), we expected e4+ to show greater hippocampal activity to subsequently remembered words at acquisition, compared to e4-. Consistent with this, activity in left hippocampus was seen to differentiate remembered and forgotten words in e4+ only. This demonstrates that at acquisition hippocampal overactivation in e4+ is detectable using a standard word-based subsequent memory paradigm; previous studies have employed pictorial stimuli, which are more likely to elicit hippocampal recruitment. Indeed, a study looking at effect of stimulus type has shown that although remembered picture stimuli activated bilateral MTL, activation to word stimuli did not reach significance in MTL at all (Kirchhoff, et al. 2000). Dennis et al. (2009) found bilateral hippocampal effects in e4+, consistent with the use of picture stimuli (picture stimuli engage both hemispheres, while word encoding is left lateralised (Kelley, et al. 1998)).

In the recognition phase, correctly identified 'New' and 'Old' words were contrasted and in line with previous studies (Filippini, et al. 2009), greater activity to 'New' words was seen in MTL regions, with differential activity also present in insula, cingulate, inferior parietal and early visual regions (Filippini, et al. 2009; Golby, et al. 2005). Novel stimuli elicited activity in right hippocampus, with activity in left hippocampus occurring at the trend level. Furthermore, a genotype by condition (Old/New) interaction was present in the right hippocampus. Follow-up tests showed that the hippocampal New>Old effect was significant in e4-, but not e4+. This contrasts with the findings of Filippini et al. (2009) who reported greater activity to

novel words in young adult e4+. However, their paradigm differed from ours in that participants were repeatedly familiarised with the 'old' stimuli. Work in healthy older e4+ (aged 58 to 65) similarly reported hippocampal overactivation in e4+ in a novelty paradigm (Fleisher, et al. 2005), while hippocampal activity in early-stage AD patients tends to not differentiate novel and familiar words (Golby, et al. 2005). It is not clear why the young adult e4+ under test here showed enhanced activity at acquisition specific to subsequently remembered items (while e4- did not), followed by a hippocampal underactivation to novel items at recognition. Clearly these results indicate that e4+ do not simply show a consistent pattern of hippocampal overactivity. Supporting evidence can be drawn from work by Mondadori et al. showing decreases in hippocampal activity across learning runs in an associative memory task, in young adult e4+ (Mondadori, et al. 2007). Interestingly, a study in healthy older e4+ (mean age 60) showed a similar pattern of findings. e4+ showed increased activity at acquisition to subsequently remembered items in prefrontal, temporal and parietal regions, while successful recognition was linked to lower activity in amygdala and prefrontal regions (Kukolja, et al. 2010). Since these older e4+ showed worse performance, this was interpreted as being indicative of premature neural decline. Although the study population was some four decades older than the one employed here, the authors reached the same conclusion, namely that the direction of e4+ effects on neural activity varies according to task phase.

A novel aspect of the current work was the inclusion of pupillometry measures. Pupil diameter indexes cognitive processing as well as general arousal state, and we collected pupil diameter throughout the acquisition phase. It has been suggested that the neural overactivation frequently observed in e4+ might be compensatory in nature

and reflect greater deployment of cognitive effort (Bondi, et al. 2005): we thus predicted genotype-specific effects in pupil diameter. Previous studies point to a reliable remembered/forgotten effect, where pupil diameter is greater for words that are subsequently remembered: this is thought to reflect the higher level of cognitive effort engaged to words that are subsequently remembered (Papesh, et al. 2012). Our pupillometry results showed this remembered/forgotten effect, but in e4- only.

Although there was no condition by genotype interaction, genotype-specific analyses showed that in e4+, there was no relationship between pupil diameter and whether a word was subsequently remembered or forgotten: allocation of cognitive effort to a stimulus did not predict whether it was subsequently remembered. When pupil diameter was introduced as a covariate in the fMRI analyses, it was seen to explain variance across three separate clusters in occipital lobe and precuneus, but effects were genotype-specific. Activity in extrastriate regions showed a positive relationship with pupil diameter, but only in e4-. This suggests that, in this group, greater pupil diameter is linked to enhanced processing of the word stimulus and a higher likelihood that it is subsequently remembered. A posterior midline region (encompassing posterior cuneus and superior parietal regions) showed a negative relationship across all participants. In addition, we found that activity in precuneus showed a negative relationship in e4- only. This is consistent with previous work linking DMN downregulation to subsequent memory success. The precuneus and posterior cingulate cortex form a core node of the DMN; DMN downregulation might signal a shift in attention from internal processes to external stimuli, thus increasing the likelihood of subsequent recall (Anticevic, et al. 2010; Daselaar, et al. 2004; Otten and Rugg 2001). Greater coactivation within the DMN has been previously demonstrated in young adult e4+ during the resting state (Filippini, et al. 2009;

Sheline, et al. 2010). These coactivation differences might mean that DMN shows less deactivation when attention is directed to external stimuli in e4+, which could underlie the pupillometry effects found here. Interestingly, Lustig et al. (2003) used an incidental encoding task to show that, whereas young adults showed precuneus deactivation to remembered items, healthy older adults did not. Here, precuneus activity did not covary with pupil diameter in e4+, suggesting a lack of responsivity similar to that seen in older adults, a pattern we have identified previously in mid-age e4+ (Evans, et al. 2014). However it should be noted that, since the fMRI data showed no overall main effects of genotype within the DMN, this interpretation requires further exploration.

In conclusion, we have shown that previous findings of hippocampal overactivity in young adult e4+ to subsequently remembered items generalise to a standard word-based paradigm. Typically, hippocampal activity in the acquisition phase to subsequently remembered items is shown when the paradigm includes tests of source memory or associative memory, rather than straightforward recognition judgements, suggesting that hippocampus underlies recollection, rather than familiarity-based decisions (Shrager, et al. 2008). Consequently, hippocampal activation to remembered items depends on the nature of the incidental task: when the task promotes the formation of rich episodic memories, hippocampal activation is evident (de Chastelaine and Rugg 2015). Given that e4+ showed hippocampal activity to remembered stimuli whereas e4- did not, this suggests that e4+ require hippocampal recruitment during incidental encoding if items are to be successfully recovered at recognition. This over-recruitment occurred in the context of genotype-specific effects in the pupillometry data, with links between pupil diameter, neural activity and

cognitive performance disrupted in e4+. This could be due to coactivation differences within DMN reported elsewhere. These findings (that hippocampal recruitment, rather than the deployment of cognitive effort, differentiates remembered from forgotten words in e4+) need to be explored further. Since hippocampal overactivation did not map onto pupillometry measures, it seems that if this overactivity is compensatory, it involves a mechanism not linked to cognitive effort. Indeed, deployment of cognitive effort did not link to subsequent memory performance in e4+. Interestingly, e4+ showed the opposite pattern in the recognition phase, with hippocampal activity now failing to differentiate 'new' and 'old' items. In contrast, e4- showed the normal novelty effect with hippocampus activating to novel stimuli. Although this also needs to be replicated, it does suggest that an account that posits consistent hippocampal overrecruitment in e4+ might be overly simple: while studies have reported that e4+ may recruit the hippocampus even when it is not appropriate to task demands (Rusted, et al. 2013; Trachtenberg, et al. 2012a), here e4+ failed to recruit hippocampus when it was task relevant, suggesting that hippocampal recruitment in e4+ is inconsistent, certainly abnormal, and is not always in the direction of overactivity. More work is needed to elucidate the relationship between e4 genotype, neural activity patterns and cognitive performance, but this study provides further evidence that, in young adulthood, *APOE* genotype influences brain activation patterns even when behavioural performance differences are absent.

Acknowledgements

This study was funded by a BBSRC project grant to Jenny Rusted (BB/L009242/1).

The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript. The authors have no conflicts of interest to declare. We thank Dan Goodwin for assistance with the data analysis, completed with support of a Junior Research Bursary awarded through the Sussex University Alzheimers Society Doctoral Training School.

References

- Alnaes D, Sneve MH, Espeseth T, Endestad T, van de Pavert SH, Laeng B. (2014): Pupil size signals mental effort deployed during multiple object tracking and predicts brain activity in the dorsal attention network and the locus coeruleus. *J Vis* 14(4).
- Anticevic A, Repovs G, Shulman GL, Barch DM. (2010): When less is more: TPJ and default network deactivation during encoding predicts working memory performance. *Neuroimage* 49(3):2638-48.
- Bauer LM, Olheiser EL, Altarriba J, Landi N. (2009): Word type effects in false recall: concrete, abstract, and emotion word critical lures. *Am J Psychol* 122(4):469-81.
- Bondi MW, Houston WS, Eyster LT, Brown GG. (2005): fMRI evidence of compensatory mechanisms in older adults at genetic risk for Alzheimer disease. *Neurology* 64(3):501-8.
- Bookheimer SY, Strojwas MH, Cohen MS, Saunders AM, Pericak-Vance MA, Mazziotta JC, Small GW. (2000): Patterns of brain activation in people at risk for Alzheimer's disease. *N Engl J Med* 343(7):450-6.
- Bunce D, Anstey KJ, Burns R, Christensen H, Eastaer S. (2011): Does possession of apolipoprotein E varepsilon4 benefit cognitive function in healthy young adults? *Neuropsychologia* 49(7):1693-7.
- Caselli RJ, Dueck AC, Osborne D, Sabbagh MN, Connor DJ, Ahern GL, Baxter LC, Rapcsak SZ, Shi J, Woodruff BK, Locke DE, Snyder CH, Alexander GE, Rademakers R, Reiman EM. (2009): Longitudinal modeling of age-related memory decline and the APOE epsilon4 effect. *N Engl J Med* 361(3):255-63.
- Daselaar SM, Prince SE, Cabeza R. (2004): When less means more: deactivations during encoding that predict subsequent memory. *Neuroimage* 23(3):921-927.
- Davies G, Harris SE, Reynolds CA, Payton A, Knight HM, Liewald DC, Lopez LM, Luciano M, Gow AJ, Corley J, Henderson R, Murray C, Pattie A, Fox HC, Redmond P, Lutz MW, Chiba-Falek O, Linnertz C, Saith S, Haggarty P, McNeill G, Ke X, Ollier W, Horan M, Roses AD, Ponting CP, Porteous DJ, Tenesa A, Pickles A, Starr JM, Whalley LJ, Pedersen NL, Pendleton N, Visscher PM, Deary IJ. (2012): A genome-wide association study implicates the APOE locus in nonpathological cognitive ageing. *Mol Psychiatry*.

- de Chastelaine M, Rugg MD. (2015): The effects of study task on prestimulus subsequent memory effects in the hippocampus. *Hippocampus* 25(11):1217-23.
- Dennis NA, Browndyke JN, Stokes J, Need A, Burke JR, Welsh-Bohmer KA, Cabeza R. (2009): Temporal lobe functional activity and connectivity in young adult APOE epsilon4 carriers. *Alzheimers Dement*.
- Evans S, Dowell NG, Tabet N, Tofts PS, King SL, Rusted JM. (2014): Cognitive and neural signatures of the APOE E4 allele in mid-aged adults. *Neurobiol Aging* 35(7):1615-23.
- Filbey FM, Chen G, Sunderland T, Cohen RM. (2010): Failing compensatory mechanisms during working memory in older apolipoprotein E-epsilon4 healthy adults. *Brain Imaging Behav* 4(2):177-88.
- Filippini N, MacIntosh BJ, Hough MG, Goodwin GM, Frisoni GB, Smith SM, Matthews PM, Beckmann CF, Mackay CE. (2009): Distinct patterns of brain activity in young carriers of the APOE-epsilon4 allele. *Proc Natl Acad Sci U S A* 106(17):7209-14.
- Fleisher AS, Houston WS, Eyler LT, Frye S, Jenkins C, Thal LJ, Bondi MW. (2005): Identification of Alzheimer disease risk by functional magnetic resonance imaging. *Arch Neurol* 62(12):1881-8.
- Golby A, Silverberg G, Race E, Gabrieli S, O'Shea J, Knierim K, Stebbins G, Gabrieli J. (2005): Memory encoding in Alzheimer's disease: an fMRI study of explicit and implicit memory. *Brain* 128(Pt 4):773-87.
- Just MA, Carpenter PA. (1993): The intensity dimension of thought: pupillometric indices of sentence processing. *Can J Exp Psychol* 47(2):310-39.
- Kelley WM, Miezin FM, McDermott KB, Buckner RL, Raichle ME, Cohen NJ, Ollinger JM, Akbudak E, Conturo TE, Snyder AZ, Petersen SE. (1998): Hemispheric specialization in human dorsal frontal cortex and medial temporal lobe for verbal and nonverbal memory encoding. *Neuron* 20(5):927-36.
- Kirchhoff BA, Wagner AD, Maril A, Stern CE. (2000): Prefrontal-temporal circuitry for episodic encoding and subsequent memory. *J Neurosci* 20(16):6173-80.
- Kukolja J, Thiel CM, Eggermann T, Zerres K, Fink GR. (2010): Medial temporal lobe dysfunction during encoding and retrieval of episodic memory in non-demented APOE epsilon4 carriers. *Neuroscience* 168(2):487-97.
- Lawrence NS, Ross TJ, Hoffmann R, Garavan H, Stein EA. (2003): Multiple neuronal networks mediate sustained attention. *J Cogn Neurosci* 15(7):1028-38.
- Lustig C, Snyder AZ, Bhakta M, O'Brien KC, McAvoy M, Raichle ME, Morris JC, Buckner RL. (2003): Functional deactivations: change with age and dementia of the Alzheimer type. *Proc Natl Acad Sci U S A* 100(24):14504-9.
- Maldjian JA, Laurienti PJ, Kraft RA, Burdette JH. (2003): An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. *Neuroimage* 19(3):1233-9.
- Marchant NL, King SL, Tabet N, Rusted JM. (2010): Positive effects of cholinergic stimulation favor young APOE epsilon4 carriers. *Neuropsychopharmacology* 35(5):1090-6.
- Mondadori CR, de Quervain DJ, Buchmann A, Mustovic H, Wollmer MA, Schmidt CF, Boesiger P, Hock C, Nitsch RM, Papassotiropoulos A, Henke K. (2007): Better memory and neural efficiency in young apolipoprotein E epsilon4 carriers. *Cereb Cortex* 17(8):1934-47.

- O'Dwyer L, Lambertson F, Matura S, Tanner C, Scheibe M, Miller J, Rujescu D, Prvulovic D, Hampel H. (2012): Reduced hippocampal volume in healthy young ApoE4 carriers: an MRI study. *PLoS One* 7(11):e48895.
- Otten LJ, Rugg MD. (2001): When more means less: neural activity related to unsuccessful memory encoding. *Curr Biol* 11(19):1528-30.
- Papesh MH, Goldinger SD, Hout MC. (2012): Memory strength and specificity revealed by pupillometry. *Int J Psychophysiol* 83(1):56-64.
- Rocchi A, Pellegrini S, Siciliano G, Murri L. (2003): Causative and susceptibility genes for Alzheimer's disease: a review. *Brain Res Bull* 61(1):1-24.
- Rusted JM, Evans SL, King SL, Dowell N, Tabet N, Tofts PS. (2013): APOE e4 polymorphism in young adults is associated with improved attention and indexed by distinct neural signatures. *Neuroimage* 65:364-73.
- Schlemmer KB, Kulke F, Kuchinke L, Van Der Meer E. (2005): Absolute pitch and pupillary response: effects of timbre and key color. *Psychophysiology* 42(4):465-72.
- Sheline YI, Morris JC, Snyder AZ, Price JL, Yan Z, D'Angelo G, Liu C, Dixit S, Benzinger T, Fagan A, Goate A, Mintun MA. (2010): APOE4 allele disrupts resting state fMRI connectivity in the absence of amyloid plaques or decreased CSF Abeta42. *J Neurosci* 30(50):17035-40.
- Shrager Y, Kirwan CB, Squire LR. (2008): Activity in both hippocampus and perirhinal cortex predicts the memory strength of subsequently remembered information. *Neuron* 59(4):547-53.
- Stening E, Persson J, Eriksson E, Wahlund LO, Zetterberg H, Soderlund H. (2016): Apolipoprotein E 4 is positively related to spatial performance but unrelated to hippocampal volume in healthy young adults. *Behav Brain Res* 299:11-8.
- Su YY, Liang X, Schoepf UJ, Varga-Szemes A, West HC, Qi R, Kong X, Chen HJ, Lu GM, Zhang LJ. (2015): APOE Polymorphism Affects Brain Default Mode Network in Healthy Young Adults: A STROBE Article. *Medicine (Baltimore)* 94(52):e1734.
- Thompson PM, Hayashi KM, de Zubicaray G, Janke AL, Rose SE, Semple J, Herman D, Hong MS, Dittmer SS, Doddrell DM, Toga AW. (2003): Dynamics of gray matter loss in Alzheimer's disease. *J Neurosci* 23(3):994-1005.
- Tomaszewski Farias S, Harrington G, Broomand C, Seyal M. (2005): Differences in functional MR imaging activation patterns associated with confrontation naming and responsive naming. *AJNR Am J Neuroradiol* 26(10):2492-9.
- Trachtenberg AJ, Filippini N, Cheeseman J, Duff EP, Neville MJ, Ebmeier KP, Karpe F, Mackay CE. (2012a): The effects of APOE on brain activity do not simply reflect the risk of Alzheimer's disease. *Neurobiol Aging* 33(3):618 e1-618 e13.
- Trachtenberg AJ, Filippini N, Ebmeier KP, Smith SM, Karpe F, Mackay CE. (2012b): The effects of APOE on the functional architecture of the resting brain. *Neuroimage* 59(1):565-72.
- Tuminello ER, Han SD. (2011): The apolipoprotein e antagonistic pleiotropy hypothesis: review and recommendations. *Int J Alzheimers Dis* 2011:726197.
- Wisdom NM, Callahan JL, Hawkins KA. (2011): The effects of apolipoprotein E on non-impaired cognitive functioning: a meta-analysis. *Neurobiol Aging* 32(1):63-74.
- Wishart HA, Saykin AJ, McAllister TW, Rabin LA, McDonald BC, Flashman LA, Roth RM, Mamourian AC, Tsongalis GJ, Rhodes CH. (2006): Regional brain atrophy in cognitively intact adults with a single APOE epsilon4 allele. *Neurology* 67(7):1221-4.

