

## Diagnosis and management of perinatal depression and anxiety in general practice: a meta-synthesis of qualitative studies

Article (Accepted Version)

Ford, Elizabeth, Lee, Suzanne Elizabeth, Shakespeare, Judy and Ayers, Susan (2017) Diagnosis and management of perinatal depression and anxiety in general practice: a meta-synthesis of qualitative studies. *British Journal of General Practice*, 67. e538-e546. ISSN 0960-1643

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/68053/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

### **Copyright and reuse:**

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

**A meta-synthesis of qualitative studies on general practitioners' diagnosis and management of perinatal depression and anxiety.**

**Elizabeth Ford <sup>1\*</sup>, Suzanne Lee <sup>2</sup>, Judy Shakespeare <sup>3</sup>, & Susan Ayers <sup>2</sup>**

1) Division of Primary Care and Public Health, Brighton and Sussex Medical School, Brighton, UK

2) Centre for Maternal and Child Health Research, School of Health Sciences, City, University of London, London, EC1V 0HB, UK

3) Royal College of General Practitioners, 30 Euston Square, London, NW1 2FB UK

\*Corresponding Author: Dr E. Ford, Brighton and Sussex Medical School, Department of Primary Care and Public Health, Mayfield House, Village Way, Falmer, Brighton, BN1 9PH. [e.m.ford@bsms.ac.uk](mailto:e.m.ford@bsms.ac.uk).

**Word count: 2439 excluding quotes**

**Abstract (233 words)**

**Background:** Up to 20% of women experience anxiety and depression during the perinatal period. In the UK, management of perinatal mental health falls under the remit of general practitioners (GPs).

**Aim:** This review aimed to synthesise the available information from qualitative studies on GPs' attitudes, recognition and management of perinatal anxiety and depression.

**Design & Setting:** Meta-synthesis of the available published qualitative evidence on GPs recognition and management of perinatal anxiety and depression.

**Method:** A systematic search was conducted on Embase, Medline, PsycInfo, Pubmed, Scopus and Web of Science, and grey literature was searched using Google, Google Scholar and British Library EThOS. Papers and reports were eligible for inclusion if they reported qualitatively on GPs' diagnosis or treatment of perinatal anxiety or depression. The synthesis was constructed using meta-ethnography.

**Results:** Five themes were established from five eligible papers: Labels: diagnosing depression; clinical judgement versus guidelines; care and management; use of medication; and Isolation: The role of other professionals. GPs considered perinatal depression as a psychosocial phenomenon, and were reluctant to label disorders and medicalise distress. GPs relied on their own clinical judgement more than guidelines. They reported helping patients make informed choices about treatment, and inviting women back regularly for GP visits. GPs felt isolated when dealing with perinatal mental health issues.

**Conclusion:** GPs often do not have timely access to appropriate psychological therapies and use several strategies to mitigate this shortfall. Training needs to focus on these issues and needs to be evaluated to consider if this makes a difference to outcomes for women.

## Introduction

The perinatal period lasts from the onset of pregnancy until twelve months after birth. Perinatal depressive and anxiety disorders are common: about 18% of pregnant women have depression during pregnancy (1) and 13–19% of new mothers have major or minor depression in the first year after delivery (1, 2). Anxiety is also common, with 8% experiencing generalized anxiety disorder (GAD), 3% experiencing panic disorder and 3% experiencing obsessive compulsive disorder (OCD) in pregnancy. Following birth, up to 8% experience GAD, 9% of women experience panic, 2–3% experience new onset OCD and 3% experience post-traumatic stress disorder (PTSD) (3-6). The consequences of perinatal disorders are potentially more severe and far-reaching than these disorders at other times in women's lives, having an adverse impact on the whole family if left untreated (7). Perinatal mental health is a strategic priority for health policy: while much data on costs are still missing, a recent UK report found that the annual cost to UK society of perinatal depression was £73 822 per case (\$104 574) (8), of which 70% was due to the increased risk of psychological and developmental disturbances in children (7).

In the UK, primary care is the first point of care for patients in the National Health Service (NHS) including perinatal women. This comprises general practitioners (GPs), midwives for pregnant women, and health visitors (community nurses specialised in maternal and child health) for new mothers and infants. England's National Institute for Health and Care Excellence (NICE) guidelines recommend that all primary care practitioners ask about possible depression and anxiety when women first have the contact in pregnancy and at all subsequent perinatal contacts (9). If a perinatal mental health difficulty is identified, NICE recommends the GP as the first line of assessment and management (9).

Despite GPs being in the front line of care for mental health, and the Royal College of General Practitioners (RCGP) recognizing perinatal mental health as a clinical priority (10), very little research has looked at how well GPs recognise, differentiate and manage perinatal disorders. Our recent systematic review of quantitative literature found large gaps in the literature and no studies

on disorders other than depression (11). Qualitative research can provide a more detailed understanding of the complex factors that influence patient-clinician interactions and decision-making. A number of studies have investigated women's and health visitors' views on help-seeking and disclosure of symptoms of anxiety and depression in primary care (12-14), on women's experience of care provided after disclosure (15, 16) and their preferences for taking antidepressants (17), but viewpoints of GPs have rarely been reported (18, 19).

Following a review of quantitative observational studies in the same area (11), the aim of this review was to synthesise qualitative studies on GPs' attitudes, decision making and routine clinical practice for the diagnosis and treatment of perinatal depression and anxiety in primary care.

## **Method**

### **Search Strategy**

A systematic search was conducted conforming to the PRISMA statement, between October and December 2014 on Embase, Medline, PsycInfo, Pubmed, Scopus and Web of Science. Broad search terms were used to ensure as many articles as possible were identified (e.g. general practitioner; family physician; anxiety; depression \*natal; \*partum; pregnan\*; matern\*; etc). The grey literature was searched using the same search terms on Google, Google Scholar and British Library EThOS.

The systematic search returned 8210 papers (Figure 1). After removing duplicates and inspection of the title of each paper for relevance, 7524 papers were identified as not relevant for the inclusion to this review. The abstracts of 686 papers were screened and 24 papers were scrutinised in full by two researchers (EF and FE; acknowledged). A further 1 eligible report was identified from the grey literature search.

- Figure 1 about here -

## **Eligibility**

Papers and reports were eligible for inclusion if they reported qualitatively on General Practitioners' (GPs; UK, Australia and Netherlands) or Family Physicians' (FPs; US and Canada) attitudes, decision making or routine clinical practice for the diagnosis or treatment of perinatal anxiety or depression in primary care. We defined "qualitative" very broadly to mean any results reported as text rather than numbers and mixed methods studies were included if they reported results analysed qualitatively. Papers were ineligible if they were published before 1990, did not report original research, were not published in English, GPs or FPs were not the main participants or reported as a separate group, or they reported interventions or quantitative results. At the full text stage, studies were excluded if they were not an empirical study (n=5), if they did not include GPs as the main participant (n=4), if they were randomised controlled trials evaluating an intervention (n=4) and if they reported on quantitative methods only (n=9) (studies were excluded for more than one reason so N>19). One Brazilian paper (20) was excluded because primary care in the Brazilian healthcare system was non-comparable with general practice as described in other papers.

## **Quality assessment**

There are no widely agreed criteria for quality of qualitative research (21), or quality reporting in metasyntheses (22). A checklist, based on that of Atkins et al. (21) was used to indicate the range of quality of studies and provide a means of testing the contribution of papers to the final meta-synthesis (23) but no studies were excluded on quality grounds (24). Out of 11 possible points, all studies scored 9 or 10. The checklist and results are shown in Supplementary Table A.

## **Analytic strategy**

The synthesis was constructed using the process of meta-ethnography described by Noblit and Hare (25). The papers were read and quotes identified by SL and EF. They were then re-read and key themes were identified by SL. Tables were constructed for each paper showing first and second order constructs for each theme. The definitions of these constructs was taken from Malpass et al. (23) where first order constructs are considered to be participants' "views, accounts

and interpretations”, i.e. direct quotes from participants. Second order constructs are considered to be “authors’ views and interpretations... of patients’ views”, i.e. analytic commentary on the first order constructs.

Using these tables, studies were then translated into one another using the processes of reciprocal and refutational translation (25). Quotes from participants were used to support the credibility of the new themes and to demonstrate their traceability back to the originals (26). To bring fresh insights and new understandings a line of argument synthesis was carried out so that the translated themes were organised into a logical and coherent order (27). All authors read and agreed thematic structure of results. The structure of themes is given in Supplementary Table B.

## **Results**

### **Studies**

Five papers were found which met eligibility criteria, reporting on views from 323 GPs (Table 1). Three papers, reporting on depression only, used interviews to elicit GPs’ views (28-30). One paper reported content analysis of open questions in a survey (31) and the fifth, (32) a non-peer reviewed report, covered perinatal mental health more generally. These papers were included due to the early stage of research in the area but their findings were used to support themes found in the other studies rather than initiating themes. Three papers focused on the postnatal period (28, 29, 31), one on pregnancy (30) and one on the “perinatal period” (32).

-insert Table 1 here-

### **Findings**

Five key themes were established from the data: **Labels: diagnosing depression; clinical judgement versus guidelines; care and management; use of medication; and Isolation: The role of other professionals.** Table 2 shows which themes were drawn from which studies.

- insert Table 2 here -

### **Labels: Diagnosing Depression**

GPs described conceptualising depression in psychosocial rather than biomedical terms and could be reluctant to identify the condition with a diagnostic label: *"I call it emotional turmoil rather than depression, psychological disturbance, at various stages after the birth, and I don't think of them as adjustment disorders, and often they are what I would think of as 'existential crises'"* (28, 29). This could reflect an overall approach to management and a preference for non-pharmacological interventions: *"I don't want to medicalise it too much really I think it needs to be an informal sort of network because I do think most of the time people do recover from it if they are just given some support rather than medication"* (28, 29). However it could also result from a necessarily pragmatic perspective in the face of limited service availability: *"If I call it depression, I need to do something. There's no one to refer to, so I would rather call it something else and manage her myself"* (28, 29).

GPs also referred to women's reactions in the face of diagnosis and how these could influence their definition of the condition. Some women were wary of being labelled even when they were presenting in distress: *"I mean, if they deny that they have got a problem but are still in tears, it becomes very difficult, because you can't treat somebody if they don't accept that there's something to treat"* (29). Others could be more willing to acknowledge there was something wrong: *"And equally others will just come in and say 'My husband said I've got to get this sorted out, and I need a tablet to calm me down' or whatever"* (29).

### **Clinical judgement versus guidelines**

GPs reported frequently relying on their own judgement in the detection of depression and anxiety: *"I think any kind of flatness, it's a difficult thing to explain, isn't it? You can just tell by having a conversation, just chatting to them"* (28, 29). Clinical intuition was considered a reliable tool for identifying women with symptoms in preference to formal detection instruments such as the EPDS (33) but there was some reluctance to consciously ask about symptoms: *"So I'm not saying I actively look for it, but I am hoping my antennae would tell me if there was a problem"* (28, 29).



This preference for the use of clinical judgement also extended to decisions about treatment where clinical judgement was again seen as a more appropriate decision-making tool than formal guidelines: *"I'm not a robot and doctors aren't programmed to be robots... and you get to know your patients and you know who needs an antidepressant and who doesn't"* (30). Sometimes guidelines were not followed because it was considered there was a lack of evidence to support them and the advice of trusted colleagues was perceived as more reliable: *"Depression. Most information is 'personal decision' i.e. no good evidence. Reasons for decision - local psychiatrist opinion, [hospital] pharmacist's opinion. Difficult finding up to date info"* (31). Guidelines were also not regarded as the best way of identifying the optimum management plan for individual patients: *"NICE guidelines are useful but I think you need to put your own experience into play as well, a lot of the time NICE guidelines are very strict and if you go strictly by the guidelines then quite often you don't necessarily give the patient what they need or what help they need"* (30). This reliance on individual judgement could lead to concerns about professional accountability: *"There is no clear professional guidance either and you always feel a little bit isolated when that's the case and a little bit at risk because you're kind of working off your own experience"* (30).

### **Care and management**

Some GPs described ensuring they made time for women with depression or anxiety: *"Once you kind of know they're in distress you don't just give them one session, you ask them to come back always, you get them to come back two weeks later to see how they're doing"* (29). While this approach was considered generally beneficial, it also raised its own issues: *"It's quite time consuming from the GP's point of view that you end up seeing them much more often than you would if they weren't on medication"* (30). GPs acknowledged they relied on using medication, together with seeing the patient regularly, more frequently than was ideal due to a lack of other treatment options: *"I mean, it's best if it's a multiple approach rather than just drugs. Unfortunately that's all we can offer"* (29). There was perceived to be a shortfall in the provision of talking therapies available for women: *"Services are too stretched and referrals are refused"* (32).

GPs reported that they generally involved women in decisions about their care: *"Postnatal depression. Antidepressant prescribed after long discussion with patient re: prob. areas and current literature/discussion re: safety and proven side effects. I was happy with the decision and I felt the patient was happy"* (31). This was perceived as empowering for women and likely to improve compliance with treatment and improve outcomes: *"It means giving patients the freedom and the confidence and the information they need to make their own decisions . . . I think if we can't give patients empowerment then they can't really be well or stay well"* (30). It was acknowledged that this approach should be tailored according to the needs of individual women: *'There's the doctor centred consult where it's 'What do you think doctor?' and I say what I think and I give you what I think and you go away happy or there is a different type of patient who like the patient centred consult which involves the patient's agenda. I think the key in general practice is to pick up on the cue of which patient wants which particular style"* (30). GPs also identified an occasional need for further intervention in the interests of safety: *'Patient empowerment is good, but you have to, if you felt it was harming to themselves or to their baby you would have to maybe take stronger action"* (30).

GPs' approach to the care of women with depression could be influenced by personal experience: *"Tragically it is only because of my own personal experience of severe postnatal depression 8 years ago and my struggle to find help and treatment... has the perinatal mental health of my patients become a priority for me... I am very sensitive to this in my patients and have a high pick up rate and aim to provide excellent multidisciplinary care for patient and her baby/family"* (32). It could also be altered by increased awareness of the issue: *"It is quite recent that after a workshop I became more aware of this and since then I have diagnosed about 5-7 ladies and looked after them including referral to perinatal mental health service in our area"* (32).

### **Use of medication**

GPs recognised their use of medication was influenced by a lack of other services: *"If I had easier access to counselling . . . my use of antidepressants would be much less"* (30). Some described

anxieties regarding prescribing medication to breastfeeding or pregnant mothers: *"Concerns about SSRI during breastfeeding by both me and patient. Decision making process is always fraught and made difficult by conflicting information"* (31). This anxiety occurred more frequently in relation to psychotropic drugs than other kinds of medication used in the perinatal period. There was however an acknowledgement that antidepressants were a necessary intervention for some women: *"if I felt that somebody's mental state was such that they were at risk, that their quality of life was . . . so bad that they weren't going to have a good pregnancy, I would have no problem with prescribing"* (30). GPs' concerns about prescribing for breastfeeding women sometimes resulted in women being given unnecessarily cautious advice regarding breastfeeding, but others took an evidence-based approach and stressed the importance of continued breastfeeding: *"Postnatal depression. Prescribed Zoloft [sertraline] advised to continue breastfeeding. Benefit outweigh risks. I felt okay with decision"* (31).

When GPs did wish to prescribe antidepressants, this could be met with reluctance by women: *"Patient's reluctance despite reassurance +++ . No problem for me, but patient very reluctant to take anything"* (31). Women's concerns sometimes resulted in them making decisions about their medication without consultation with their GPs: *"Women will just stop if they are on antidepressants and find out they are pregnant . . . I know they shouldn't have done that but they just panicked and said 'right, OK I'm pregnant now no more tablets'"* (30).

### ***Isolation: The role of other professionals***

GPs reported concerns that changes to the organisation of perinatal healthcare services, in particular their decreased contact with health visitors, had led to a worsening of service quality: *"[I now have] much less opportunity [to identify women]. [I] used to do joint clinics with [the] health visitor [but these have] now stopped so communication with other healthcare professionals [is] poor. I feel I am seeing fewer patients with post-natal depression which cannot be correct"* (32). Concerns included lack of continuity of service: *"Where we used to have a health visitor who was assigned to us, who we could discuss cases with, we are now assigned to a local team, so it could be anybody and it could change from day to day who the patient's health visitor is and which team they are working*

for" (28). There was also uncertainty about both their own role and that of health visitors under the new system: *"I feel my role has been marginalised since joint working with health visitors has effectively stopped"* (32); *"Because I think [health visitors] seem very constrained on what they are prepared to do really. I think that they seem just to play not a very non-interventionalist role and see themselves as being preventative, which I think is quite tragic"* (28).

Other professionals were sometimes consulted for advice regarding the management of women: *"The pharmacist at [hospital] excellent - gives various sources of information and good opinion re: overall management. If not in, she always rings you back - very reliable"* (31). This happened more frequently when the GP knew and trusted the individual professional. Otherwise, advice from others was not always perceived as useful: *"Pharmacist[s] tend to be too conservative and advise against taking anything. Also, they sometimes provide advice against what I say and alarm patients"* (31).

## **Discussion**

This meta-synthesis has highlighted that GPs consider perinatal depression as a psychosocial phenomenon rather than a biomedical one, leading to a reluctance to label disorders and medicalise distress. This finding is congruent with other commentaries on recognition and management of depression in UK general practice (34). Practitioners vary considerably in the threshold at which they will label patients as cases needing treatments because depressive symptoms are widely distributed through the population and change quickly (35). GPs see a range of social problems leading to distress and sadness, so doubt the effectiveness of antidepressants (35) and doubt that patients' problems are solvable with medical treatment (36). This can lead them to approach disclosures of mental health symptoms with reassurance, or a "watch and wait" approach.

Women may perceive this response as their symptoms being minimised and dismissed, (37-39) following which they may become reluctant to pursue treatment (40). "Watch and wait" is also potentially an inappropriate strategy in the perinatal period when suicide is a real risk (41) and

disorders may have profound impacts on the child's emotional and behavioural development (7). Some evidence suggests that when trusting relationships with healthcare professionals have time to develop the risk of dismissing new or important symptoms is diminished (42). It could be argued that rather than offering lesser treatments for perinatal women with anxiety or depression, GPs should be more proactive about initiating treatment during this vulnerable period, compared to at other times in a woman's life.

The second theme suggests that GPs rely on their own clinical judgement more than established or evidence-based guidelines. However, doctors' confidence in their decisions is not always related to their accuracy (43). When guidelines are not used in practice, unconscious biases can occur throughout doctors' interactions with patients, such as selectively gathering and interpreting evidence that confirms a diagnosis and ignoring evidence that might disconfirm it (44). The adoption of evidenced-based approaches and decision or screening tools may improve the quality of doctors' reasoning, but more research is needed to confirm this. Insight via education appears the major means in which to avoid distorted decision-making processes (45).

GPs reported on helping their patients to make informed choices about treatment, and on attempting to plug the gap in availability of "talking" therapies by inviting women to come back regularly for GP visits. They prescribed antidepressants despite recognition that a psychological therapy may be more appropriate. This suggests a tension between what GPs consider best practice and what they can practicably offer. Studies suggest GPs are aware of patients' dislike and reluctance to take antidepressants (35, 46), and would prefer to offer patients treatments aligned with their preferences.

The final theme suggests that GPs feel isolated when dealing with perinatal mental health issues. Over recent years midwives' and especially health visitors' methods of working have moved from case-loading and affiliation to a particular GP surgery, to corporate team working, where it is harder for professional relationships to develop. This may have reduced collaboration between different specialties, and may risk women losing out on joined up care. For example, Chew-Graham

et al., (28) reported health visitors as having negative attitudes to GPs, and as saying that GPs do not have a “sympathetic attitude” and would “just write a prescription”. Given that health visiting services are now commissioned by the local authority rather than the NHS, the co-ordination and continuity of care are becoming harder rather than easier within the primary care and community environment.

### **Implications for Research**

Research with GPs on how they manage perinatal depression is currently sparse, and we found none exploring perinatal anxiety or PTSD. Future research is needed at all levels of the primary care pathway, from recognition of psychological distress, to outcomes of treatment both within primary care and following referral to specialist services (47). Training and resource interventions should be evaluated to see if they improve outcomes for women, their infants and their families.

### **Clinical Implications**

Continuity of care and trusting relationships are found to be important in the literature on women’s perception of help-seeking. However, it is unlikely that GPs will have more routine antenatal contact with pregnant women in order to develop a sense of continuity, (48) or closer working relationships with midwives and health visitors in the near future. One potential strategy is for practices to have a GP lead for maternity and maternal mental health who regularly meets with local midwives and health visitors and coordinates strategy within the practice, and is visibly available for patients to consult with about perinatal mental health issues. A key issue for GPs is also to have specialist community perinatal mental health services available to refer patients to in a timely way. Very recently, there has been considerable investment in specialist perinatal mental health by the current UK government, for example, development of Mother and Baby units and specialist community teams, but little so far to address the common mental health problems which are usually managed in primary care.

### **Strengths and Limitations**

Our search strategy was comprehensive so it is likely we captured all available literature, and our methodology for synthesising the papers was robust. However, only one researcher screened titles and abstracts, which may have made study selection unreliable. The small number of studies and small samples mean that these findings only represent a narrow range of views, are not generalizable, and are likely to be subject to selection bias. Additionally, because of the small range of literature available, we included a lower quality study which used open-ended survey responses and a non-peer reviewed report. The survey study was focussed on prescribing for breastfeeding women rather than perinatal mental health, thus results from this study could only support rather than initiate themes. Additionally we found no evidence for how perinatal anxiety is recognised or managed in primary care. Studies all originated from English speaking countries (UK and Australia) and given that four of the five were UK based, these results are very UK focussed. Much more research is needed in this area to confirm these findings and set them in context, and explore how GPs manage perinatal mental health in other countries.

## **Conclusions**

This meta-synthesis shows that GPs consider perinatal depression within the context of women's lives and are frustrated at the lack of talking therapy resources they have available. It is clear that GPs try to plug the gap in mental health services by inviting women back regularly, thus developing a potentially therapeutic trusting relationship. Much more research is needed in this area, and particularly in how GPs manage perinatal anxiety, to inform training and resource interventions. Where interventions are implemented, they must be evaluated to consider if they make a difference to outcomes for women.

## **Competing interests**

The authors have no competing interests.

## **Acknowledgements**

We gratefully acknowledge the help of Maithreyee Vipulanathan and Fatin Elias in conducting the literature searches.

## References

1. Gavin NI, Gaynes BN, Lohr KN, Meltzer-Brody S, Gartlehner G, Swinson T. Perinatal depression: a systematic review of prevalence and incidence. *Obstet Gynecol* 2005;106(5, Part 1):1071-83.
2. O'Hara MW, Swain AM. Rates and risk of postpartum depression; a meta-analysis. *Int Rev Psychiatry*. 1996;8(1):37-54.
3. Zambaldi CF, Cantilino A, Montenegro AC, Paes JA, de Albuquerque TLC, Sougey EB. Postpartum obsessive-compulsive disorder: prevalence and clinical characteristics. *Compr Psychiatry* 2009; 50(6): 503-9
4. Ross LE, McLean LM. Anxiety disorders during pregnancy and the postpartum period: A systematic review. *J Clin Psychiatry*. 2006;67(8):1285-98.
5. Grekin R, O'Hara MW. Prevalence and Risk Factors of Postpartum Posttraumatic Stress Disorder: A Meta-Analysis. *Clin Psychol Rev*. 2014;34(5):389-401.
6. Farr SL, Dietz PM, O'Hara MW, Burley K, Ko JY. Postpartum anxiety and comorbid depression in a population-based sample of women. *J Womens Health* 2014;23(2):120-8.
7. Stein A, Pearson RM, Goodman SH, Rapa E, Rahman A, McCallum M, et al. Effects of perinatal mental disorders on the fetus and child. *Lancet*. 2014;384(9956):1800-19.
8. Bauer A, Parsonage M, Knapp M, Lemmi V, Adelaja B. The costs of perinatal mental health problems. 2014. [http://www.centreformentalhealth.org.uk/pdfs/Costs\\_of\\_perinatal\\_mh.pdf](http://www.centreformentalhealth.org.uk/pdfs/Costs_of_perinatal_mh.pdf)
9. National Institute for Health and Care Excellence. Antenatal and postnatal mental health: clinical management and service guidance. London UK: National Institute for Health and Care Excellence; 2014.
10. Ayers S, Shakespeare J. Should perinatal mental health be everyone's business? *Prim Health Care Res Dev*. 2015;16(04):323-5.
11. Ford E, Shakespeare J, Elias F, Ayers S. Recognition and management of perinatal depression and anxiety by general practitioners: A systematic review. *Fam Pract*. submitted.
12. Dennis CL, Chung-Lee L. Postpartum depression help-seeking barriers and maternal treatment preferences: a qualitative systematic review. *Birth*. 2006;33(4):323-31.
13. Slade P, Morrell CJ, Rigby A, Ricci K, Spittlehouse J, Brugha TS. Postnatal women's experiences of management of depressive symptoms: a qualitative study. *Br J Gen Pract*. 2010;60(580):e440-e8.
14. Russell K, Ashley A, Chan G, Gibson S, Jones R. *Maternal Mental Health – Women's Voices*. London, UK: Royal College of Obstetricians and Gynaecologists, 2017.
15. Shakespeare J, Blake F, Garcia J. How do women with postnatal depression experience listening visits in primary care? A qualitative interview study. *J Reprod Infant Psychol*. 2006;24(02):149-62.
16. Sword W, Busser D, Ganann R, McMillan T, Swinton M. Women's care-seeking experiences after referral for postpartum depression. *Qual Health Res*. 2008;18(9):1161-73.
17. Turner KM, Sharp D, Folkes L, Chew-Graham C. Women's views and experiences of antidepressants as a treatment for postnatal depression: a qualitative study. *Fam Pract*. 2008;25(6):450-5.



18. Oates MR, Cox JL, Neema S, Asten P, Glangeaud-Freudenthal N, Figueiredo B, et al. Postnatal depression across countries and cultures: a qualitative study. *Br J Psychiatry*. 2004;184(46):s10-s6.
19. Boots Family Trust Alliance. Perinatal Mental Health: Experiences of Women and Health Professionals 2013 [22 October 2016]. Available from: <http://www.bftalliance.co.uk/the-report/>.
20. Santos Junior HP, Gualda DMR, Silveira MdFA, Hall WA. Postpartum depression: the (in) experience of Brazilian primary healthcare professionals. *Journal of advanced nursing* 2013; **69**(6): 1248-58.
21. Atkins S, Lewin S, Smith H, Engel M, Fretheim A, Volmink J. Conducting a meta-ethnography of qualitative literature: lessons learnt. *BMC Med Res Methodol*. 2008;8(1):21.
22. Dheensa S, Metcalfe A, Williams RA. Men's experiences of antenatal screening: A metasynthesis of the qualitative research. *Int J Nurs Stud*. 2013;50(1):121-33.
23. Malpass A, Shaw A, Sharp D, Walter F, Feder G, Ridd M, et al. "Medication career" or "Moral career"? The two sides of managing antidepressants: A meta-ethnography of patients' experience of antidepressants. *Soc Sci Med*. 2009;68(1):154-68.
24. Sandelowski M, Docherty S, Emden C. Focus on qualitative methods Qualitative metasynthesis: issues and techniques. *Res Nurs Health*. 1997;20:365-72.
25. Noblit GW, Hare RD. *Meta-ethnography: Synthesizing qualitative studies*: Sage; 1988.
26. Jensen LA, Allen MN. Meta-synthesis of qualitative findings. *Qual Health Res*. 1996;6(4):553-60.
27. Walsh D, Downe S. Meta-synthesis method for qualitative research: a literature review. *J Adv Nurs*. 2005;50(2):204-11.
28. Chew-Graham C, Chamberlain E, Turner K, Folkes L, Caulfield L, Sharp D. GPs' and health visitors' views on the diagnosis and management of postnatal depression: a qualitative study. *Br J Gen Pract*. 2008;58(548):169-76.
29. Chew-Graham CA, Sharp D, Chamberlain E, Folkes L, Turner KM. Disclosure of symptoms of postnatal depression, the perspectives of health professionals and women: a qualitative study. *BMC Fam Pract*. 2009;10(7):[9p.].
30. McCauley C-O, Casson K. A qualitative study into how guidelines facilitate general practitioners to empower women to make decisions regarding antidepressant use in pregnancy. *International Journal of Mental Health Promotion*. 2013;15(1):3-28.
31. Jayawickrama HS, Amir LH, Pirotta MV. GPs' decision-making when prescribing medicines for breastfeeding women: Content analysis of a survey. *BMC Res Notes*. 2010;3:82.
32. Khan L. *Falling through the gaps: perinatal mental health and general practice*. London UK: Centre for Mental Health; 2015.
33. Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatry*. 1987;150(6):782-6.
34. Thomas-MacLean R, Stoppard JM. Physicians' constructions of depression: inside/outside the boundaries of medicalization. *Health*. 2004;8(3):275-93.
35. Kendrick T. Why can't GPs follow guidelines on depression? *Brit Med J*. 2000;320(7229):200-1.
36. Maxwell M. Women's and doctors' accounts of their experiences of depression in primary care: the influence of social and moral reasoning on patients' and doctors' decisions. *Chronic Illn*. 2005;1(1):61-71.
37. McIntosh J. Postpartum depression: women's help-seeking behaviour and perceptions of cause. *J Adv Nurs*. 1993;18(2):178-84.
38. Mauthner NS. Postnatal depression: how can midwives help? *Midwifery*. 1997;13(4):163-71.

39. Amankwaa LC. Postpartum depression among African-American women. *Issues Ment Health Nurs.* 2003;24(3):297-316.
40. Edge D, Baker D, Rogers A. Perinatal depression among black Caribbean women. *Health Soc Care Community.* 2004;12(5):430-8.
41. Knight M, Kenyon S, Brocklehurst P, et al. *Saving Lives, Improving Mothers' Care - Lessons learned to inform future maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2009–12.* Oxford: National Perinatal Epidemiology Unit, University of Oxford, 2014.
42. Megnin-Viggars O, Symington I, Howard L, Pilling S. Experience of care for mental health problems in the antenatal or postnatal period for women in the UK: a systematic review and meta-synthesis of qualitative research. *Arch Women's Ment Health.* 2015;18(6):745-59.
43. Berner E, Graber M. Overconfidence as a cause of diagnostic error in medicine *Am J Med.* 2008;121(5 Suppl):s2-s3.
44. Moskowitz AJ, Kuipers BJ, Kassirer JP. Dealing with Uncertainty, Risks, and Tradeoffs in Clinical Decisions: A Cognitive Science Approach. *Ann Intern Med.* 1988;108(3):435-49.
45. Hall KH. Reviewing intuitive decision-making and uncertainty: the implications for medical education. *Med Educ.* 2002;36(3):216-24.
46. Hyde J, Calnan M, Prior L, Lewis G, Kessler D, Sharp D. A qualitative study exploring how GPs decide to prescribe antidepressants. *Br J Gen Pract.* 2005;55(519):755-62.
47. Goldberg DP, Huxley P. *Mental Illness in the Community: The Pathway to Psychiatric Care.* London: Tavistock Publications Ltd; 1980.
48. The National Maternity Review. *Better Births. Improving outcomes in maternity services in England. A Five Year Forward View for maternity care.* 2016 [25 October 2016]. Available from: <https://www.england.nhs.uk/wp-content/uploads/2016/02/national-maternity-review-report.pdf>.

Table 1: Study Characteristics

Authors	Country	Methods	Sample	Sampling Approach/ Response Rate	Primary Objective
Chew-Graham et al (2008)	UK	In-depth interviews, thematic analysis.	Purposive sample of 19 GPs recruited from participants in multi-centre RCT (RESPOND - Randomised Evaluation of antidepressants and Support for women with P <sub>O</sub> stNatal Depression)	Sampling was purposive and sought to achieve maximum variation in relation to GPs' age, gender, length of time in general practice, practice size and level of deprivation.	To explore the views of GPs and health visitors on the diagnosis and management of <b>postnatal</b> depression.
Chew-Graham et al(2009)	UK	In-depth interviews, thematic analysis.	Same sample as Chew-Graham (2008) above.	As above	To explore GPs', health visitors' and women's views on the disclosure of symptoms which may indicate depression in primary care.

Jaywickerama et al (2010)	Australia	Anonymous postal survey, content analysis.	335 GPs: 70% female 37% aged 45-54 84% obtained medical degree in Australia 90% had children 49% of them (or their partners) had >12 months experience of breastfeeding.	125/640 (19.5%) GPs responded to survey and provided open ended comments on prescribing decisions, 54 GPs (8.4%) mentioned depression.	Explore GPs' decision making when they are considering recommending or prescribing medication for a breastfeeding woman.
McCauley & Casson (2013)	UK (Northern Ireland)	Semi-structured interviews, Colaizzi's process of analysis.	8 GPs: 2 male, 6 female.	10 Practice managers were invited to identify GPs who were eligible for involvement, 8 GPs were identified.	Develop an in-depth understanding of GPs' experience of using guidelines in the treatment of perinatal depression and if this enabled them to empower women to become involved in treatment decisions.
Khan (2015)	Mainly UK	Postal survey plus semi-structured	43 GPs:	The GP survey was	To better understand the

		interview with 3 survey respondents, interpretive phenomenology.	40 from England, 2 from Wales & Scotland, 1 from India. Over half had <11 years' experience in general practice, just over a third had practised for 1-3 years. Just over a quarter had >20 years' experience. 14% felt they held a partially specialist role in perinatal mental healthcare.	distributed to an unknown but large number of GPs through virtual portals. Only 43 GPs responded.	contribution of GPs to the area of perinatal mental health.
--	--	--	---	---	---

**Table 2: Themes drawn from the five studies**

<b>Theme</b> <b>Paper</b>	<b>Labels: Diagnosing Depression</b>	<b>Clinical judgement versus guidelines</b>	<b>Care and management</b>	<b>Use of medication</b>	<b>Isolation: Role of other professionals</b>
Chew-Graham et al (2008) (24)	Yes	Yes	Yes	-	Yes
Chew-Graham et al (2009) (25)	Yes	Yes	Yes	-	-
Jayawickrama et al (2010) (27)	-	Yes	Yes	Yes	Yes
McCauley & Casson (2013) (29)	-	Yes	Yes	Yes	-
Khan (2015) (28)	-	-	Yes	-	Yes



Supplementary Table B: Development of Themes

Concept	1st Order Evidence	2nd Order Evidence
<p data-bbox="203 381 327 411"><b>Labelling</b></p> <p data-bbox="203 459 360 568"><b>(Theme: Diagnosing depression)</b></p>	<p data-bbox="461 381 925 411"><b>Chew-Graham et al (2008 &amp; 2009):</b></p> <p data-bbox="461 459 1227 647">'I call it emotional turmoil rather than depression... psychological disturbance, at various stages after the birth, and I don't think of them as adjustment disorders, and often they are what I would think of as 'existential crises'.' (GP,M1)</p> <p data-bbox="461 695 1227 842">'I don't want to medicalise it too much really I think it needs to be an informal sort of network because I do think most of the time people do recover from it if they are just given some support rather than medication.' (GP, M8)</p> <p data-bbox="461 890 1227 999">'If I call it depression, I need to do something. There's no one to refer to, so I would rather call it something else and manage her myself.' (GP,M10)</p> <p data-bbox="461 1046 819 1077"><b>Chew-Graham et al (2009):</b></p> <p data-bbox="461 1125 1227 1272">"I mean, if they deny that they have got a problem but are still in tears, it becomes very difficult, because you can't treat somebody if they don't accept that there's something to treat." (B GP1)</p> <p data-bbox="461 1319 1227 1391">"and equally others will just come in and say 'my husband said I've got to get this sorted out, and I need a tablet to</p>	<p data-bbox="1256 381 1615 411"><b>Chew-Graham et al (2008):</b></p> <p data-bbox="1256 459 1966 606">All responders attributed a psychosocial aetiology to postnatal depression and demonstrated ambivalence about the status of postnatal depression as a separate condition.</p> <p data-bbox="1256 654 2018 801">GPs described a variety of strategies for managing women in the postnatal period, and how the label they used for the woman's problems determined what management strategies they employed.</p> <p data-bbox="1256 849 2024 995">GPs were reluctant to ascribe a biomedical label to women with symptoms that could indicate postnatal depression because they perceived there was a lack of resources to which they could refer women.</p> <p data-bbox="1256 1086 1615 1117"><b>Chew-Graham et al (2009):</b></p> <p data-bbox="1256 1165 2018 1353">Health professionals also attributed a psychosocial aetiology to postnatal depression and demonstrated ambivalence about the status of postnatal depression as a separate condition as compared with depressive illness at other times in a woman's life.</p>



	<p>calm me down' or whatever. You get the whole spectrum, really." (M GP2)</p>	<p>Some GPs described a reluctance to use the term postnatal depression because they felt that symptoms would recover without formal interventions, because of a lack of services or referral options, and the feeling that antidepressants were the only treatment option available.</p> <p>GPs described difficulties in using the label for postnatal depression with women, particularly referring to the stigma that they perceived women felt, and the effect on this on the consultation. Other GPs, however, described consultations where the woman was happy to accept the label.</p>
<p><b>Care</b></p> <p><b>(Theme: Care and management)</b></p>	<p><b>Chew-Graham et al (2008):</b></p> <p>'In the old days I would have said, you know, there's an ongoing relationship with the health visitor, would have said "fine", you know, they can monitor her, they can keep an eye on her that's what she used to do, you know, keep in close contact.' (GP, L1)</p> <p>'It's about normalising how they think ... I don't always offer anything ... I very rarely prescribe on the first visit ... so I think many depressions are like PND actually, they exist in the context of somebody's life and it has a meaning for them which you have to attend to ... I hate these things where they say if you've got five tick-boxes then you have 6 months of fluoxetine, that's gross.' (GP, M1)</p>	<p><b>Chew-Graham et al (2008):</b></p> <p>GPs described a variety of strategies for managing women in the postnatal period, and how the label they used for the woman's problems determined what management strategies they employed.</p> <p>Some GPs emphasised the important role health visitors could play in the management of postnatal depression and their assumptions about how they worked.</p> <p>GPs were aware of these changes in the organisation of health visiting services and described the impact on their day-to-day relationships with health visitors, as well as confusion over the expected roles and responsibilities of the health visitor.</p>

'I think it depends very much on the skills and the experience of the health visitor but I think very much about helping the women too. Providing some sort of support, I guess, someone to talk to and listen to but also perhaps, one hopes, that you are giving people some structure and some practical things to do in order to maybe to cope with crying babies and, you know, poor sleep, and perhaps lack of support in the house.' (GP, M10)

**Chew-Graham et al (2009):**

"Once you kind of know they're in distress you don't just give them one session, you ask them to come back always...you get them to come back two weeks later to see how they're doing." (L GP1)

"I mean, it's best if it's a multiple approach rather than just drugs. Unfortunately that's all we can offer." (L GP1)

"Easier not to ask, if I'm not going to see her again." (L GP1)

**Jaywickerama et al (2010):**

"Postnatal depression. Antidepressant prescribed after long discussion with patient re: prob. areas and current

**Chew-Graham et al (2009):**

Some GPs described strategies used to facilitate disclosure and offer women ongoing support.

Some GPs described a reluctance to use the term postnatal depression because they felt that symptoms would recover without formal interventions, because of a lack of services or referral options, and the feeling that antidepressants were the only treatment option available.

Both GPs and health visitors, however, described the current systems of care as hindering disclosure of symptoms of postnatal depression.

**Jaywickerama et al (2010):**

Many GPs discussed the situation and medicines issues at length with the woman herself before arriving at a decision especially with regard to postnatal depression.

literature/discussion re: safety and proven side effects. I was happy with the decision and I felt the patient was happy." (ID 22)

**McCauley & Casson (2013):**

‘you need to have the information first, know what you’re talking about, give them the information . . . I think it’s like everything if you give them enough information and then give them a bit of support . . . you’re empowering them and helping them to make better decisions and help with their compliance.’ (P7)

‘it’s quite time consuming from the GP’s point of view that you end up seeing them much more often than you would if they weren’t on medication.’ (P2)

‘it means giving patients the freedom and the confidence and the information they need to make their own decisions . . . just in general. I think if we can’t give patients empowerment . . . then they can’t really be well or stay well.’ (P2)

‘Patient empowerment . . . is good, but you have to . . . if you felt it was harming to themselves or to their baby you would have to maybe take stronger action. (P8)

‘how you explain something differs from patient to patient and some people will want to know everything and some

**McCauley & Casson (2013):**

GPs described how providing information to women encouraged them to become involved in the decision-making process. However, they acknowledged that, as they have no specific or accurate guidance, it is difficult to confidently provide information to women to enable them to make informed decisions.

Treatment decisions are complicated, which most of the GPs identified, by the lack of available resources within the timescale of the pregnancy.

All GPs believed involving women in decision making was a major component of patient empowerment. It was recognized that empowerment was significant for women to remain well and to make healthy decisions, with GPs believing that they could give women empowerment by providing them the information to make their own decisions.

Some GPs in areas of higher deprivation reported differing views, feeling some women had a variable attitude towards involvement in decision making that was related to their attitude towards their own level of empowerment. Most GPs also expressed varying attitudes on patient empowerment, feeling that it was not always appropriate in clinical practice.

	<p>people trust you and don't want to know anything. The usual line is 'I trust you, ye know, you're the doctor.'" (P4)</p> <p>'general practice consultation . . . has two ends of a spectrum, at one end there's the doctor centred consult where it's . . . 'What do you think doctor?' and I say what I think and I give you what I think and you go away happy or there is a different type of patient who like the patient centred consult which involves the patient's agenda . . . I think the key in general practice is to pick up on the cue of which patient wants which particular style. (P3)</p> <p>'some patients like it and some patients don't and it's really engaging the patient in the consultation and in the decision making and trying to give them ownership of it . . . in Derry, an awful lot of my patients turn it back on me when I try and empower them and involve them in the decisions . . . and they say 'Aye right! You're the doctor . . . you tell me that's what they pay ye for!'" (P1)</p> <p>'you are restricted sometimes the service that you want isn't available, but certainly if I had easier access to counselling . . . so I would know that someone who is depressed and vulnerable is able to go and see someone that week or the next week, on a weekly basis . . . my use of antidepressants would be much less.' (P4)</p> <p><b>Khan (2015):</b></p>	<p>They described what they communicated to women was influenced by their own perception of how much information they felt some women wanted and were able to understand... They described how they used their professional experience to determine the level of involvement women wanted to have and this influenced how decisions were made.</p> <p>Some GPs in areas of higher deprivation reported differing views, feeling some women had a variable attitude towards involvement in decision making that was related to their attitude towards their own level of empowerment. Most GPs also expressed varying attitudes on patient empowerment, feeling that it was not always appropriate in clinical practice.</p> <p>Treatment decisions are complicated, which most of the GPs identified, by the lack of available resources within the timescale of the pregnancy.</p> <p><b>Khan (2015):</b></p> <p>In qualitative responses, a minority of GPs noted that 'lived experience' or increased training had noticeably altered their effectiveness in identifying women with such difficulties.</p>
--	---	---

<p>It is quite recent that after a workshop I became more aware of this and since then I have diagnosed about 5-7 ladies and looked after them including referral to perinatal mental health service in our area. I try to review them myself to keep the continuity of care. (GP survey respondent)</p> <p>Tragically it is only because of my own personal experience of severe postnatal depression 8 years ago and my struggle to find help and treatment (eventually by paying privately to see a Perinatal Psychiatrist and subsequent reading of NICE guidelines, personal reading etc) has the perinatal mental health of my patients become a priority for me... I am very sensitive to this in my patients and have a high pick up rate and aim to provide excellent multidisciplinary care for patient and her baby/family. (GP survey respondent)</p> <p>We are lucky to have fairly easily accessible clinics locally, but I know from personal experience that these are definitely not available nationwide. In [the] distant past [we had one] attached to our practice – which was when we did have a working postnatal depression protocol. (GP, survey respondent)</p> <p>Specialist care is patchy, and the waiting times are unacceptable. (GP survey respondent)</p> <p>Services are too stretched and referrals are refused. (GP survey respondent)</p>	<p>Both mothers and GPs also made references to postcode inconsistencies in responses and in the range of care available.</p> <p>Nearly half of GPs responding to the RCGP survey reported not trusting the quality of local referral routes with over a third of survey respondents feeling that the greatest barrier to providing women with effective support was a lack of reliable pathways in their local area.</p>
--	---

	<p>I have or have had several patients who have experienced post-natal mental health problems. Our local service is purely antenatal, and for those under consultant care, which to me seems to miss when these problems often develop. The local priority feels low. We have a specialist psychologist within our psychology department - but no access to her as GPs. (GP survey respondent)</p>	
<p><b>Use of medication</b>  <b>(Theme: Use of medication)</b></p>	<p><b>Jaywickerama et al (2010):</b></p> <p>"Concerns about SSRI during breastfeeding by both me and patient. Decision making process is always fraught and made difficult by conflicting information." (ID 115)</p> <p>"I think the risks of depression (postpartum) often outweigh the risks of the antidepressants. I felt that there are no right answers to the problem." (ID 24)</p> <p>"Patient's reluctance despite reassurance +++. No problem for me, but patient very reluctant to take anything." (ID 14)<sup>7</sup></p> <p>"Patient concern is very high" (ID 164)</p> <p>"Public perception is they can't take anything. This may partly be impacting on low uptake of breastfeeding." (ID 5)</p> <p>"Recently prescribed Cipramil [citalopram] for postnatal depression to a woman who was breastfeeding. I felt the risk to the baby was low &amp; the drug was important to the woman. I felt reassured that I have been to talks where</p>	<p><b>Jaywickerama et al (2010):</b></p> <p>GPs either reported positive feelings which were mainly associated with a certainty about the decision or negative feelings when they were less certain. Negative feelings (<i>difficult, concern, doubt</i>) were more evident with prescribing antidepressants than with antibiotics.</p> <p>The fifth organising theme concerned issues of prescribing for one person, the breastfeeding woman, thereby exposing the vulnerable "third party" - the breastfed child - to the effects of the medicine.</p>

	<p>psychiatrists have said they use this drug in lactating women." (ID 78)</p> <p>"Be careful &amp; know the potential dangers." (ID 68)</p> <p><b>McCauley &amp; Casson (2013):</b></p> <p>'women will just stop if they are on antidepressants and find out they are pregnant . . . I know they shouldn't have done that but they just panicked and said 'right, OK I'm pregnant now no more tablets.' (P7)</p> <p>'if I felt that somebody's mental state was such that they were at risk, that their quality of life was . . . so bad that they weren't going to have a good pregnancy, I would have no problem with prescribing.' (P5)</p> <p>"SSRI in a breastfeeding woman. That it was acceptable. I had to look back at past Australian Doctor [magazine] articles b/c [because] the online sources of MIMS info was too overcautious. OK once I had read the article. I feel able to make an informed decision." (ID 161)</p> <p>"A woman needed Flagyl [metronidazole] for ?anaerobic infection. Information accessed via MIMS Annual (internet). Decided to express + discard for 1 wk + 3 days + formula feed, then resumed thereafter". (ID 304)</p>	<p><b>McCauley &amp; Casson (2013):</b></p> <p>They described how women felt anxious about the effects of medication on their baby's development and often abruptly discontinued their antidepressants.</p> <p>A woman's quality of life and her enjoyment of her pregnancy would be an influence in the use of antidepressants.</p> <p>Non-availability of easily accessible, evidence based, up to date information on medicines in breastfeeding was mentioned. GPs often mentioned that their sources of information were conflicting and often "over cautious".</p> <p>GPs advised cessation of breastfeeding and the introduction of infant formula in several instances, even in situations that did not warrant such measures.</p> <p>However many GPs stressed the importance of continued breastfeeding together with medicines.</p>
--	--	--

	<p>"Postnatal depression. Prescribed Zoloft [sertraline] advised to continue breastfeeding. Benefit outweigh risks. I felt Okay with decision." (ID 138)</p>	
<p><b>Judgement v guidelines</b></p> <p><b>(Theme: Clinical judgement versus guidelines)</b></p>	<p><b>Chew-Graham et al (2008 &amp; 2009):</b></p> <p>"So I'm not saying I actively look for it, but I am hoping my antennae would tell me if there was a problem." (M GP5)</p> <p>"I think any kind of flatness...it's a difficult thing to explain, isn't it?...You can just tell by having a conversation... just chatting to them." (B HV1)</p> <p><b>Jaywickerama et al (2010):</b></p> <p>"Depression. Most information is 'personal decision' i.e. no good evidence. Reasons for decision - local psychiatrist opinion, RWH pharmacist's opinion. Difficult finding up to date info." (ID 152)</p>	<p><b>Chew-Graham et al (2008):</b></p> <p>GPs and health visitors described a reliance on instinct or clinical intuition which would alert them to the possibility of pnd; they did not then go on to use a schedule such as the Edinburgh Postnatal Depression Scale. Neither did they use such a schedule to screen women. Thus, postnatal depression was viewed as a social response to birth and there was an apparent reluctance by GPs and health visitors to actively look for depression in postnatal women.</p> <p><b>Chew-Graham et al (2009):</b></p> <p>GPs and health visitors described a reliance on instinct or clinical intuition, which would alert them to the possibility of postnatal depression, rather than using formal screening instruments or actively seeking out symptoms of depression.</p> <p><b>Jaywickerama et al (2010):</b></p> <p>Non-availability of easily accessible, evidence based, up to date information on medicines in breastfeeding was mentioned. GPs often mentioned that their sources of information were conflicting and often "over cautious".</p>



	<p><b>McCauley &amp; Casson (2013):</b></p> <p>‘there is no clear professional guidance either and you always feel a little bit isolated when that’s the case and . . . a little bit at risk because your . . . kinda working off your own experience...and nobody can argue with your own experience but again nobody can support it except yourself.’ (P2)</p> <p>‘NICE guidelines are useful but I think . . . you need to put your own experience into play as well, a lot of the time NICE guidelines are very . . . strict and if you go . . . strictly by the guidelines then quite often you don’t necessarily . . . give the patient what they need or what help they need.’ (P8)</p> <p>‘I’m not a robot and doctors aren’t programmed to be robots. You don’t just type into a pile of stuff into my head and I just spit it all out . . . and ye get to know your patients and you know who needs an antidepressant and who doesn’t.’ (P6)</p> <p>‘I’m not aware of what the guidelines are for this . . . I presume there probably are guidelines but at times there’s that many guidelines that it can be slightly overwhelming, and ye don’t have time to read them all’ (P3)</p> <p>‘we’re not really getting any definite guidelines . . . the NICE guidelines aren’t great in this regard . . . a bit vague and not pinpointing one specific antidepressant.’ (P7)</p>	<p><b>McCauley &amp; Casson (2013):</b></p> <p>Most GPs recognized that they often relied on their own professional experience rather than guidelines in order to make complex risk–benefit treatment decisions for pregnant women.</p> <p>Some GPs acknowledged that they found guidelines not only generic but also restrictive, stating that no drug was safe, leaving them with limited options in cases of severe depression. They described how guidelines did not enable GPs to practice with the flexibility and knowledge that their training had encouraged and, if followed strictly, patient need would not be met.</p> <p>It was recognized by all the GPs that guidelines were not always applicable as the benefits and risks of treatment will be different for each individual woman. Their own knowledge of individual women and their needs was a key element of how perinatal depression, in practice, was managed.</p> <p>The majority of GPs interviewed identified low usage of guidelines in practice. Most GPs described how they received a large number of guidelines each week, limiting the time available to read all in detail and, as a result, did not feel that they had full knowledge of guidelines for perinatal depression.</p>
--	--	---

	<p>'I suppose in a way guidelines protect you a bit, in terms of if you do something that is recommended in the guidelines you have that protection . . . you're using them in that way, practising defensive medicine which I don't particularly like . . . you should be able to justify things outta your own knowledge without having to . . . fall back on guidelines.' (P5)</p>	<p>All GPs reported that the guidelines they had read in relation to the treatment of depression in pregnancy lacked specific and clear direction on treatment in the perinatal period.</p> <p>However, they acknowledged that, despite the limitations of the guidelines, they represent best practice advice. They described all guidelines by providing a professional reference point, which can be used as a defence against litigation in case of adverse reactions. Some GPs acknowledged that, in this regard, they offer protection, but it was not their ideal way to practice.</p>
<p><b>Other professionals</b></p> <p><b>(Theme: Isolation: the role of other professionals)</b></p>	<p><b>Chew-Graham et al (2008):</b></p> <p>'because I think they seem very constrained on what they are prepared to do really. I think that they seem just to play not a very noninterventionalist role and see themselves as being preventative, which I think is quite tragic because there is lots of ... if you don't integrate sort of preventative curative resources it's not a great service really.' (GP, M6)</p> <p>'That's another difficulty because the set up of health visitors is being reorganised so where we used to have a health visitor who was assigned to us, who we could discuss cases with, we are now assigned to a local team, so it could be anybody and it could change from day to day who the patient's health visitor is and which team they are</p>	<p><b>Chew-Graham et al (2008):</b></p> <p>Other GPs reported observing an unwillingness of health visitors to manage women with postnatal depression.</p> <p>GPs were aware of these changes in the organisation of health visiting services and described the impact on their day-to-day relationships with health visitors, as well as confusion over the expected roles and responsibilities of the health visitor.</p> <p>GPs described their experience of health visitors declining to support women with what was dismissed as a 'mental health problem', although they agreed that health visiting teams still had a role in offering practical support to women with postnatal depression.</p>

working for. Which makes it very difficult to work, and I see no sense in it. Also, they seem to have dropped a lot of their duties in relation to new mothers. And, I do wonder what health visitors are doing now.' (GP, M7)

'I would say that of all the practitioners here they are the least integrated and they are the least, erm, we get least communication with them ... there's nothing wrong with them, they're perfectly nice and friendly, but there's just not that clinical sort of connectiveness really.' (GP, B3)

'Well, our health visitors tend to say that it's a mental health problem "nothing really to do with me", which is disappointing really. They do go in and offer support but it's very vague what that support is. They cover some practical things around sort of nursery services for the children, stuff like that and it depends which health visitor team it is because not all will take an interest.' (GP, M7)

**Jaywickerama et al (2010):**

"Depression ... Consulted RWH book, psychiatrist, discussion with pt. [patient] re risk etc." (ID 199)

"The pharmacist at RWH excellent - gives various sources of information and good opinion re: overall management. If not in, she always rings you back - very reliable." (ID 301)

**Jaywickerama et al (2010):**

Before prescribing for a breastfeeding woman, many GPs needed to check sources for information on safety of the medicine; this was not straight forward as sources gave conflicting responses, and sometimes pharmacists' opinion on medicines in breastfeeding was at odds with their own decision.

Non-availability of easily accessible, evidence based, up to date information on medicines in breastfeeding was

	<p>"I would appreciate ready access to detailed information - books often get misplaced, so internet access would be great." (ID 60)</p> <p>"We need a dedicated reliable easy access source." (ID 164)</p> <p>"Depression. Efexor [venlafaxine]. Checked with Box Hill Hospital pharmacist via phone. Got the most reliable and up to date info but the information took hours to obtain. ie. too long." (ID 321)<sup>3</sup></p> <p>"Pharmacist[s] tend to be too conservative and advise against taking anything. Also, they sometimes provide advice against what I say and alarm patients" (ID 246)</p> <p>"Had to prescribe an antidepressant. Discussed situation with patient and her psychiatrist ... OK with decision as it involved team care co-ordination" (ID 104)</p> <p><b>Khan (2015):</b></p> <p>I feel my role has been marginalised since joint working with health visitors has effectively stopped. (GP, survey respondent)</p> <p>[We have a] brilliant midwife who runs antenatal clinics in the surgery. This [facilitates]...good communication</p>	<p>mentioned. GPs often mentioned that their sources of information were conflicting and often "over cautious".</p> <p>Several GPs brought up the issue of pharmacists' advice regarding medicines in breastfeeding conflicting with their decision.</p> <p>GPs felt that certain situations warranted involvement of several parties in the decision-making process rather than a quick decision on their part. Although this would involve more time and work for GPs, they thought this would help to make a more appropriate and safe decision and increase mothers' compliance with the recommended/prescribed medicine.</p> <p><b>Khan (2015):</b></p> <p>A few GPs also referred in qualitative comments to their increasing distance from antenatal care due to organisational changes placing some midwives and health visitors outside practice surgeries</p> <p>Some women and GPs described benefitting from close team work approaches supporting mothers' mental health: recent changes in the organisation of perinatal care (particularly reduced GP co-location with midwives and health visitors in some areas) and more unreliable communication and information sharing was seen</p>
--	---	--

	<p>between midwife and GP, excellent team work and good continuity of care. All mothers-to-be with perinatal mental health issues are identified during [the] booking appointment with [the] midwife – [then the] form [is] faxed to [the] local secondary care team to add to records. (GP, survey respondent)</p> <p>[It is important to identify women] at any opportunity - the midwife sees patients during their pregnancy, so we may not see the patient until after she has delivered, unless there are problems with her mental health. (GP, survey respondent)</p> <p>[I now have] much less opportunity [to identify women]. [I] used to do joint clinics with [the] health visitor [but these have] now stopped so communication with other healthcare professionals [is] poor. I feel I am seeing fewer patients with post-natal depression which cannot be correct. (GP, survey respondent)</p>	<p>sometimes to limit GP involvement in earlier identification and support:</p>
--	---	---

Figure 1. Flow diagram of study selection

