Reheating with a composite Higgs boson

Article (Published Version)

Croon, Croon, Sanz, Verónica and Tarrant, Ewan R M (2016) Reheating with a composite Higgs boson. *Physical Review D*, 94 (4). 045010. ISSN 2470-0010

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/69463/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Reheating with a composite Higgs boson

Djuna Croon, Verónica Sanz, and Ewan R. M. Tarrant

Department of Physics and Astronomy, University of Sussex, Brighton BN1 9QH, UK

(Received 28 October 2015; published 19 August 2016)

The flatness of the inflaton potential and lightness of the Higgs boson could have the common origin of the breaking of a global symmetry. This scenario provides a unified framework of Goldstone inflation and composite Higgs models, where the inflaton and the Higgs particle both have a pseudo-Goldstone boson nature. The inflaton reheats the Universe via decays to the Higgs and subsequent secondary production of other SM particles via the top and massive vector bosons. We find that inflationary predictions and perturbative reheating conditions are consistent with cosmic microwave background data for sub-Planckian values of the fields, as well as opening up the possibility of inflation at the TeV scale. We explore this exciting possibility, leading to an interplay between collider data cosmological constraints.

DOI: [10.1103/PhysRevD.94.045010](https://doi.org/10.1103/PhysRevD.94.045010)**I. INTRODUCTION**

Scalar fields are popular protagonists in cosmological theories. They play chief roles in the leading paradigms for important events, such as inflation and electroweak symmetry breaking. However, it has been long known that fundamental scalars suffer radiative hierarchy problems: for theory to match observations, one requires an unnatural cancellation of UV corrections. In inflation, this radiative instability can be quantified by the tension between the Lyth bound [1] on the slow roll phase of the field, pushing towards $\Delta\phi > M_p$, and the measurement of CMB anisotropies, which indicate $\Lambda_{\text{inf}} \lesssim 10^{15}$ GeV. For electroweak symmetry breaking (EWSB), one usually considers the large separation of scales between the Higgs mass and the Planck scale as an illustration, as the latter is where the theory should be cut off for an elementary Higgs.

Here we will discuss the appeal of pseudo-Goldstone bosons (pGBs) for the dynamical generation of scales in both paradigms. The realization that Goldstone bosons can solve hierarchy problems is not new: for EWSB, there is popular branch of model building that goes by composite Higgs theory which postulates a new strongly coupled sector of which the Higgs is a bound state [2] (for a review see [3]). The effective theory then has a cutoff, such that the Higgs mass is not sensitive to effects above the compositeness scale.

Likewise, in inflationary model building “Natural Inflation” provides an inflaton candidate protected from UV corrections using essentially the same mechanism with an axionic GB [4]. Alas, vanilla Natural Inflation requires trans-Planckian scales to predict the measured cosmic microwave background (CMB) spectrum and thus has

questionable value as a valid effective theory.¹ In [12] the idea of a pGB inflaton was generalized, and it was shown there and in [13] that different models may realize inflation compatible with data from the cosmic microwave background (CMB) without the issues that the original Natural Inflation has.

In this paper we will show how both mechanisms can be unified, thus realizing radiative stability for both models in a single simple setup. We will explore the minimal symmetry breaking pattern that realizes a Higgs $SU(2)$ doublet and an inflaton singlet. We discuss both the generation of an inflaton potential and reheating in this model. Interestingly, both can be fully perturbative processes. The inflationary predictions are shown to be compatible with the latest CMB data by Planck [14] without the necessity of introducing trans-Planckian scales in the effective theory. After inflation the inflaton decays into Higgs bosons, which subsequently decay into the Standard Model particles. Importantly, we find that the question if reheating can take place perturbatively crucially depends on the CP assignment in the model.

We will finish by showing how the model naturally connects to electroweak physics. The inflaton mass and couplings to the Higgs could be of the same order, leading to the possibility of looking for the inflaton through their mixing with the Higgs.

In Fig. 1 we show a graphic of the relevant scales in our model. The global symmetry is broken at the scale f , which is below the Planck scale at which we expect a UV

¹There have been several proposals to explain the trans-Planckian decay constant while maintaining the simple potential and the explanatory power of the model. Among these are Extra-natural Inflation [5], hybrid axion models [6,7], N-flation [8,9], axion monodromy [10] and other pseudonatural inflation models in supersymmetry [11].

FIG. 1. Relevant scales: pseudo-Goldstone bosons naturally realize mass hierarchies. CMB data and constraints on perturbative reheating allow us to relate the complete spectrum to the symmetry breaking scale f and the Planck scale M_P .

completion in the form of a theory of quantum gravity. The scale of inflation is then expected to be parametrically smaller than f , as we will show. The Coleman Weinberg masses of the goldstone boson inflaton and Higgs are fixed by CMB and electroweak data respectively. Likewise, the values of the coefficients of the (self-)couplings in the potential can be fixed in light of the data, modulo the scale of inflation. This is a free parameter in our model. As usual for slow roll inflation, it is most naturally found around the grand unified theory scale (10^{15} GeV), but can be as low as $\sim 10^5$ GeV if one allows for a degree of tuning.

Finally we would like to highlight some recent developments that may be of interest to the reader. In [15] a dynamical solution to the electroweak hierarchy problem was proposed, in terms of a Higgs boson coupling to an inflaton and an axionlike field. Although critics have pointed out several shortcomings, among which the necessity of a very large number of e-foldings and the low cutoff (which makes one arguably expect new physics around the EW scale) [16], the scanning mechanism is a new facet worth investigating. As the model behind the mechanism bares similarities with our setup, it seems like a worthwhile exercise to look for a realization in the present context. A second recent result that is interesting in the present context is the observation in [17] that the Higgs-inflaton coupling $c_4 h^2 \eta^2$ may drastically alter the Higgs dynamics in the early Universe, thereby stabilizing the electroweak vacuum. As we will see the coupling c_4 will automatically be present in our model.

II. THE LAGRANGIAN OF THE HIGGS AND THE INFLATON

A. Inflaton-Higgs couplings for perturbative reheating

The condition that the inflaton field must decay completely into relativistic particles to complete the reheating process dictates the interaction structure in a successful theory of inflation. After the end of inflation, the inflaton field η begins to oscillate about the minimum of its potential with amplitude $\Phi(t)$. The Universe is completely dominated by the zero mode, $\langle \eta(t) \rangle$, which may be interpreted as a

condensate of nonrelativistic zero-momentum η particles of mass m_η . The condensate oscillation amplitude decays as $\Phi(t) \sim t^{-1}$ due to the Hubble expansion and due to interactions with the Higgs field. Trilinear couplings, $\frac{1}{2} \sigma \eta h^2$, and quartic couplings, $\frac{1}{2} g^2 \eta^2 h^2$, with the Higgs are to be expected on fairly general grounds, as we argue in the following section. As we will show in Sec. IV, provided that the coupling constants σ, g^2 and the amplitude $\Phi(t)$ are small enough such that nonperturbative particle production processes are absent, the energy loss experienced by the condensate can be described by the Boltzmann equation

$$\frac{d}{dt}(a^3 \rho_\eta) = -\frac{\sigma^2 \Phi_0^2 m_\eta}{64\pi} - \frac{g^4 \Phi_0^4 m_\eta}{128\pi a^3}, \quad (1)$$

where a is the scale factor and Φ_0 is the initial amplitude of the inflaton oscillations at the start of reheating. The contribution from the quartic interaction decreases as $a^{-3} \sim t^{-2}$, which, as is well known [18–20], poses a major problem for theories which do not contain a trilinear interaction. Specifically, since the Hubble rate decreases as $H \sim a^{-3/2} \sim t^{-1}$, volume dilution due to the Hubble expansion takes place faster than the annihilation process $\phi\phi \rightarrow \chi\chi$ can drain energy from the condensate and so reheating never completes. In order to successfully reheat the universe, a trilinear coupling must be present. We will use this result as a guiding principle when constructing the Lagrangian for the composite Higgs model.

B. Symmetry breaking: The minimal coset

The inflaton and Higgs correspond to five scalar degrees of freedom which could come from the breaking of $SO(6)$ to $SO(5)$ or, equivalently $SU(4)$ to $Sp(4)$. This breaking pattern is very popular in building models of composite Higgs, as it preserves custodial symmetry.

The breaking gives rise to five Goldstone bosons, transforming as a **5** of $SO(5)$. The most general vacuum which breaks $SO(6) \rightarrow SO(5) \sim SU(4) \rightarrow Sp(4)$ as shown in Ref. [21] is given by²

$$\Sigma_0 = \begin{pmatrix} 0 & e^{i\alpha} \cos(\theta) & \sin(\theta) & 0 \\ -e^{i\alpha} \cos(\theta) & 0 & 0 & \sin(\theta) \\ -\sin(\theta) & 0 & 0 & -e^{-i\alpha} \cos(\theta) \\ 0 & -\sin(\theta) & e^{-i\alpha} \cos(\theta) & 0 \end{pmatrix} \quad (2)$$

where α and θ are real angles. One recovers a well-known choice of vacuum in Composite Higgs models [22] in the limit $\alpha \rightarrow \text{mod}(\pi)$ and $\theta \rightarrow \text{mod}(\pi)$.

²The discussion in Ref. [21] assumes the presence of CP conserving vacua, as well as CP breaking vacua, such that the Pfaffian of the inflaton is real.

In fact, the vacuum in which we have $\theta = \text{mod}(\pi)$ has an enhanced custodial symmetry [21], as in this case the unbroken generators generate $SU(2) \times SU(2) \subset Sp(4)$. Likewise, the limit $\alpha = \text{mod}(\pi)$ parametrizes the conservation of CP by the vacuum.

One can then parametrize the Goldstone bosons via the field $\Sigma(x)$,

$$\Sigma(x) = e^{i\Pi^a(x)T_\perp^a/\sqrt{2}f}\Sigma_0, \quad (3)$$

where $\Pi^a(x)$ are the Goldstone fields with decay constant f , corresponding to the broken $SO(6) \cong SU(4)$ generators

$T_\perp^a$. A linear combination of three of the Goldstone fields is eaten by the Standard Model gauge fields such that the corresponding generators can be recognized as their longitudinal components. The two remaining Goldstone bosons remain in the spectrum as massless scalar fields and couple via the broken generators $T_\perp^4$ and $T_\perp^5$.³

$$T_\perp^4 = \begin{pmatrix} 0 & \sigma_2 \\ \sigma_2 & 0 \end{pmatrix}, \quad T_\perp^5 = \begin{pmatrix} c_\theta e^{i\alpha} \mathbb{1}_2 & -is_\theta \sigma_2 \\ is_\theta \sigma_2 & c_\theta e^{i\alpha} \mathbb{1}_2 \end{pmatrix}. \quad (4)$$

Expanding the matrix exponential, we obtain

$$\Sigma(x) = \begin{pmatrix} c_\pi + \frac{\sqrt{2}f i e^{i\alpha} c_\theta s_\pi \eta}{\sqrt{\pi_a^2}} & 0 & 0 & \frac{i\sqrt{2}f s_\pi (-ih - s_\theta \eta)}{\sqrt{\pi_a^2}} \\ 0 & c_\pi + \frac{\sqrt{2}f i e^{i\alpha} c_\theta s_\pi \eta}{\sqrt{\pi_a^2}} & \frac{i\sqrt{2}f s_\pi (ih + s_\theta \eta)}{\sqrt{\pi_a^2}} & 0 \\ 0 & \frac{i\sqrt{2}f s_\pi (s_\theta \eta - ih)}{\sqrt{\pi_a^2}} & c_\pi - \frac{i\sqrt{2}f e^{i\alpha} c_\theta s_\pi \eta}{\sqrt{\pi_a^2}} & 0 \\ \frac{i\sqrt{2}f s_\pi (ih - s_\theta \eta)}{\sqrt{\pi_a^2}} & 0 & 0 & c_\pi - \frac{i\sqrt{2}f e^{i\alpha} c_\theta s_\pi \eta}{\sqrt{\pi_a^2}} \end{pmatrix} \Sigma_0 \quad (5)$$

where we have suppressed space-time dependence of the fields $h = h(x)$ and $\eta = \eta(x)$, and where we use the shorthands

$$h(x)^2 + \eta(x)^2 = \pi_a^2 \quad \text{and} \quad s_\pi = \sin\left(\frac{\sqrt{\pi_a^2}}{\sqrt{2}f}\right), \quad c_\pi = \cos\left(\frac{\sqrt{\pi_a^2}}{\sqrt{2}f}\right) \\ s_\theta = \sin(\theta), \quad c_\theta = \cos(\theta). \quad (6)$$

We will further assume that gauging the theory breaks $SU(4)$ to the Standard Model group⁴ $SU(2)_L \times U(1)_Y$ and $U(1)_\eta$. This latter shift symmetry for η will assure that it does not get a potential from gauge bosons. Then the kinetic term becomes

$$\frac{f^2}{8} \text{Tr}|D_\mu \Sigma|^2 = \frac{1}{2} \frac{(\eta \partial_\mu h - h \partial_\mu \eta)^2}{h^2 + \eta^2} \\ + \frac{g^2}{4} h^2 \left(W_\mu^+ W^{-\mu} + \frac{1}{\cos^2 \theta_w} Z_\mu Z^\mu \right) \\ \approx \frac{1}{2} (\partial_\mu h)^2 + \frac{1}{2} (\partial_\mu \eta)^2 + \frac{1}{2} \frac{(h \partial_\mu h + \eta \partial_\mu \eta)^2}{1 - h^2 - \eta^2} \\ + \frac{g^2}{4} h^2 \left(W_\mu^+ W^{-\mu} + \frac{1}{\cos^2 \theta_w} Z_\mu Z^\mu \right) \quad (7)$$

³Here we use generalized expressions from Ref. [21]; obtained by assuming the general vacuum [Eq. (A17)] in the rotation Eq. (B25).

⁴Here we do not address the color group $SU(3)_c$.

where the following field redefinitions are made:

$$h^2 s_\pi^2 f^2 / \pi_a^2 \rightarrow h^2 \quad \eta^2 s_\pi^2 f^2 / \pi_a^2 \rightarrow \eta^2 \\ (\partial_\mu h s_\pi f / \sqrt{\pi_a^2})^2 \rightarrow (\partial_\mu h)^2 (\partial_\mu \eta s_\pi f / \sqrt{\pi_a^2})^2 \rightarrow (\partial_\mu \eta)^2 \quad (8)$$

corresponding to dropping the operators with more than four powers in the field (they will be effectively suppressed by f). For the sigma model, there is an equivalence between the original and rotated fields. However, the rotated fields couple to gauge bosons as in (7) and are as such the physically relevant choice.

At this level, the η and h fields are true Goldstone bosons. (Small) explicit breaking of the symmetry will generate a Coleman-Weinberg contribution to the scalar potential, via gauge and Yukawa interactions. This potential accounts, then, for resummations of loops of gauge bosons and fermions. Rather than considering the fully generic case, we can use the information from the previous section as prior information about what a Lagrangian which gives perturbative reheating will look like. In particular, the necessity of terms with odd powers of the singlet η in the scalar potential implies that the singlet η has specific transformation properties under CP that differ from the composite Higgs model. This can be understood in the following way: if we for a moment assume that CP is unbroken, we can set $\alpha = 0$. As we will see, the way we parametrize the coupling between η and (Dirac) fermions can schematically be written as

$$\eta \bar{F}(c_{\text{even}} + ic_{\text{odd}}\gamma_5)F. \quad (9)$$

Clearly, for $c_{\text{odd}} = 0$, η behaves as a scalar, such that the trilinear interaction ηh^2 is allowed by the symmetry. However in the composite Higgs case ($c_{\text{odd}} \neq 0$) where η behaves as a (partial) pseudoscalar, the term ηh^2 breaks CP .

In contrast, the breaking of the enhanced custodial symmetry by taking $\theta \neq 0$ does not have such a direct impact on the predictions for perturbative reheating. It is expected to give rise to mass mixing, i.e. terms of the form $V \ni c_i \eta h$. Deviations from custodial symmetry in the Higgs sector are rather constrained by low-energy data and it will therefore be practical to assume $\theta = 0$ in the following. This choice corresponds to identifying the Higgs with the bi-doublet under the subgroup $SO(4) \cong SU(2)_L \times SU(2)_R$, and η with the singlet: $\mathbf{1} \oplus \mathbf{4} = (\mathbf{1}, \mathbf{1}) \oplus (\mathbf{2}, \mathbf{2})$.

As the scalar η does not couple to the $SU(2)_L$ gauge group, see Eq. (7), couplings to gauge bosons do not help with generating a cubic term. The difference in dynamics between the different vacua has to come from the couplings to fermions.

As an example, we implement the fermions in a **6** of $SU(4)$ [corresponding to the vector representation of $SO(6)$]. Other options for fermion representations, such as **4** and the **10**, have their own difficulties to address [22].

The **6** of $SU(4)$ decomposes as $(\mathbf{2}, \mathbf{2}) \oplus (\mathbf{1}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{1})$ under $SU(2)_L \times SU(2)_R$, such that we can implement the fermions as [22]

$$\Psi_q = \frac{1}{2} \begin{pmatrix} 0 & Q \\ -Q^T & 0 \end{pmatrix} \quad \Psi_u = \Psi_u^+ + \Psi_u^- \quad \Psi_u^\pm = \frac{1}{2} \begin{pmatrix} \pm U & 0 \\ 0 & U \end{pmatrix} \quad (10a)$$

$$\Psi_{q'} = \frac{1}{2} \begin{pmatrix} 0 & Q' \\ -Q'^T & 0 \end{pmatrix} \quad \Psi_d = \Psi_d^+ + \epsilon_d \Psi_d^- \quad \Psi_d^\pm = \frac{1}{2} \begin{pmatrix} \pm D & 0 \\ 0 & D \end{pmatrix} \quad (10b)$$

where $Q = (0, q_L)$, $Q' = (q_L, 0)$, $U = u_R i\sigma_2$ and $D = d_R i\sigma_2$. The $\epsilon_{u,d}$ are complex free parameters defining the embedding of the quarks into the singlets, and consecutively the CP -assignment of η . In the limit $|\epsilon_{u,d}| = 1$ the fermions have definite charges under $U(1)_\eta$ and it is therefore expected that η is massless.

The coupling of Σ to fermions will be of the form

$$\begin{aligned} \mathcal{L}_{\text{eff}} = & \sum_{r=q,u,q',d} [\Pi'_0 \text{Tr}[\bar{\Psi}_r \not{p} \Psi_r] + \Pi'_1 \text{Tr}[\bar{\Psi}_r \Sigma] \not{p} \text{Tr}[\Psi_r \Sigma^\dagger]] \\ & + M_u \text{Tr}[\bar{\Psi}_q \Sigma] \text{Tr}[\Psi_u \Sigma^\dagger] + M_d \text{Tr}[\bar{\Psi}_{q'} \Sigma] \text{Tr}[\Psi_d \Sigma^\dagger]. \end{aligned} \quad (11)$$

C. Composite Higgs limit: CP assignment in the fermion sector

As we show in the Appendix, loops of fermions and gauge bosons will generate a Coleman-Weinberg potential at one loop, which will be of the form [22]

$$V(\kappa, h) = a_1 h^2 + \lambda h^4 + |\kappa|^2 (a_2 + a_3 h^2 + a_4 |\kappa|^2) \quad \text{where} \quad \kappa = \sqrt{f^2 - \eta^2 - h^2} + i\epsilon_t \eta \quad (12)$$

where a_i are dimensionful constants dependent on the form factors of the UV theory as given in the Appendix. Here ϵ_t is the parameter that defines the embedding of the up-type fermion in the global symmetry and determines the mass and CP assignment of η , as we demonstrated above. It is easy to see that the scenario in which ϵ_t is real is distinctly different from the case in which it can be complex. For $\epsilon_t \in \mathbb{R}$, we find that η behaves like a pseudoscalar [$c_{\text{odd}} \neq 0$ and $c_{\text{even}} = 0$ in (9)], and we can expand (12) to obtain the following CP and custodially symmetric potential:

$$V(\eta, h) = m_h^2 h^2 + \lambda_h h^4 + m_\eta^2 \eta^2 + \lambda_\eta \eta^4 + c_4 \eta^2 h^2. \quad (13)$$

Here, in terms of the parameters above we have defined

$$m_h^2 = (a_1 + a_3 - a_2 - a_4), \quad (14a)$$

$$\lambda_h = (\lambda - a_3 + a_4), \quad (14b)$$

$$m_\eta^2 = (1 - \epsilon_t^2)(-a_2 - a_4), \quad (14c)$$

$$\lambda_\eta = (1 - \epsilon_t^2)^2 a_4, \quad (14d)$$

$$c_4 = (1 - \epsilon_t^2)(-a_3 + 2a_4), \quad (14e)$$

and as announced the trilinear term is absent. If we allow for complex coupling to fermions,

$$\epsilon_t = \epsilon_t^{\text{RE}} + i\epsilon_t^{\text{IM}} \quad (15)$$

where $\epsilon_t^{\text{IM}} \neq 0$, we will find η has $c_{\text{even}} \neq 0$ in (9).⁵ In this case the scalar potential will include a trilinear interaction and a tadpole for η , both of which multiply ϵ_t^{IM} ,

$$V = c_{\text{tad}} \eta + m_\eta^2 \eta^2 + \tilde{c}_\eta \eta^3 + \lambda_\eta \eta^4 + m_h^2 h^2 + \lambda_h h^4 + c_3 \eta h^2 + c_4 \eta^2 h^2 \quad (16)$$

where

$$\tilde{c}_\eta = 4a_4 \epsilon_t^{\text{IM}} (1 - (\epsilon_t^{\text{RE}})^2) \sqrt{f^2 - \eta^2 - h^2}, \quad (17a)$$

⁵In the boundary case $\epsilon_t^{\text{RE}} = 0$, $\epsilon_t^{\text{IM}} \neq 0$ η behaves like a scalar.

$$c_3 = (4a_4 - 2a_3)(\epsilon^{\text{IM}})\sqrt{f^2 - \eta^2 - h^2}, \quad (17b)$$

$$c_4 = (a_3 - 2a_4)(\epsilon^{\text{RE}})^2 - 4a_4(\epsilon^{\text{IM}})^2 + 2a_4 - a_3, \quad (17c)$$

and the other coefficients remain as above. The tadpole and trilinear interaction term violate CP for $\epsilon_i^{\text{RE}} \neq 0$. We may shift away the tadpole $c_{\text{tad}}\eta$ by an appropriate vacuum expectation value v_η , which solves

$$c_{\text{tad}} + 2m_\eta^2 v_\eta + 3\tilde{c}_\eta v_\eta^2 + 4\lambda_\eta v_\eta^3 = 0. \quad (18)$$

This will also shift the parameters,

$$m_\eta^2 \rightarrow m_\eta^2 + 3\tilde{c}_\eta v_\eta + 6\lambda_\eta v_\eta^2, \quad (19a)$$

$$\tilde{c}_\eta \rightarrow \tilde{c}_\eta + 4v_\eta \lambda_\eta, \quad (19b)$$

$$m_h^2 \rightarrow m_h^2 + c_3 v_\eta + c_4 v_\eta^2, \quad (19c)$$

$$c_3 \rightarrow c_3 + 2c_4 v_\eta. \quad (19d)$$

In terms of the shifted parameters the potential becomes

$$V = m_\eta^2 \eta^2 + \tilde{c}_\eta \eta^3 + \lambda_\eta \eta^4 + m_h^2 h^2 + \lambda_h h^4 + c_3 \eta h^2 + c_4 \eta^2 h^2. \quad (20)$$

This potential has the required form to be a suitable candidate for inflation followed by perturbative reheating.

D. Spontaneously broken CP by the inflaton ($\alpha \neq 0$)

For the composite Higgs vacuum discussed above $\alpha = 0$ and CP is unbroken by the vacuum. Here we relax this constraint, and consider the CP breaking vacuum with CP breaking in the model to

$$0 < \alpha \leq 1/2\pi \quad (21)$$

For $\alpha = 1/2\pi$ both fields have a quadratic term and do not interact. For the open interval, $0 < \alpha < 1/2\pi$, we indeed find the same potential as at the end of the previous sector, to fourth order in the fields:

$$V(\eta, h) = m_\eta^2 \eta^2 + \tilde{c}_\eta \eta^3 + \lambda_\eta \eta^4 + m_h^2 h^2 + \lambda_h h^4 + c_3 \eta h^2 + c_4 \eta^2 h^2. \quad (22)$$

The coefficients are in general nonzero, except for at $\alpha = 1/4\pi$. We refer the reader to the Appendix for a discussion, and an example computation. Importantly, in these vacua we are not required to introduce explicit CP breaking by a complex fermion representation to get the η -odd terms as we were for $\alpha = 0$, that is, we may have either $\in \mathbb{R}$ or $\in \mathbb{C}$.

In these vacua the η field couples directly to fermions as

$$(\eta \bar{u}_R \not{p} u_R) \in \mathcal{L}, \quad (23)$$

an effect proportional to $(1-^2)$. Indeed, is seen that the odd powers of η in the potential (which includes the trilinear coupling) are multiplied by $(1-^2)$ and $(b_1 - b_2)^2$ for some constants b_i (from the linear and the second order expansion of the logarithm respectively). This combination plays the role that ϵ_i^{IM} played in the previous section, as an order parameter of CP breaking.

As expected from periodicity, the two quadrants $0 < \alpha < 1/2\pi$ and $1/2\pi < \alpha < \pi$ are equivalent, modulo a redefinition of the fields⁶:

$$\eta \rightarrow -\eta \quad \text{and} \quad h \rightarrow -h. \quad (24)$$

We demonstrate this explicitly in the Appendix.

We will finish this section with a comment on the appearance of domain walls [23]. As we introduced the possibility of breaking CP spontaneously, one may be worried that these will be present, and become energetically important. However, if the vacuum breaks CP spontaneously, it does it at the scale of symmetry breaking f . But, as we will see in the next section, we expect inflation to occur below this scale, $\Lambda_{\text{inf}} < f$, hence the domain walls will be diluted during inflation.

III. INFLATION

In this section we study inflation due to the field η . As the scale of inflation will turn out to be much larger than the electroweak scale, the Higgs field would be stabilized at the minimum of its potential during inflation, and so we set $h = 0$. Hence, we neglect the dynamics of the Higgs field during inflation, and the model is effectively single field. We can canonically normalize the inflationary sector via the field redefinition

$$\phi = f \arcsin(\eta/f), \quad (25)$$

such that the scalar potential becomes, in the unbroken CP limit,

$$V_{CP}(\phi) = m_\eta^2 f^2 \sin(\phi/f)^2 + \lambda_\eta f^4 \sin(\phi/f)^4. \quad (26)$$

This is equivalent to the Goldstone inflation [13] potential

$$V(\phi) = \Lambda^4 (\sin^2(\phi/f) - \tilde{\beta} \sin^4(\phi/f)), \quad (27)$$

if we identify

⁶Because of custodial symmetry, which shows up here as a Z_2 symmetry for h , $h \rightarrow -h$ is a symmetry over the whole range. The latter substitution is therefore made for free.

FIG. 2. Form of the potential: Here $\lambda_\eta f^2 \gtrsim -1/2m_\eta^2$.

$$\lambda_\eta f^4 = -\tilde{\beta}\Lambda^4 \quad \text{and} \quad m_\eta^2 f^2 = \Lambda^4.$$

In Fig. 2 we show a plot of the form of the potential, for the moment with $\tilde{c}_\eta/m_\eta^2 = 0$. This model would lead to inflation with $f < M_p$ (where M_p is the reduced planck mass) and spectral index within the bounds allowed by Planck (at 2σ) [14],

$$n_s = [0.948-0.982] \quad \text{for } \tilde{\beta} \lesssim 1/2 \rightarrow \lambda_\eta f^2 \gtrsim -1/2m_\eta^2.$$

As in Goldstone inflation, the sensitivity to the exact value of $\tilde{\beta}$ that predicts the right spectral index is a function of $(f/M_p)^2$:

$$4 \times 10^{-4} \left(\frac{f}{M_p} \right)^2 < \delta\tilde{\beta} < 3 \times 10^{-3} \left(\frac{f}{M_p} \right)^2 \quad \text{where} \quad \delta\tilde{\beta} = 1/2 - \tilde{\beta}. \quad (28)$$

As in [13], this feeds into the amount of tuning needed in the model, which we will discuss below.

Likewise, the model has the initial condition for the start of slow roll as a function of $(f/M_p)^2$,

$$\phi_i - 1/2\pi f = (0.020-0.025) \left(\frac{f}{M_p} \right)^2 M_p. \quad (29)$$

As in all models of Goldstone inflation, the tensor to scalar ratio will also be subject to fine-tuning, but its value is generically very small:

$$r \approx 10^{-6} (f/M_p)^4. \quad (30)$$

A measurement of CMB tensor modes would fix the symmetry breaking scale f (as well as the scale of inflation, as usual) in our model.

FIG. 3. Model predictions in the n_s - r plane: Planck 2015 2σ bounds [14]. For convenience, we have set $M_p = 1$ here. In green: the TT spectrum and polarization data at low- ℓ (lowP); in pink the combined spectra TT, TE, EE + lowP.

In the CP breaking fermion implementation described above there is an additional term

$$\begin{aligned} V_{CP}(\phi) &= \tilde{c}_\eta \sin^3(\phi/f) \sqrt{1 - \sin^2(\phi/f)} \\ &= \tilde{c}_\eta \sin^3(\phi/f) \cos(\phi/f). \end{aligned} \quad (31)$$

This term imposes modulations on the potential with period πf , as seen from Fig. 2. Increasing the CP breaking in the model corresponds to increasing the value of the tensor to scalar ratio r . The bound $r < 0.1$ gives

$$\tilde{c}_\eta \leq \mathcal{O}(10^{-1}) m_\eta^2 f^2. \quad (32)$$

The effect of the CP breaking term is illustrated for an order of magnitude below this bound in Fig. 3.

The scale of inflation is related to the amplitude of the scalar power spectrum, as measured by Planck [14],

$$A_s = \frac{\Lambda^4}{24\pi^2 M_p^4 \epsilon} = \frac{e^{3.089}}{10^{10}} \quad (33)$$

where ϵ is the first slow roll parameter. For our case [Eq. (30)], where $r = 16\epsilon$ in the slow roll approximation) this implies

$$\Lambda \approx 10^{15} \left(\frac{f}{M_p} \right) \text{ GeV}. \quad (34)$$

Interestingly, we can see from this relation that the onset of inflation is related to the scale of the symmetry breaking: $\Lambda \sim 10^{-3} f$. That is, fitting to the CMB data implies a mass gap of roughly 3 orders of magnitude between the two scales.

A. Tuning

Following convention, tuning can be expressed numerically using the Barbieri-Giudice [24] parametrization as follows:

FIG. 4. Fine-tuning: numerically defined as in (35).

$$\Delta = \left| \frac{\partial \log n_s}{\partial \log \tilde{\beta}} \right| = \left| \frac{\tilde{\beta}}{n_s} \frac{\partial n_s}{\partial \tilde{\beta}} \right| \approx [8.1-8.5] \left(\frac{f}{M_p} \right)^{-2}. \quad (35)$$

See Fig. 4 below. It is seen that the parameters are sensitive to the square of the ratio of scales.

However, the relation $\tilde{\beta} \approx .5$ can be seen as a consequence of a symmetry in the sector responsible for the breaking of the global symmetry $SO(6)/SO(5)$. This would agree with naturalness in the 't Hooft interpretation. In this case the fact that the small deviation $\delta\tilde{\beta}$ is sensitive to the relation of the scales f and M_p implies that a symmetry in the sector is broken at the same time as $SO(6)/SO(5)$. In [13] we related this symmetry to the spectrum of resonances in the composite sector.

When we identify the other scalar resonance with the Higgs, we introduce a second source of tuning, between the electroweak scale v and the symmetry breaking scale f . This source of tuning coincides with the tuning in the minimal and the next to minimal composite Higgs model, and is a function of $(v/f)^2$; see for instance [2]. As this is a tuning of the parameters in the Higgs potential, which are independent combinations of the input parameters (the form factors, vacuum angles, and fermion representation), this tuning is independent and additive. The Barbieri-Giudice function will then take the form $\Delta_{\text{total}} = c_1(M_p/f)^2 + c_2(f/v)^2$, where c_1 and c_2 are $\mathcal{O}(1)$ constants. This suggests that the Barbieri-Giudice function is minimized for $\Delta_{\text{total}}(f^2 = \sqrt{4c_1/c_2} M_p v) \sim 10^{16}$, which is a large, but technically natural fine-tuning.

IV. REHEATING

At the end of inflation, the inflation field approaches, overshoots and begins to oscillate about the minimum of its potential. At this stage, the Universe is completely dominated by the zero mode of the oscillating inflaton field $\langle\phi(t)\rangle$. Interactions with the Higgs field, which we have so far neglected, lead to dissipation which drains energy from $\langle\phi(t)\rangle$, and excites relativistic Higgs particles. We refer to

these collective processes as reheating (see e.g., [18,25] for reviews). The calculation that we present in the below section is semiclassical: we treat the inflaton condensate as a classical source in the mode equations for the quantum fluctuations of the Higgs field. This treatment neglects many of the complicated processes which are present during the reheating phase, such as thermal corrections, rescatterings of the produced Higgs particles on the inflaton condensate, and the thermalization process. As we discuss at the end of this section, these effects can in general modify the rate of decay of the condensate. Our approach does however provide an estimate for the perturbative decay rate of $\langle\phi(t)\rangle$ into Higgs particles, and allows us to estimate the reheating temperature T_R .

A. Equations of motion

To begin, we study the classical inflaton background. As a first approximation, we neglect interactions with the Higgs field and set $h = 0$. As before, the inflaton sector can be canonically normalized through the field redefinition $\eta(t) = f \sin(\phi(t)/f)$. We neglect excitations of the inflaton field, $\delta\phi$, and so for simplicity label the zero mode $\phi(t) \equiv \langle\phi(t)\rangle$ which obeys the usual Klein Gordon equation:

$$\ddot{\phi} + 3H\dot{\phi} + \left. \frac{\partial V}{\partial \phi} \right|_{h=0} = 0, \quad (36)$$

where the potential is given by Eq. (27). After inflation, the inflaton field approaches, overshoots and begins to oscillate about its minimum. This region of the potential, where $\phi/f \ll 1$, is essentially quadratic:

$$V_{h=0}(\phi) \approx \frac{1}{2} m_\phi^2 \phi^2, \quad m_\phi^2 \equiv 2m_\eta^2 \approx 2 \times 10^{-14} \left(\frac{f}{M_p} \right)^2 M_p^2, \quad (37)$$

where we have used the Planck constraint on the amplitude of scalar power spectrum [Eq. (34)] to determine the mass m_ϕ in terms of the scale f . To describe the oscillations, notice that Eq. (36) can be written as

$$\frac{d^2}{dt^2} (a^{3/2} \phi) + \left[m_\phi^2 - \left(\frac{9}{4} H^2 + \frac{3}{2} \dot{H} \right) \right] (a^{3/2} \phi) = 0. \quad (38)$$

At the onset of oscillation, $m_\phi^2 \gg H^2, \dot{H}$ and under this condition, Eq. (38) has the damped sinusoidal solution:

$$\phi(t) = \frac{\Phi_0}{a^{3/2}(t)} \sin(m_\phi t + \vartheta), \quad \Phi_0 \approx 0.6 \left(\frac{f}{M_p} \right) M_p. \quad (39)$$

The numerical value for the initial amplitude, Φ_0 , was obtained by matching the above solution with an exact numerical integration of Eq. (36)—see the left-hand panel of Fig. 5 for illustration. Subscript zero denotes evaluation at the onset of oscillations (start of reheating), and we set

FIG. 5. Left: comparison between the exact numerical solution of Eq. (36) and the approximate analytic solution Eq. (39). Right: comparison between the exact “mass” [the coefficient of the term linear in χ of Eq. (44)] and $M_{\text{eff}}^2(t)$ as defined in Eq. (53).

$a_0 = 1$. The scale factor, averaged over many oscillations, grows as $a(t) \sim t^{2/3}$, while the energy density of the field decreases as

$$\rho_\phi(t) = \frac{1}{2}\dot{\phi}^2(t) + \frac{1}{2}m_\phi^2\phi^2(t) \approx \frac{m_\phi^2\Phi_0^2}{2a^3}. \quad (40)$$

We see that the vacuum energy of the inflaton field exists as spatially coherent oscillations, which can be interpreted as a condensate of nonrelativistic zero-momentum ϕ particles. The amplitude of the oscillations decay due to the Hubble expansion and also due to production of Higgs particles. We can obtain an estimate for this particle production rate by considering propagation of Higgs fluctuations, h_k , in the background of the classical inflaton condensate.

We begin by canonically normalizing the Higgs kinetic sector [given by Eq. (7)] by performing the following field redefinition:

$$\partial_\mu \chi(x) = \sqrt{\frac{f^2 - \eta^2(t)}{f^2 - \eta^2(t) - h^2(x)}} \partial_\mu h(x), \quad (41)$$

such that

$$h(x) = f \cos(\phi(t)/f) \sin \hat{\chi}(x), \quad \hat{\chi}(x) \equiv \frac{\chi(x)}{f \cos^2(\phi(t)/f)}. \quad (42)$$

We will henceforth drop the space-time labels and write $\chi = \chi(x)$, $\phi = \phi(t)$: it is to be understood that the Higgs is inhomogeneous, while the inflaton condensate is homogeneous, and described by Eq. (39). Under these field redefinitions we obtain

$$\mathcal{L} = -\frac{1}{2}\partial_\mu \chi \partial^\mu \chi - \frac{1}{2}[1 + \sin^2(\phi/f) \tan^2 \hat{\chi}] \partial_\mu \phi \partial^\mu \phi - [\sin(\phi/f) \tan \hat{\chi}] \partial_\mu \chi \partial^\mu \phi - V(\phi, \chi), \quad (43)$$

where the potential is given by Eq. (22). The canonically normalized Higgs equation of motion is obtained by varying the action with respect to χ :

$$\ddot{\chi} - \frac{\nabla^2}{a^2} \chi + 3H\dot{\chi} = -\frac{\partial V(\phi, \chi)}{\partial \chi} + \sin(\phi/f) \tan \hat{\chi} \frac{\partial V(\phi)}{\partial \phi} \Big|_{h=0} - \frac{\dot{\phi}^2}{f^2} K(\phi, \chi), \quad (44)$$

where

$$K(\phi, \chi) \equiv \frac{f \sin \hat{\chi} \cos^2 \hat{\chi} \cos^4(\phi/f) + 2\chi \cos \hat{\chi} \sin^2(\phi/f) - f \sin \hat{\chi} \cos(\phi/f) + f \sin \hat{\chi} \cos^3(\phi/f)}{\cos^3(\phi/f) \cos^3 \hat{\chi}}. \quad (45)$$

In deriving Eq. (44), we have used Eq. (36) to eliminate $\ddot{\phi}$ which arises from the variation of the action. The task at hand is to solve Eq. (44) given the inflaton background Eq. (39). This is made tractable by expanding the right-hand side of Eq. (44) about $\phi/f = 0$, and about $\chi/f = 0$:

$$\ddot{\chi} - \frac{\nabla^2}{a^2} \chi + 3H\dot{\chi} \approx -\left[m_\chi^2 + \sigma\phi + g^2\phi^2 + \frac{\dot{\phi}^2}{f^2}\right]\chi + \dots, \quad (46)$$

where we have defined

$$m_\chi^2 \equiv 2m_h^2, \quad \sigma \equiv 2c_3, \quad g^2 \equiv 2[m_h^2/f^2 - m_\eta^2/f^2 + c_4]. \quad (47)$$

The expansion in ϕ/f is permitted since the amplitude of the inflaton oscillations are small with respect to the scale

f : $\Phi_0/a^{3/2}(t) \sim 0.6f/a^{3/2}(t)$. The expansion in χ/f is permitted since we assume that the Higgs field is stabilized at the minimum of its potential throughout inflation, $\langle \chi(\mathbf{x}, t) \rangle = 0$. Furthermore we consider perturbative reheating only: we restrict ourselves to regions of parameter space where the coupling constants σ and g^2 are small enough such that resonant enhancement of Higgs modes is not possible. This ensures that $\chi \ll f$ throughout reheating. We will discuss the conditions for perturbative reheating shortly. Notice that inflaton mass, m_η^2 , and the Higgs mass, m_h^2 , enter the definition of the coupling g^2 : their presence may be traced back to canonical normalization of the Higgs kinetic term.

For the analysis of Eq. (46) it is convenient to define a comoving field

$$\mu_k(\tau) \equiv a(\tau)\chi_k(\tau), \quad (48)$$

and to work in conformal time, which is related to cosmic time by an integral over the scale factor:

$$t(\tau) = \int_{\tau_0}^{\tau} d\tau' a(\tau'). \quad (49)$$

According to standard arguments, we may decompose this field into creation and annihilation operators:

$$\mu(\tau, \mathbf{x}) = \int \frac{d^3k}{(2\pi)^{3/2}} [a_{\mathbf{k}}\mu_k(\tau) + a_{-\mathbf{k}}^\dagger\mu_k^*(\tau)]e^{i\mathbf{k}\cdot\mathbf{x}}, \quad (50)$$

where the mode functions obey

$$\mu_k''(\tau) + \omega_k^2(\tau)\mu_k(\tau) = 0, \quad (51)$$

and where a prime denotes differentiation with respect to conformal time. The time dependent frequency is given by

$$\omega_k^2(\tau) \equiv k^2 + a^2 M_{\text{eff}}^2(\tau) - \frac{a''}{a}, \quad \frac{a''}{a} = \frac{a^2}{6M_{\text{p}}^2}(\rho_\phi - 3P_\phi), \quad (52)$$

where $P_\phi \simeq 0$ is the pressure of the field, and we have defined the effective mass:

$$M_{\text{eff}}^2(t) \equiv m_\chi^2 + \frac{\sigma\Phi_0}{a^{3/2}(t)}\sin(m_\phi t + \vartheta) + \frac{g^2\Phi_0^2}{a^3(t)}\sin^2(m_\phi t + \vartheta) + \frac{\Phi_0^2 m_\phi^2}{f^2 a^3(t)}\cos^2(m_\phi t + \vartheta). \quad (53)$$

The final term on the right-hand side of $M_{\text{eff}}^2(t)$ is the leading contribution from $\dot{\phi}^2/f^2$: we have neglected terms which decay faster than a^{-3} . In the right panel of Fig. 5, we plot the effective mass against the coefficient of the term linear in χ of Eq. (44), which demonstrates the accuracy of this expansion. Equations of the type (51), with time dependent mass (53) have been extensively studied in the context of (p)reheating after inflation. For certain regions of $\{\sigma, g^2, \Phi_0\}$ parameter space, the mode functions experience exponential growth as parametric instability develops, a phenomenon known as parametric resonance [18,20,26,27]. To be specific, when any one of the three terms in $M_{\text{eff}}^2(t)$ is dominant, the oscillator equation (51) may be written

$$\frac{d^2\mu_k}{dz^2} + [A_k - 2q_i \cos(2z)]\mu_k = 0, \quad (54)$$

$$q_0 \equiv \frac{\Phi_0^2}{4f^2 a^3}, \quad q_3 \equiv \frac{\sigma\Phi_0}{m_\phi^2 a^{3/2}}, \quad q_4 \equiv \frac{g^2\Phi_0^2}{4m_\phi^2 a^3}, \quad A_k \equiv \frac{k^2 + m_\chi^2}{m_\phi^2 a^2} + 2q_{(0,4)}, \quad (55)$$

following a time redefinition of the form $z \equiv m_\phi t + \text{const.}$ Here we have ignored terms proportional to H/m_ϕ (recall that $H \ll m_\phi$ during reheating). Equation (54) is known as the Mathieu equation, which is known to possess instability bands for certain values of A_k and q_i . For $q_i \gg 1$, a large region of parameter space is unstable and broad parametric resonance can develop. Throughout this paper we restrict ourselves to regions of parameter space where $q_i \ll 1$, such that nonperturbative preheating processes are negligible. With $\Phi_0 \approx 0.6f$, we find $q_0 = 0.09$, and so parametric instability cannot be triggered by this term. Meanwhile, $q_{3,4} \ll 1$ requires

$$\sigma \ll \frac{m_\phi^2}{\Phi_0}, \quad g^2 \ll \left(\frac{m_\phi}{\Phi_0}\right)^2, \quad (56)$$

or, in terms of the original parameters of the potential (22)

$$c_3 \ll m_\eta^2/f, \quad m_\eta^2/f^2 + c_4 \ll 10m_\eta^2/f^2. \quad (57)$$

This relation for the smallness of the CP breaking term c_3 in terms of the inflaton mass is consistent with the similar relation for c_η found in the previous section. Likewise, the constraint on c_4 is consistent with our expectations from the computation of the potential, as can be verified with the appendix. We always ensure that the above bounds are respected, and do not consider parametric resonance in this paper.

If we regard the inflaton condensate ϕ to be a collection of zero-momentum inflaton “particles,” then the effective mass $M_{\text{eff}}^2(t)$ has a physical interpretation in terms of Feynman diagrams:

These diagrams describe the three-leg, $-\frac{1}{2}\sigma\phi\chi^2$, and four-leg, $-\frac{1}{2}g^2\phi^2\chi^2$, interaction terms which reside in the canonically normalized Lagrangian—Eq. (43). Since we have not quantized the inflaton, there are no ϕ propagators, which allows for tree-level diagrams only. These diagrams describe the perturbative decay of a single inflaton particle with mass m_ϕ into two Higgs particles of comoving momentum $k \sim am_\phi/2$, and the annihilation of a pair of ϕ particles into a pair of χ particles with comoving momentum $k \sim am_\phi$ respectively. We use the term inflaton particle rather loosely here, since what we are really describing is creation of Higgs particles from a classical

inflaton condensate. This diagrammatic representation does however offer intuition for the physical processes at work.

B. Bogoliubov calculation

We wish to solve Eq. (51) with frequency (52). Our calculation closely follows that of Ref. [20]. First, we notice that since the inflaton condensate behaves like a collection of nonrelativistic particles with zero pressure, $P_\phi \approx 0$, and so we have $a''/a \approx 2a^2H^2$. Therefore, for the modes $k^2 \sim a^2m_\phi^2$ which we expect to be produced, we can safely neglect a''/a , given that $H \ll m_\phi$ during reheating. In the adiabatic representation, the solution to the mode equation Eq. (51) may be written in the WKB form (see e.g. [18,20]):

$$\mu_k(\tau) = \frac{\alpha_k(\tau)}{\sqrt{2\omega_k(\tau)}} e^{-i\Psi_k(\tau)} + \frac{\beta_k(\tau)}{\sqrt{2\omega_k(\tau)}} e^{+i\Psi_k(\tau)}, \quad (58)$$

where the accumulated phase is given by

$$\Psi_k(\tau') \equiv \int_{\tau_0}^{\tau'} d\tau'' \omega_k(\tau''). \quad (59)$$

Equation (58) is a solution of Eq. (51) provided that the Bogoliubov coefficients satisfy the following coupled equations:

$$\begin{aligned} \alpha'_k(\tau) &= \beta_k(\tau) \frac{w'_k(\tau)}{2w_k(\tau)} e^{+2i\Psi_k(\tau)}, \\ \beta'_k(\tau) &= \alpha_k(\tau) \frac{w'_k(\tau)}{2w_k(\tau)} e^{-2i\Psi_k(\tau)}, \end{aligned} \quad (60)$$

which also implies that

$$\mu'_k(\tau) = -i\alpha_k(\tau) \sqrt{\frac{w_k(\tau)}{2}} e^{-i\Psi_k(\tau)} + i\beta_k(\tau) \sqrt{\frac{w_k(\tau)}{2}} e^{+i\Psi_k(\tau)}. \quad (61)$$

The Wronskian condition, $W[\mu_k(t), \mu_k^*(t)] = i$, demands that the Bogoliubov coefficients are normalized as $|\alpha_k(t)|^2 - |\beta_k(t)|^2 = 1$. In this basis, the Hamiltonian of the χ field is instantaneously diagonalized. The single particle mode occupation number n_k is defined as the energy of the mode, $\frac{1}{2}|\mu'_k|^2 + \frac{1}{2}\omega_k^2|\mu_k|^2$, divided by the frequency of the mode:

$$n_k(\tau) = \frac{|\mu'_k(\tau)|^2 + \omega_k^2(\tau)|\mu_k(\tau)|^2}{2\omega_k(\tau)} - \frac{1}{2} = |\beta_k(\tau)|^2. \quad (62)$$

The $-1/2$ corresponds to subtraction of the zero-point energy, and the last equality is obtained via substitution of the WKB solution (58). In terms of the classical mode functions, creation of Higgs particles occurs due to departure from the initial positive-frequency solution: the initial conditions therefore at $\tau = \tau_0$ (the start of reheating) are

then $\alpha_k = 1$, $\beta_k = 0$, and so $n_k(\tau_0) = 0$. Since we work in the perturbative regime specified by Eq. (56) the mode occupation numbers remain small, $|\beta_k(\tau)|^2 \ll 1$, and so we can iterate Eq. (60) to obtain

$$\beta_k(\tau) \approx \int_{\tau_0}^{\tau} d\tau' \frac{\omega'_k(\tau')}{2\omega_k(\tau')} e^{-2i\Psi_k(\tau')}. \quad (63)$$

In the perturbative regime we can approximate

$$\Psi_k(\tau') \approx k \int_{\tau_0}^{\tau'} d\tau'' \sqrt{1 + \left(\frac{a(\tau'')m_\chi}{k} \right)^2}, \quad (64)$$

while for the frequency we have

$$\begin{aligned} \frac{\omega'_k}{2\omega_k} &\approx \frac{a^{3/2}(\tau')\Phi_0 m_\phi}{4k^2} \\ &\times \left[\frac{\sigma + 2\Phi_0(g^2 - m_\phi^2/f^2)a^{-3/2}(\tau') \sin(m_\phi t(\tau') + \vartheta)}{1 + a^2(\tau')m_\chi^2/k^2} \right] \\ &\times \cos(m_\phi t(\tau') + \vartheta), \end{aligned} \quad (65)$$

where we have neglected terms containing derivatives of the scale factor. Inserting these results into Eq. (63) gives

$$\begin{aligned} \beta_k(\tau) &= \frac{\sigma\Phi_0 m_\phi}{8k^2} \int_{\tau_0}^{\tau} \frac{d\tau' a^{3/2}(\tau')}{1 + a^2(\tau')m_\chi^2/k^2} [e^{+i\psi_{3,k}^-(\tau')} + e^{-i\psi_{3,k}^+(\tau')}] \\ &+ \frac{(g^2 - m_\phi^2/f^2)\Phi_0^2 m_\phi}{8ik^2} \\ &\times \int_{\tau_0}^{\tau} \frac{d\tau'}{1 + a^2(\tau')m_\chi^2/k^2} [e^{+i\psi_{4,k}^-(\tau')} - e^{-i\psi_{4,k}^+(\tau')}], \end{aligned} \quad (66)$$

where we have defined the phases

$$\begin{aligned} \psi_{3,k}^\pm(\tau) &\equiv \pm 2\Psi_k(\tau) + m_\phi t(\tau) + \vartheta, \\ \psi_{4,k}^\pm(\tau) &\equiv \pm 2\Psi_k(\tau) + 2(m_\phi t(\tau) + \vartheta). \end{aligned} \quad (67)$$

As discussed in Ref. [20] (see also [18]), the integrals in Eq. (66) can be evaluated using the method of stationary phase: they are dominated near the instants $\tau_{3,k}$ and $\tau_{4,k}$ where

$$\begin{aligned} \left. \frac{d}{d\tau} \psi_{3,k}^-(\tau) \right|_{\tau_{3,k}} &= 0, \Rightarrow k = \frac{1}{2} m_\phi a(\tau_{3,k}) \sqrt{1 - 4\delta_M^2}, \\ \left. \frac{d}{d\tau} \psi_{4,k}^-(\tau) \right|_{\tau_{4,k}} &= 0, \Rightarrow k = m_\phi a(\tau_{4,k}) \sqrt{1 - \delta_M^2}, \end{aligned} \quad (68)$$

where we have defined $\delta_M \equiv m_\chi/m_\phi$. For the 3-leg interaction, the above result corresponds to the creation of a pair of Higgs particles with momentum $k \sim am_\phi/2$

from an inflaton with mass m_ϕ at the instant $\tau_{3,k}$ of the resonance between the mode k and the inflaton condensate. A similar interpretation may be given for the 4-leg interaction. Upon performing the integrals, we find

$$n_k(\tau) = \frac{\pi\sigma^2\Phi_0^2 m_\phi}{32k^4} (1 - 4\delta_M^2) \frac{a^3(\tau_{3,k})}{a'(\tau_{3,k})} + \frac{\pi(g^2 - m_\phi^2/f^2)^4 \Phi_0^2 m_\phi}{64k^4} \frac{(1 - \delta_M^2)}{a'(\tau_{4,k})} \\ + \frac{\pi\sigma(g^2 - m_\phi^2/f^2) \Phi_0^3 m_\phi}{32k^4} \sqrt{2(1 - 4\delta_M^2)(1 - \delta_M^2)} \mathcal{I}(\tau_{3,k}\tau_{4,k}), \quad (69)$$

where we have defined

$$\mathcal{I}(\tau_{3,k}\tau_{4,k}) \equiv \sqrt{\frac{a^3(\tau_{3,k})}{a'(\tau_{3,k})a'(\tau_{4,k})}} \sin[\psi_{4,k}^-(\tau_{4,k}) - \psi_{3,k}^-(\tau_{3,k})]. \quad (70)$$

As discussed in [20], the oscillatory term $\mathcal{I}(\tau_{3,k}\tau_{4,k})$ represents the interference between the two decay channels ($\phi \rightarrow \chi\chi$ and $\phi\phi \rightarrow \chi\chi$) of the inflaton. It is present because we have treated the inflaton as a classical oscillating source, and not an honest collection of particles.

C. Boltzmann equations

Since $m_\phi \gg m_\chi$ the Higgs particles are relativistic when produced. This means we can effectively treat them as a bath of radiation with g_* number of degrees of freedom. We define the comoving energy density in the Higgs field as

$$a^4 \rho_\chi \equiv \int_0^\infty \frac{d^3k}{(2\pi)^3} \omega_k n_k \\ = \frac{\sigma^2 \Phi_0^2 m_\phi}{64\pi} (1 - 4\delta_M^2) \int_0^\infty \frac{dk}{k^2} \sqrt{k^2 + a^2(\tau) m_\chi^2} \frac{a^3(\tau_{3,k})}{a'(\tau_{3,k})} \\ + \frac{(g^2 - m_\phi^2/f^2)^2 \Phi_0^4 m_\phi}{128\pi} (1 - \delta_M^2) \int_0^\infty \frac{dk}{k^2} \sqrt{k^2 + a^2(\tau) m_\chi^2} \frac{1}{a'(\tau_{4,k})} \\ + \frac{\sigma(g^2 - m_\phi^2/f^2) \Phi_0^3 m_\phi}{64\pi} \sqrt{2(1 - 4\delta_M^2)(1 - \delta_M^2)} \int_0^\infty \frac{dk}{k^2} \sqrt{k^2 + a^2(\tau) m_\chi^2} \mathcal{I}(\tau_{3,k}\tau_{4,k}). \quad (71)$$

At first glance these integrals appear divergent. This however is not the case, as can be seen from the requirement that the Higgs particles be produced perturbatively. Equation (68) enforces

$$\frac{1}{2} m_\phi a_0 \sqrt{1 - 4\delta_M^2} < k < \frac{1}{2} m_\phi a(\tau) \sqrt{1 - 4\delta_M^2}, \quad \text{for } \phi \rightarrow \chi\chi, \quad (72)$$

$$m_\phi a_0 \sqrt{1 - \delta_M^2} < k < m_\phi a(\tau) \sqrt{1 - \delta_M^2}, \quad \text{for } \phi\phi \rightarrow \chi\chi. \quad (73)$$

Hence, the limits of the first and the third integrals on the right-hand side of Eq. (71) should be replaced by the limits of Eq. (72), while those of the second integral should be replaced by Eq. (73). Once again neglecting derivatives of a , we obtain

$$\frac{d}{d\tau} (a^4 \rho_\chi) \approx a^2 \frac{\sigma^2 \Phi_0^2 m_\phi}{64\pi} \sqrt{1 - 4\delta_M^2} \\ + a^{-1} \frac{(g^2 - m_\phi^2/f^2)^2 \Phi_0^4 m_\phi}{128\pi} \sqrt{1 - \delta_M^2}, \quad (74)$$

where we have discarded the interference term since it vanishes when averaged over time. Replacing factors of a using $\rho_\phi \approx m_\phi^2 \Phi_0^2 / (2a^3)$, we are left with the familiar Boltzmann equation:

$$a^{-4} \frac{d}{dt} (a^4 \rho_\chi) \approx \Gamma_{\phi \rightarrow \chi\chi} \rho_\phi + 2 \frac{[\sigma_{\phi\phi \rightarrow \chi\chi} v]_{v=0}}{m_\phi} \rho_\phi^2, \quad (75)$$

where

$$\Gamma_{\phi \rightarrow \chi\chi} = \frac{\sigma^2}{32\pi m_\phi} \sqrt{1 - 4 \frac{m_\chi^2}{m_\phi^2}}, \\ [\sigma_{\phi\phi \rightarrow \chi\chi} v]_{v=0} = \frac{(g^2 - m_\phi^2/f^2)^2}{64\pi m_\phi^2} \sqrt{1 - \frac{m_\chi^2}{m_\phi^2}}. \quad (76)$$

The decay rate $\Gamma_{\phi \rightarrow \chi\chi}$ agrees with the tree-level result obtained from Quantum Field Theory (QFT). The cross section $\sigma_{\phi\phi \rightarrow \chi\chi}$ also agrees with QFT so long as the Feynman amplitude is evaluated at zero relative velocity, $v = 0$.

Note that ϕ , as a CP odd particle, could have couplings to vector bosons as an axion. For example, it could have couplings to gluons and photons as

$$\mathcal{L}_{CP} = \frac{c_\gamma \alpha}{f} \phi F_{\mu\nu} \tilde{F}^{\mu\nu} + \frac{c_s \alpha_s}{f} \phi \text{Tr} G_{\mu\nu} \tilde{G}^{\mu\nu} \quad (77)$$

as well as to W and Z bosons. These couplings could be generated by triangle diagrams involving fermionic degrees of freedom coupled to SM gauge interactions. Whether these are present or not is a highly model dependent question, whereas we have focused in this paper on interactions between the Goldstone bosons (the Higgs and the inflaton). We refer the reader to Refs. [28,29] for a thorough analysis of preheating due to nonzero couplings to gauge bosons.

Conservation of energy demands $a^{-3} \frac{d}{dt} (a^3 \rho_\phi) = -a^{-4} \frac{d}{dt} (a^4 \rho_\chi)$, which gives

$$\frac{d}{dt} (a^3 \rho_\phi) = -\Gamma_{\phi \rightarrow \chi\chi} (a^3 \rho_\phi) - 2 \frac{[\sigma \phi \phi \rightarrow \chi\chi v]_{v=0}}{m_\phi a^3} (a^3 \rho_\phi)^2. \quad (78)$$

If the trilinear interaction is absent ($\sigma = 0$) we can integrate Eq. (78) to show that $a^3 \rho_\phi \rightarrow \text{const}$ as $t \rightarrow \infty$. This means that the inflaton does not completely decay: volume dilution due to the Hubble expansion takes place faster than the annihilation process $\phi\phi \rightarrow \chi\chi$ can drain energy from the inflaton condensate. In order to successfully reheat the Universe, the trilinear coupling must be present. Indeed, in the absence of $\phi\phi \rightarrow \chi\chi$ annihilations (if $g^2 = m_\phi^2/f^2$), we can integrate Eq. (78) to show that $a^3 \rho_\phi \sim e^{-\Gamma t}$: in a time of order $\Gamma_{\phi \rightarrow \chi\chi}^{-1}$ the inflaton has decayed completely. For the remainder of this section we set $g^2 = m_\phi^2/f^2$ in order to place order-of-magnitude bounds on the model parameters.

Up to this point we have neglected the decay of the Higgs to the SM. The dominant channel is $\chi \rightarrow b\bar{b}$, with width

$$\Gamma_{\chi \rightarrow b\bar{b}} = \frac{3m_\chi}{8\pi} \left(\frac{m_b}{v_\chi} \right)^2 \left(1 - \frac{4m_b^2}{m_\chi^2} \right)^{3/2} \sim 5 \text{ MeV}. \quad (79)$$

Since $m_\chi \gg m_b$, the $b\bar{b}$ decay products are produced relativistically:

$$a^{-4} \frac{d}{dt} (a^4 \rho_b) = \Gamma_{\chi \rightarrow b\bar{b}} \rho_b. \quad (80)$$

With $\phi\phi \rightarrow \chi\chi$ processes absent, energy conservation demands

$$\begin{aligned} a^{-4} \frac{d}{dt} (a^4 \rho_\chi) &\approx \Gamma_{\phi \rightarrow \chi\chi} \rho_\phi - \Gamma_{\chi \rightarrow b\bar{b}} \rho_b, \\ a^{-3} \frac{d}{dt} (a^3 \rho_\phi) &= -\Gamma_{\phi \rightarrow \chi\chi} \rho_\phi. \end{aligned} \quad (81)$$

Equations (80) and (81) are the final Boltzmann equations describing perturbative reheating in the composite

Higgs model. The approximations involved in their derivation will begin to break down when the energy density of the decay products becomes comparable to the energy density of the inflaton condensate. Furthermore, as pointed out in [30], and discussed in detail in [31,32], $\Gamma_{\phi \rightarrow \chi\chi}$ develops a temperature dependence due to interactions (which we have not accounted for) between the decay products and the condensate. Indeed, as the decay products thermalize via scatterings and further decays, they acquire a temperature dependent “plasma” mass $m_p(T)$ of the order $\sim \lambda T^2$, where λ is a typical coupling constant for a particle in the plasma. The presence of these “thermal” masses prevent decay of the condensate if $m_\phi^2 \approx \lambda T^2$: the decay process becomes kinematically forbidden. An important consequence of these finite temperature corrections is that the reheating temperature, T_R (the temperature at the onset of the radiation dominated phase) is generally higher compared to the naive estimate obtained via setting $\Gamma = H$ (see the following section).

In addition to the effect of thermal masses, the produced χ particles can “rescatter” off the oscillating condensate $\langle \phi \rangle$ to excite $\delta\phi$ particles. This opens another possible channel for decay of the condensate. We illustrate this schematically in Fig. 6 for the case of the 4-leg interaction. In the language of our Bogoliubov calculation, this process corresponds to the term $\chi^2 \phi \delta\phi$ which results from expanding ϕ about the mean field: $\phi(x) = \phi(t) + \delta\phi(x)$. There is also a subdominant process of the type $\chi\chi \rightarrow \delta\phi \delta\phi$, which is phase space suppressed. Such processes, which we have neglected in this work, will promote the decay rate $\Gamma_{\phi \rightarrow \chi\chi}$ from a constant to a function of time and temperature. To include these processes would require recourse to nonequilibrium thermal field theory, which is beyond the scope of this paper. Having acknowledged these caveats,

FIG. 6. Schematic illustration of possible inflaton-Higgs interactions. The vacuum energy of the inflaton field exists as spatially coherent oscillations, which can be interpreted as a condensate of nonrelativistic zero-momentum ϕ particles. The condensate decays via three-leg, $-\frac{1}{2}\sigma\phi\chi^2$, and four-leg, $-\frac{1}{2}g^2\phi^2\chi^2$, interactions. The Bogoliubov calculation presented in Sec. IV B treats the condensate as a classical source, and so rescattering processes between the produced Higgs particles and the condensate which excite $\delta\phi$ particles are ignored.

we use the Boltzmann equations (80) and (81) to place rough bounds on our model parameters only.

D. Parameter constraints from reheating

Combining the Planck constraint on the inflaton mass, Eq. (37), with the bound (56), we find that for reheating to proceed perturbatively,

$$\left(\frac{\sigma}{M_p}\right)^2 \ll 10^{-27} \left(\frac{f}{M_p}\right)^2, \quad (82)$$

where we have used $\Phi_0 \sim 0.6f$. This provides an upper bound on the trilinear coupling σ in terms of the scale f . A lower bound on σ can be obtained from the condition that the Universe be totally radiation dominated before the Big Bang Nucleosynthesis (BBN) epoch. This requires knowledge of the reheating temperature T_R , which may be estimated as follows: Reheating completes at time t_c , when the Hubble rate $H^2 = \rho/3M_p^2 \sim t_c^{-2}$ drops below the decay rate $\Gamma_{\phi \rightarrow \chi\chi}$. The density of the Universe at this moment is then

$$\rho(t_c) \simeq 3M_p^2 H^2(t_c) = 3M_p^2 \Gamma_{\phi \rightarrow \chi\chi}^2. \quad (83)$$

Provided that the Higgs particles are produced in thermal and chemical equilibrium, the temperature of the Higgs plasma is T_R . Treating this ultrarelativistic gas of particles with Bose-Einstein statistics, the energy density of the Universe in thermal equilibrium is then

$$\rho(T_R) \simeq \left(\frac{\pi^2}{30}\right) g_* T_R^4, \quad (84)$$

where the factor $g_*(T_R) \sim 10^2$ – 10^3 depends on the number of ultrarelativistic degrees of freedom. Comparing Eqs. (83) and (84) we arrive at

$$T_R \approx 0.1 \sqrt{\Gamma_{\phi \rightarrow \chi\chi} M_p}. \quad (85)$$

In order not to spoil the success of BBN, the Universe must be completely dominated by relativistic particles before the BBN epoch. This constrains the reheating temperature to be $T_R \gtrsim 5$ MeV [33,34], which in turn implies⁷

$$\Gamma_{\phi \rightarrow \chi\chi} \gtrsim 10^{-40} M_p. \quad (86)$$

Combining Eqs. (37), (76), (86) we find

$$\left(\frac{\sigma}{M_p}\right)^2 \gtrsim 10^{-45} \left(\frac{f}{M_p}\right)^2. \quad (87)$$

Finally, combining this temperature bound with the bound for perturbative reheating Eq. (82), we find

⁷We note that since T_R also enters expressions for the primordial observables, the lower bound on $\Gamma_{\phi \rightarrow \chi\chi}$ given by Eq. (86) may be tightened if our model were to be confronted with CMB data—see for example Ref. [35].

$$f \gg 10^{-18} M_p. \quad (88)$$

V. TEV INFLATON AND ITS CONSEQUENCES

With the inflaton and Higgs doublet originated by the breaking of the same global symmetry, the Coleman-Weinberg contributions to their potential are naturally of the same order. Therefore, we would expect the mass of both particles to be not far from each other, $m_\eta \sim m_h$, as well as similar size couplings. From perturbative reheating we require $m_\eta > 2m_h$ as well as a condition on the cubic coupling Eq. (57), namely

$$\frac{c_3}{f} \ll \left(\frac{m_\eta}{f}\right)^2, \quad (89)$$

which is technically natural as the parameter c_3 breaks the symmetry $\eta \rightarrow -\eta$.

Inflation would also impose a bound on the mass of the inflaton with respect to the scale of breaking, see Eqs. (34) and (26), $m_\eta/f \simeq 10^{-6}$, a hierarchy which is again technically natural. On the other hand, in our inflationary potential we could have added a constant term, a *phenomenological* cosmological constant which could change this condition and allow closer values of f and m_η .

One should also keep in mind that inflation cannot last to reach energies around the MeV when the very predictive theory of big-bang nucleosynthesis takes on [36]. Another constraint to keep in mind is the generation of baryon asymmetry in the Universe, which in the context of electroweak baryogenesis (see Ref. [37] and references therein) would require inflation to end some time before the electroweak scale. One additional attractive feature of this model is that the conditions for reheating, which in turn require CP violation, could be helpful for baryogenesis, e.g. see Ref. [38] for a study of electroweak baryogenesis in a similar model.

If the inflaton is heavier than the Higgs doublet, one can integrate it out leading to an effective field theory (EFT). In Ref. [39] one can find a more general discussion on the EFT due the presence of a singlet like η , and its phenomenology.

Interestingly, the cubic term c_3 is the main player in the reheating discussion as well as the collider phenomenology. The cubic term, when the Higgs acquires a vacuum expectation value v , would lead to a mixing of the singlet with the Higgs, resulting in two mass eigenstates with an admixture of η and h . The mixing angle is given by

$$s_\theta \simeq \frac{c_3 v}{m_\eta^2}. \quad (90)$$

The mixing, then, changes the way the physical SM-like Higgs behaves, as well as induces new couplings of the heavy η -like state to vector bosons and fermions. Detailed studies from electroweak precision tests (EWPT) at LEP, as well as current constraints from the measurement of the

FIG. 7. Present and future 95% C.L. exclusion limits in the (m_η, c_3) plane from ATLAS and CMS measurements of Higgs signal strengths (denoted by Run1 indirect) and from EWPT (denoted by LEP). Values above the red-dashed line are excluded at 95% C.L. by the combination (EWPT and Higgs signal strength). Above the green line may also be excluded by constraints from heavy scalar searches at LHC, although these limits could be evaded in the presence of new decay modes for η . Also shown is the projected exclusion reach from Higgs signal strengths at the 14 TeV run of LHC with $\mathcal{L} = 300 \text{ fb}^{-1}$ and at HL-LHC with $\mathcal{L} = 3000 \text{ fb}^{-1}$ in blue, as well as projections from measurements of the S and T oblique parameters with ILC-*GigaZ* and FCC-ee in dashed-blue.

Higgs properties imposes strong bounds on this mixing. Moreover, the heavier state can be searched for directly and the reach for these searches is related to the amount of mixing.

In Fig. 7, we show current and future constraints on these parameters. They include (1) a χ^2 fit to Higgs coupling measurements [40–49], (2) the 95% C.L. exclusion prospects for LHC at 14 TeV with $\mathcal{L} = 300 \text{ fb}^{-1}$ and $\mathcal{L} = 3000 \text{ fb}^{-1}$, by assuming that future measurements of Higgs signal strengths will be centered at the SM value, and use the projected CMS sensitivities, (3) a fit to the oblique parameters S , T , U using the best-fit values and standard deviations from the global analysis of the GFitter Group [50], and finally (4) future limits on EW precision observables from e^+e^- colliders (see e.g. [51]), ILC and FCC-ee. The corrections to S and T from the inflaton-Higgs mixing given by

$$\begin{aligned} \Delta S &= \frac{1}{\pi} s_\theta^2 \left[-H_S \left(\frac{m_h^2}{m_Z^2} \right) + H_S \left(\frac{m_\eta^2}{m_Z^2} \right) \right], \\ \Delta T &= \frac{g^2}{16\pi^2 c_W^2 \alpha_{\text{EM}}} s_\theta^2 \left[-H_T \left(\frac{m_h^2}{m_Z^2} \right) + H_T \left(\frac{m_\eta^2}{m_Z^2} \right) \right] \end{aligned} \quad (91)$$

with the functions $H_S(x)$ and $H_T(x)$ defined in Appendix C of [52].

Regarding future colliders, we assumed a SM best-fit value, and interpreted the ILC *GigaZ* program’s expected precision is $\sigma_S = 0.017$ and $\sigma_T = 0.022$ [50,53] and the FCC-ee prospects of $\sigma_S = 0.007$ and $\sigma_T = 0.004$ [54]. As one can see, colliders are sensitive to relatively large values of the triple coupling, whereas perturbative reheating is sensitive to lower values of the coupling.

Finally, note that in the explicit CP breaking scenario, there would be direct couplings of the inflaton to SM fermions (ϵ_f) and these would be proportional to c_3 ; see Eq. (17b).

VI. CONCLUSIONS

We have presented a single model that can realize inflation, perturbative reheating, and electroweak symmetry breaking in a natural way. In the minimal model the five Goldstone bosons from the global symmetry breaking $SO(6) \sim SU(4) \rightarrow SO(5) \sim Sp(4)$ play the role of a Higgs doublet and an inflaton singlet. We have argued that a trilinear coupling between the latter (η) and two Higgs bosons (h) is necessary for successful reheating, and shown under which condition this term can be present. In particular, the model needs to have broken CP , which can be realized spontaneously or explicitly. A detailed derivation of the scalar potential for h and η arising from loops of $SU(2)$ gauge bosons and fermions in the **6** of $SU(4)$ was given in the first section.

The CMB results [14] allow us to relate the parameters in our model, and explain mass hierarchies. A range of energy scales for inflation, or equivalently for the mass of the inflaton, was presented in the second section. To the merit of the model, none of the relevant scales are expected to be affected by quantum gravity.

The motive of perturbative reheating further fixes the parameters in the potential. For a particular range of parameter space [given by Eq. (57)] parametric instability is not triggered and nonperturbative effects are subdominant. With a Bogoliubov calculation [20] we find the single particle occupation numbers, and as usual the evolution of the fields is established using Boltzmann equations. We finished this section by an exposition of the numerical constraints on the reheating temperature and the model parameters from perturbativity (86)–(88).

We have also explored the possibility of TeV values of the inflaton mass and coupling to the Higgs. As an effective theory, the inflaton’s effect at low energies is inducing a mixing effect in the Higgs particle properties, an effect which is constrained by precise electroweak data as well as the LHC. We discussed the future reach for colliders on the inflaton-Higgs parameter space, finding that while perturbative reheating explores a region of small mixing, colliders are most sensitive to large values of this parameter.

The model building presented in this paper hints at interesting opportunities for further studies. The fact that the model is able to address and connect normally unrelated

cosmological events in a natural way shows that the considerations here may indeed tempt the reader to further inquiry, in the light of recent developments. As mentioned in the Introduction, the discussion of cosmological relaxation by an interplay between the Higgs and a pGB [15] offers an attractive example. Other directions include an investigation of the changed evolution of the Higgs dynamics and its implications on electroweak stability [17], possible UV completions for which the present theory is a boundary condition at low energy (on which we commented in [13]), as well as the implications of CP violation and the inflaton degree of freedom for electroweak baryogenesis.

ACKNOWLEDGMENTS

We would like to thank Marco Drewes, Hitoshi Murayama, and Josemi No for discussions. This work is supported by the Science Technology and Facilities Council (STFC) under Grant No. ST/L000504/1.

APPENDIX: COMPUTATION OF THE SCALAR POTENTIAL

1. Composite Higgs vacuum

At one loop, the Coleman-Weinberg potential due to up-type quarks coupling to Σ as in (11) is given by⁸

$$V(h, \eta) = -2N_c \int \frac{d^4 p}{(2\pi)^4} \log(p^2 \Pi_{u_L} \Pi_{u_R} - |\Pi_{u_L u_R}|^2) \quad (\text{A1})$$

where we have used new form factors for simplicity, which are just rotations of the original parameters in the Lagrangian (11):

$$\Pi_{u_L} = \frac{\Pi_0^q + \Pi_0^{q'}}{2} - \Pi_1^q \frac{\text{Tr}[\bar{\Psi}_q \Sigma] \text{Tr}[\Psi_q \Sigma^\dagger]}{\bar{u}_L \not{p} u_L}, \quad (\text{A2a})$$

$$\Pi_{u_R} = \Pi_0^u - \Pi_1^u \frac{\text{Tr}[\bar{\Psi}_u \Sigma] \text{Tr}[\Psi_u \Sigma^\dagger]}{\bar{u}_R \not{p} u_R}, \quad (\text{A2b})$$

$$\Pi_{u_L u_R} = M_1^u \frac{\text{Tr}[\bar{\Psi}_q \Sigma] \text{Tr}[\Psi_u \Sigma^\dagger]}{\bar{u}_L u_R}. \quad (\text{A2c})$$

As explained in the main text, we refer to Ψ as the fermion multiplets in the 6 of $SU(4)$.

⁸In general there will be contributions from down type quarks and gauge bosons as well. In fact, it should be noted that at least one other fermion generation is needed to make the CP assignment physical [55]. However, these will not lead to different couplings in the scalar potential, and here we take them to be subleading corrections to the coefficients.

If we assume the ratios form factors fall off rapidly enough with momentum to make the integrals converge, we may expand the logarithms to find the following Lagrangian to fourth order in the fields⁹:

$$V(\phi, h) = a_1 h^2 + \lambda h^4 + |\kappa|^2 (a_2 + a_3 h^2 + a_4 |\kappa|^2) \quad (\text{A3})$$

where $\kappa = \sqrt{f^2 - h^2 - \eta^2} + i\epsilon_1 \eta$. The coefficients are given by integrals over the form factors of the fields contributing to the Coleman-Weinberg potential: the gauge bosons, and the up-type and down-type fermions. If we assume the contributions are dominated by the heaviest up-type quark, which we will call the top as in the Standard Model (while this quark is not necessarily identified with the Standard Model top), the coefficients are given by

$$a_1 = -2f^2 N_c \int \frac{d^4 p}{(2\pi)^4} \frac{1}{\Pi_0} (-4\Pi_1^q \Pi_0^t), \quad (\text{A4a})$$

$$a_2 = -2f^2 N_c \int \frac{d^4 p}{(2\pi)^4} \frac{1}{\Pi_0} (-2\Pi_0^q \Pi_1^t - 2\Pi_1^{q'} \Pi_1^t), \quad (\text{A4b})$$

$$a_3 = -2N_c \int \frac{d^4 p}{(2\pi)^4} \frac{1}{\Pi_0} \left(-\frac{4|M_t^1|^2}{p^2} + \frac{8\Pi_0^q \Pi_1^q \Pi_0^t \Pi_1^t}{\Pi_0} + \frac{8\Pi_1^q \Pi_1^{q'} \Pi_0^t \Pi_1^t}{\Pi_0} + 16\Pi_1^q \Pi_1^t \right), \quad (\text{A4c})$$

$$a_4 = -2N_c \int \frac{d^4 p}{(2\pi)^4} \frac{1}{\Pi_0} \left(\frac{2(\Pi_0^q)^2 (\Pi_1^t)^2}{\Pi_0} + \frac{4\Pi_0^q \Pi_1^{q'} (\Pi_1^t)^2}{\Pi_0} + \frac{2(\Pi_1^{q'})^2 (\Pi_1^t)^2}{\Pi_0} \right), \quad (\text{A4d})$$

$$\lambda = -2N_c \int \frac{d^4 p}{(2\pi)^4} \frac{1}{\Pi_0} \left(\frac{8(\Pi_1^q)^2 (\Pi_0^t)^2}{\Pi_0} \right), \quad (\text{A4e})$$

where Π_0 is the relevant field independent factor:

$$\Pi_0 = \frac{1}{2} \Pi_0^t (\Pi_0^q + \Pi_0^{q'}) \quad (\text{A5})$$

i.e., a function of the different propagation terms for the fermions, the first terms in the fermion Lagrangian (11). Also, note we have defined

$$p \rightarrow p/f \quad (\text{A6})$$

for simplicity.

⁹This is a common assumption, motivated by the fact that higher order terms are expected to be suppressed by squares of ratios of form factors. In other words, this falls under the same assumption as the convergence of the integrals.

2. CP breaking vacuum

Here we repeat the exercise in the previous section to compute the coefficients of the CP breaking vacuum potential,

$$V(\eta, h) = c_{\text{tad}}\eta + m_\eta^2\eta^2 + \tilde{c}_\eta\eta^3 + \lambda_\eta\eta^4 + m_h^2h^2 + \lambda_hh^4 + c_3\eta h^2 + c_4\eta^2h^2. \quad (\text{A7})$$

The coefficients c_i are in general nonzero, except for at $\alpha = 1/4\pi$. Below we compute the parameters in an example with $\alpha = 1/3\pi$ case. As argued in the main text, the $\alpha = 2/3\pi$ case can be obtained from this by making the substitution $\eta \rightarrow -\eta$ in the potential¹⁰:

$$c_{\text{tad}} = -2N_c f^3 \int \frac{d^4 p}{(2\pi)^4} \frac{1}{2\Pi_0} \sqrt{3}\eta \Pi_1^t ((\epsilon_u - 4)\epsilon_u - 1)(\Pi_0^q + \Pi_0^{q'}), \quad (\text{A8a})$$

$$m_\eta^2 = -2N_c f^2 \int \frac{d^4 p}{(2\pi)^4} \frac{1}{\Pi_0} \left(\frac{3\Pi_1^t ((\epsilon_u - 4)\epsilon_u - 1)^2 (\Pi_0^q + \Pi_0^{q'})^2}{8\Pi_0} - \Pi_1^t (\epsilon_u^2 - 1)(\Pi_0^q + \Pi_0^{q'}) \right), \quad (\text{A8b})$$

$$\tilde{c}_\eta = -2N_c f \int \frac{d^4 p}{(2\pi)^4} \left(-\frac{\sqrt{3}\eta^3 (\Pi_1^t)^2 (\epsilon_u^2 - 1)((\epsilon_u - 4)\epsilon_u - 1)(\Pi_0^q + \Pi_0^{q'})(\Pi_0^q + \Pi_0^{q'})}{2\Pi_0^2} \right), \quad (\text{A8c})$$

$$\lambda_\eta = -2N_c \int \frac{d^4 p}{(2\pi)^4} \frac{(\Pi_1^t)^2 (\epsilon_u^2 - 1)^2 (\Pi_0^q + \Pi_0^{q'})^2}{2\Pi_0^2}, \quad (\text{A8d})$$

$$m_h^2 = -2N_c \int \frac{d^4 p}{(2\pi)^4} \frac{(p^2 (\Pi_1^t (\epsilon_u^2 + 3)(\Pi_0^q + 2\Pi_1^q + \Pi_0^{q'}) - 2\Pi_1^q \Pi_0^t) - 2M_t^2 (\epsilon_u^2 + 3))}{2p^2 \Pi_0}, \quad (\text{A8e})$$

$$\lambda_h = -2N_c f^2 \int \frac{d^4 p}{(2\pi)^4} \frac{(\epsilon_u^2 + 3)(M_t^2 - p^2 \Pi_1^q \Pi_1^t)}{p^2 \Pi_0} + \frac{(\Pi_1^t (\epsilon_u^2 + 3)(\Pi_0^q + 2\Pi_1^q + \Pi_0^{q'}) - 2\Pi_1^q \Pi_0^t)^2}{8\Pi_0^2}, \quad (\text{A8f})$$

$$c_3 = -2N_c f \int \frac{d^4 p}{(2\pi)^4} \left(\frac{\sqrt{3}((\epsilon_u - 4)\epsilon_u - 1)(M_t^2 - p^2 \Pi_1^q \Pi_1^t)}{p^2 \Pi_0} \right. \quad (\text{A8g})$$

$$\left. + \frac{\sqrt{3}\Pi_1^t ((\epsilon_u - 4)\epsilon_u - 1)(\Pi_0^q + \Pi_0^{q'}) (\Pi_1^t (\epsilon_u^2 + 3)(\Pi_0^q + 2\Pi_1^q + \Pi_0^{q'}) - 2\Pi_1^q \Pi_0^t)}{4\Pi_0^2} \right) \quad (\text{A8h})$$

$$c_4 = -2N_c \int \frac{d^4 p}{(2\pi)^4} \left(\frac{2(\epsilon_u^2 - 1)(p^2 \Pi_1^q \Pi_1^t - M_t^2)}{p^2 \Pi_0} \right. \quad (\text{A8i})$$

$$\left. - \frac{\eta^2 h^2 \Pi_1^t (\epsilon_u^2 - 1)(\Pi_0^q + \Pi_0^{q'}) (\Pi_1^t (\epsilon_u^2 + 3)(\Pi_0^q + 2\Pi_1^q + \Pi_0^{q'}) - 2\Pi_1^q \Pi_0^t)}{2\Pi_0^2} \right) \quad (\text{A8j})$$

where again Π_0 is the relevant field independent factor, here given by

$$\Pi_0 = \frac{1}{2} (\Pi_0^q + \Pi_0^{q'}) (\Pi_0^t - 2\Pi_1^t (\epsilon_u^2 + 1)). \quad (\text{A9})$$

As explained in the main text, the tadpole term can be shifted away by an appropriate shift in the other parameters, corresponding to a vev for η :

$$c_{\text{tad}} + 2m_\eta^2 v_\eta + 3\tilde{c}_\eta v_\eta^2 + 4\lambda_\eta v_\eta^3 = 0.$$

¹⁰These are again the parameters before shifting away the tadpole term, in exactly the same way as above.

The new parameters will then be given in terms of the quoted parameters as

$$m_\eta^2 \rightarrow m_\eta^2 + 3\tilde{c}_\eta v_\eta + 6\lambda_\eta v_\eta^2, \quad (\text{A10a})$$

$$\tilde{c}_\eta \rightarrow \tilde{c}_\eta + 4v_\eta \lambda_\eta, \quad (\text{A10b})$$

$$m_h^2 \rightarrow c_3 v_\eta + c_4 v_\eta^2, \quad (\text{A10c})$$

$$c_3 \rightarrow c_3 + 2c_4 v_\eta. \quad (\text{A10d})$$

-
- [1] D. H. Lyth, *Phys. Rev. Lett.* **78**, 1861 (1997).
[2] K. Agashe, R. Contino, and A. Pomarol, *Nucl. Phys.* **B719**, 165 (2005).
[3] B. Bellazzini, C. Csi, and J. Serra, *Eur. Phys. J. C* **74**, 2766 (2014).
[4] K. Freese, J. A. Frieman, and A. V. Olinto, *Phys. Rev. Lett.* **65**, 3233 (1990).
[5] N. Arkani-Hamed, H.-C. Cheng, P. Creminelli, and L. Randall, *Phys. Rev. Lett.* **90**, 221302 (2003).
[6] A. D. Linde, *Phys. Rev. D* **49**, 748 (1994).
[7] J. E. Kim, H. P. Nilles, and M. Peloso, *J. Cosmol. Astropart. Phys.* **01** (2005) 005.
[8] S. Dimopoulos, S. Kachru, J. McGreevy, and J. G. Wacker, *J. Cosmol. Astropart. Phys.* **08** (2008) 003.
[9] E. J. Copeland, A. Mazumdar, and N. J. Nunes, *Phys. Rev. D* **60**, 083506 (1999).
[10] E. Silverstein and A. Westphal, *Phys. Rev. D* **78**, 106003 (2008).
[11] N. Arkani-Hamed, H.-C. Cheng, P. Creminelli, and L. Randall, *J. Cosmol. Astropart. Phys.* **07** (2003) 003.
[12] D. Croon and V. Sanz, *J. Cosmol. Astropart. Phys.* **02** (2015) 008.
[13] D. Croon, V. Sanz, and J. Setford, *J. High Energy Phys.* **10** (2015) 020.
[14] P. Ade *et al.* (Planck Collaboration), *arXiv:1502.02114*.
[15] P. W. Graham, D. E. Kaplan, and S. Rajendran, *Phys. Rev. Lett.* **115**, 221801 (2015).
[16] J. R. Espinosa, C. Grojean, G. Panico, A. Pomarol, O. Pujol, and G. Servant, *Phys. Rev. Lett.* **115**, 251803 (2015).
[17] C. Gross, O. Lebedev, and M. Zatta, *Phys. Lett. B* **753**, 178 (2016).
[18] L. Kofman, A. D. Linde, and A. A. Starobinsky, *Phys. Rev. D* **56**, 3258 (1997).
[19] D. I. Podolsky, G. N. Felder, L. Kofman, and M. Peloso, *Phys. Rev. D* **73**, 023501 (2006).
[20] J. Braden, L. Kofman, and N. Barnaby, *J. Cosmol. Astropart. Phys.* **07** (2010) 016.
[21] J. Galloway, J. A. Evans, M. A. Luty, and R. A. Tacchi, *J. High Energy Phys.* **10** (2010) 086.
[22] B. Gripaios, A. Pomarol, F. Riva, and J. Serra, *J. High Energy Phys.* **04** (2009) 070.
[23] V. A. Rubakov and M. E. Shaposhnikov, *Phys. Lett.* **125B**, 136 (1983).
[24] R. Barbieri and G. F. Giudice, *Nucl. Phys.* **B306**, 63 (1988).
[25] R. Allahverdi, R. Brandenberger, F.-Y. Cyr-Racine, and A. Mazumdar, *Annu. Rev. Nucl. Part. Sci.* **60**, 27 (2010).
[26] J. H. Traschen and R. H. Brandenberger, *Phys. Rev. D* **42**, 2491 (1990).
[27] Y. Shtanov, J. H. Traschen, and R. H. Brandenberger, *Phys. Rev. D* **51**, 5438 (1995).
[28] A. Linde, S. Mooij, and E. Pajer, *Phys. Rev. D* **87**, 103506 (2013).
[29] P. Adshead, J. T. Giblin, T. R. Scully, and E. I. Sfakianakis, *J. Cosmol. Astropart. Phys.* **12** (2015) 034.
[30] E. W. Kolb, A. Notari, and A. Riotto, *Phys. Rev. D* **68**, 123505 (2003).
[31] M. Drewes, *J. Cosmol. Astropart. Phys.* **11** (2014) 020.
[32] M. Drewes and J. U. Kang, *Nucl. Phys.* **B875**, 315 (2013).
[33] M. Kawasaki, K. Kohri, and N. Sugiyama, *Phys. Rev. Lett.* **82**, 4168 (1999).
[34] K. Ichikawa, M. Kawasaki, and F. Takahashi, *Phys. Rev. D* **72**, 043522 (2005).
[35] J. Martin, C. Ringeval, and V. Vennin, *Phys. Rev. Lett.* **114**, 081303 (2015).
[36] S. Hannestad, *Phys. Rev. D* **70**, 043506 (2004).
[37] A. G. Cohen, D. B. Kaplan, and A. E. Nelson, *Annu. Rev. Nucl. Part. Sci.* **43**, 27 (1993).
[38] J. R. Espinosa, B. Gripaios, T. Konstandin, and F. Riva, *J. Cosmol. Astropart. Phys.* **01** (2012) 012.
[39] M. Gorbahn, J. M. No, and V. Sanz, *J. High Energy Phys.* **10** (2015) 036.
[40] G. Aad *et al.* (ATLAS Collaboration), *Phys. Rev. D* **90**, 112015 (2014).
[41] V. Khachatryan *et al.* (CMS Collaboration), *Eur. Phys. J. C* **74**, 3076 (2014).
[42] G. Aad *et al.* (ATLAS Collaboration), *Phys. Rev. D* **91**, 012006 (2015).
[43] S. Chatrchyan *et al.* (CMS Collaboration), *Phys. Rev. D* **89**, 092007 (2014).
[44] ATLAS Collaboration, Report No. ATLAS-CONF-2013-030, 2013.
[45] S. Chatrchyan *et al.* (CMS Collaboration), *J. High Energy Phys.* **01** (2014) 096.

- [46] T. A. collaboration (ATLAS Collaboration) (2013).
- [47] S. Chatrchyan *et al.* (CMS Collaboration), [Phys. Rev. D **89**, 012003 \(2014\)](#).
- [48] T. A. collaboration (ATLAS Collaboration) (2013).
- [49] S. Chatrchyan *et al.* (CMS Collaboration), [J. High Energy Phys. **05** \(2014\) 104](#).
- [50] M. Baak, J. Cth, J. Haller, A. Hoecker, R. Kogler, K. Mnig, M. Schott, and J. Stelzer (Gfitter Group), [Eur. Phys. J. C **74**, 3046 \(2014\)](#).
- [51] J. Fan, M. Reece, and L.-T. Wang, [J. High Energy Phys. **09** \(2015\) 196](#).
- [52] K. Hagiwara, S. Matsumoto, D. Haidt, and C. S. Kim, [Z. Phys. C **64**, 559 \(1994\); **68**, 352\(E\) \(1995\)](#).
- [53] D. M. Asner *et al.*, [arXiv:1310.0763](#).
- [54] M. Bicer *et al.* (TLEP Design Study Working Group), [J. High Energy Phys. **01** \(2014\) 164](#).
- [55] M. Redi and A. Tesi, [J. High Energy Phys. **10** \(2012\) 166](#).