

Wave instabilities in the presence of non vanishing background in nonlinear Schrodinger systems

Article (Unspecified)

Trillo, S, Toterogongora, J S and Fratalocchi, A (2014) Wave instabilities in the presence of non vanishing background in nonlinear Schrodinger systems. Scientific Reports, 4. p. 7285. ISSN 2045-2322

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/69633/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.



OPEN

Wave instabilities in the presence of non vanishing background in nonlinear Schrödinger systems

S. Trillo¹, J. S. Toterogongora² & A. Fratalocchi²

¹Dipartimento di Ingegneria, Università di Ferrara, Via Saragat 1, 44122 Ferrara, Italy, ²PRIMALIGHT, Faculty of Electrical Engineering; Applied Mathematics and Computational Science, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia.

SUBJECT AREAS:
OPTICS AND PHOTONICS
APPLIED MATHEMATICS

Received
17 September 2014

Accepted
14 November 2014

Published
3 December 2014

Correspondence and requests for materials should be addressed to A.F. (andrea.fratalocchi@kaust.edu.sa)

We investigate wave collapse ruled by the generalized nonlinear Schrödinger (NLS) equation in 1+1 dimensions, for localized excitations with non-zero background, establishing through virial identities a new criterion for blow-up. When collapse is arrested, a semiclassical approach allows us to show that the system can favor the formation of dispersive shock waves. The general findings are illustrated with a model of interest to both classical and quantum physics (cubic-quintic NLS equation), demonstrating a radically novel scenario of instability, where solitons identify a marginal condition between blow-up and occurrence of shock waves, triggered by arbitrarily small mass perturbations of different sign.

Wave collapse, i.e. the occurrence of blow up in a finite time or propagation distance^{1–3}, is a general phenomenon that appears in many contexts including self-focusing in optics^{4,5} (for an extensive review and a historical perspective, see ref. 6), plasma waves⁷, Bose-Einstein condensates^{8,9}, hydrodynamics¹⁰ and organic systems¹¹. Indeed several dispersive models with critical or supercritical nonlinearity (e.g. generalized KdV or KP equations¹², modified KP¹³, Zakharov equation¹⁴, and discrete or nonlocal version of the nonlinear Schrödinger equation^{15,16}) exhibit formation of a point singularity from a large class of initial data. The critical case is by far the most intriguing one due to the sensitivity of the collapse to perturbations. In this context the most studied case, namely the focusing cubic nonlinear Schrödinger (NLS) equation, which is critical in two transverse dimensions, is still a formidable ground to understand the dynamics of collapse^{17–20} in spite of long standing investigations^{3–6}. However, many physical classical and quantum systems need to be described in terms of generalized NLS (gNLS) equation which accounts for higher-order nonlinearities. Indeed such higher-order terms arise from different physical mechanisms in nonlinear optics (saturation of optical susceptibilities of standard material^{21–23}, local field effects²⁴, tailoring of nonlinearities in a cooled gas²⁵), dynamics of superfluids²⁶, or quantum condensed systems where they are related to higher-order atom-atom interactions^{27–29}. In gNLS systems critical collapse can occur also in one transverse dimension when quintic (or higher-order) nonlinearity is effective^{2,30,31}. Though the blow-up problem is generally addressed for bright (i.e., zero background) solutions, gNLS systems support translationally invariant solitary wave solutions with non-zero background, yet of bright type^{32–36} (complement of well known dark soliton or bubble type solutions^{37,38}), for which the problem of blow-up have been overlooked. Our aim in this paper is to establish a novel criterion for collapse valid for solutions of this type, and study the dynamics across the threshold for blow-up. This allows us to reveal a new instability scenario where opposite behaviors, such as blowup or decay into a dispersive shock wave, can be controlled by means of an arbitrarily weak perturbation which controls the variation of the power (or mass) integral of a launched perturbed solitary wavepacket. We consider regimes for which the background itself is stable (otherwise modulation instability of the background becomes the main mechanism that affects the decay dynamics of the field).

The paper is organized as follows. In Section 2 we introduce the model and develop a sufficient criterion for collapse. In Section 3, we propose an approach to characterize non-collapsing solutions based on self-similar and semiclassical scaling arguments. The prediction of our analysis are confirmed by means of numerical simulations with reference to a specific model of physical interest in Section 4.



The model and collapse criterion. We start from the following gNLS equation in dimensionless units

$$i \frac{\partial \psi}{\partial t} + \frac{\partial^2 \psi}{\partial x^2} + F(|\psi|^2) \psi = 0, \quad (1)$$

with a general nonlinearity $F(\rho)$ (henceforth $\rho = |\psi|^2$, which has physical meaning of density or intensity in optics) and study the instability scenario of localized fields ψ characterized by nonvanishing boundary conditions: $\psi(\pm\infty) \rightarrow \sqrt{\rho_0} e^{\mp i\phi_0}$. Equation (1) is conveniently normalized by setting $F(\rho_0) = 0$ at the background value ρ_0 . Invariants of motion originate from the translational symmetries along t , x and Gauge transformations, resulting in the renormalized functionals associated with energy H , momentum P , and L^2 norm M (henceforth referred to as ‘mass’, though it can physically represent number of particles or power in optics)³⁹

$$H = \int \left[\left| \frac{\partial \psi}{\partial x} \right|^2 - U(|\psi|^2) \right] dx, \quad (2)$$

$$P = \int \text{Re} \left[i\psi \frac{\partial \psi^*}{\partial x} (1 - \rho_0/|\psi|^2) \right] dx, \quad (3)$$

$$M = \int (|\psi|^2 - \rho_0) dx, \quad (4)$$

where $U(\rho) \equiv \int_{\rho_0}^{\rho} F(y) dy$.

As demonstrated by Barashenkov³⁸, traveling solitary solutions $\psi(x, t) = \phi(x - vt)$ of the gNLS equation (1) on a finite background $\rho_0 \neq 0$ are unstable when the slope of the curve $P = P(v)$ becomes negative ($dP/dv < 0$). Under the validity of such assumption, we begin our theoretical analysis by finding a sufficient criterion for collapse based on virial identities (see the review papers in Refs. 1–3, 6. We address the most common regime where, in the absence of an initial derivative of the virial (second-order momentum), the collapse of a bright disturbance with vanishing background is known to be essentially ruled by the energy (Hamiltonian) invariant of motion. Conversely, for a finite background we show that blow-up is driven by a suitable combination of renormalized energy and mass, thus allowing the existence of non collapsing solutions even for negative energy. We then employ a combination of virial identities and arguments based on self-similarity to study the leading order scaling of the dynamics in both collapsing and non collapsing regimes. In the latter case, a suitable rescaling in a semiclassical form and a diagonalization of the resulting equations of motion in terms of Riemann invariants, allow us to predict a completely novel scenario where collapse is arrested and the instability leads to the decay into dispersive shock waves, i.e. expanding fast wavetrains that are naturally emitted, owing to dispersion, to regularize steep gradients developing through the nonlinearities^{40–51}. We verified our theoretical finding on a model of physical interest, namely the defocusing-focusing Cubic-Quintic Nonlinear Schrödinger Equation (CQNLS). We demonstrate a specific class of localized pulses on a pedestal, represented by antidark solitons, marking a marginal condition crossover between collapse and shock-wave generation, controlled via (even infinitesimal) mass perturbations. These findings, other than the fundamental interest related to collapse, suggest a new avenue for arresting blow-up and generating dispersive shocks from unstable solutions with nonzero background.

We begin our theoretical analysis by writing the Hamilton-Jacobi equations of motion for the evolution of a suitable renormalized virial $V(t) = \int dx x^2 (|\psi|^2 - \rho_0)$:

$$\frac{\partial V}{\partial t} = -2i \langle \psi | x \partial / \partial x | \psi \rangle + c.c., \quad (5)$$

$$\frac{\partial^2 V}{\partial t^2} = 4[H + G(|\psi|^2)], \quad (6)$$

where we set $\langle \psi | f(x) | \psi \rangle \equiv \int dx \psi^* f(x) \psi$, and Eq. (6) is obtained by further deriving in time Eq. (5) and expressing such derivative in terms of the energy functional H , and the suitably defined functional $G(\rho) \equiv \int dx [2U(\rho) - \rho F(\rho)]$, which depends on the nonlinear response of the system. In order to derive a sufficient criterion for collapse, we expand in series the functional $G(\rho)$, obtaining

$$G(\rho) = - \sum_{m=1}^{\infty} \frac{F^{(m)}}{(m+1)!} \int dx (\rho - \rho_0)^m [\rho(m-1) + 2\rho_0], \quad (7)$$

with $F^{(m)} \equiv \frac{\partial^m F}{\partial \rho^m} \Big|_{\rho=\rho_0}$. Several different scenario can arise according to the signs of the terms in the expansion (7). An interesting case manifests when a low order defocusing nonlinearity—which supports localized solutions on a pedestal—competes with high order contributions of different sign, i.e., when $F^{(1)} < 0$ and $F^{(m)} > 0$ for $m > 2$. The following inequality then holds

$$G(\rho) \leq -MF^{(1)}\rho_0. \quad (8)$$

Then, combining Eq. (6) and (8), we find the following differential inequality

$$\frac{\partial^2 V}{\partial t^2} = 4[H + G(\rho)] \leq 4(H - MF^{(1)}\rho_0) \equiv 4H_r, \quad (9)$$

which gives an upper bound to the dynamics of the virial $V(t)$ in terms of rescaled Hamiltonian H_r , depending on the renormalized energy H and mass M . When $H_r < 0$, or equivalently:

$$H - MF^{(1)}\rho_0 < 0, \quad (10)$$

the virial $V(t) \sim Ct^2$ (C negative constant) tends to zero regardless of any initial condition $V(0)$ and $\partial V(0)/\partial t$. In order to understand the consequences of a vanishing virial on the dynamics of the field ψ , we employ a specific form of the Hölder inequality:

$$\left(\int dx x \sqrt{\rho - \rho_0} \frac{\partial \sqrt{\rho - \rho_0}}{\partial x} \right)^2 \leq \int dx \left(\frac{\partial \sqrt{\rho - \rho_0}}{\partial x} \right)^2 \cdot \int dx x^2 (\rho - \rho_0), \quad (11)$$

and through an integration by parts of Eq. (11), derive the following upper limit for the virial evolution,

$$V(t) \geq \frac{M^2}{4 \int dx \left(\frac{\partial \sqrt{\rho - \rho_0}}{\partial x} \right)^2}. \quad (12)$$

According to Eq. (12), if the virial tends to zero, the gradient norm $\int dx \left(\frac{\partial \sqrt{\rho - \rho_0}}{\partial x} \right)^2$ diverges in order to maintain an invariant mass M . This condition leads to the formation of a singularity in the field $\rho - \rho_0$. Equation (10) can be therefore regarded as a sufficient condition for blow-up in the generalized NLS equation (1) with non vanishing boundary conditions. An important observation, stemming from Eq. (12), concerns the role of the background during the collapse. In particular the gradient norm, which drives the collapsing dynamics, does not depend on the background ρ_0 . The latter is therefore unaffected by the dynamics and the density evolves by respecting the boundary conditions $\rho(\pm\infty, t) = \rho_0$.



Self-similar analysis and semiclassical argument. The differential inequality (9) can be also employed to calculate the leading order dynamics in both collapsing and non collapsing regimes. We start from the following self-similar form of the field

$$\rho(x,t) = b(t)S\left[\frac{x}{\sqrt{a(t)}}\right] + \rho_0, \quad a(0) = a_0, \quad (13)$$

which can be used to extract the scaling dynamics in the unstable regime³, and then employ the conservation of the mass M to determine the relationship between the peak value b and the width (variance) a of the field $|\psi|^2 - \rho_0$,

$$b(t)\sqrt{a(t)} = \frac{M}{I_0}, \quad (14)$$

being $I_0 = \int dx S(x)$. By substituting Eqs. (13) and (14) into Eq. (9), we obtain

$$\frac{\partial^2 a}{\partial t^2} \leq 4 \frac{I_0}{MI_2} H_r, \quad (15)$$

having introduced the constant $I_2 = \int dx x^2 S(x)$ representing the second-order moment of inertia of $S(x)$ with respect to $x = 0$. By integrating twice Eq. (15) in time, we straightforwardly obtain the following lower bound for the evolution of the peak b of $|\psi|^2 - \rho_0$

$$b(t) \geq \frac{\sqrt{M^3 I_2}}{I_0} \frac{1}{\sqrt{2I_0 H_r t^2 + a_0 M I_2}}, \quad (16)$$

which states that at leading order the field power $b(t)$ scales as $b \sim \frac{1}{t}$ for a non collapsing solution possessing $H_r > 0$. This scaling has important consequences in the development of shock wave instabilities when the input self-similar parameter is small, i.e., for $a(0) \sim 2c^2$ with $\epsilon < 1$. In this case, we can rescale Eq. (1) with $x = y \frac{\sqrt{2}}{\epsilon}$ and $\tau = \epsilon t$, thus obtaining a semiclassical form of the NLS equation with a general nonlinearity:

$$i\epsilon \frac{\partial \psi}{\partial \tau} + \frac{\epsilon^2}{2} \frac{\partial^2 \psi}{\partial y^2} + F(\rho)\psi = 0. \quad (17)$$

Equation (17) can be casted into a hydrodynamic (or so-called dispersionless) form by applying the Madelung transformation $\psi = \sqrt{\rho} e^{i \int dy u(y)}$ and taking the leading-order terms in ϵ , which yields

$$\frac{\partial \rho}{\partial \tau} + \frac{\partial(\rho u)}{\partial y} = 0, \quad \frac{\partial u}{\partial \tau} + u \frac{\partial u}{\partial y} + \frac{1}{\rho} \frac{\partial P}{\partial y} = 0, \quad (18)$$

which is physically equivalent to the system (also known as P -system) that rules the evolution of a compressible gas flow with density ρ , velocity u and pressure given through the equation of state $P = P(\rho)$. In our case such equation of state is explicitly given in terms of nonlinearity and density as $P(\rho) = \int d\rho F - \rho F(\rho) = - \int \rho \partial_\rho F(\rho) d\rho$, which in turn defines the sound velocity

$$c(\rho) = \sqrt{\frac{\partial P}{\partial \rho}} = \sqrt{-\rho \partial_\rho F} \quad (19)$$

Equivalently Eqs. (18) can be cast in the form of a system of quasi-linear equations $\partial_\tau \mathbf{q} + \mathbf{A} \partial_y \mathbf{q} = 0$, with $\mathbf{q} \equiv (\rho, u)^T$ and 2×2 matrix $\mathbf{A} = (u \rho; -\partial_\rho F u)$, which is diagonalizable in the form

$$\frac{\partial \lambda_\pm}{\partial \tau} + (u \pm c) \frac{\partial \lambda_\pm}{\partial y} = 0, \quad (20)$$

by introducing the Riemann invariants $\lambda_\pm = u \pm \int_{\rho_0}^{\rho} \frac{c(\rho)}{\rho} d\rho$ and the eigenvelocities of the system $v_\pm = u \pm c$, which correspond to the eigenvalues of the matrix \mathbf{A} . By expanding in Taylor series the nonlinearity $F(\rho)$ around ρ_0 , we obtain

$$c^2 = -\rho \left[F^{(1)} + \sum_{m=2}^{\infty} \frac{F^{(m)}}{(m-1)!} (\rho - \rho_0)^{m-1} \right]. \quad (21)$$

When Eq. (10) is not fulfilled and a collapse is not observed in the dynamics, the field density decreases rapidly (as shown above by the self-similarity argument), and higher-order terms in the sum in Eq.

(21) scale as $(\rho - \rho_0)^{m-1} \sim \frac{1}{t^{m-1}}$, $m \geq 2$, thus becoming negligible.

As a consequence the velocity c is determined by the lowest power of ρ in the leading order term $F^{(1)} \equiv \partial_\rho F|_{\rho=\rho_0}$. For leading-order cubic nonlinearities where $F(\rho) = \chi(\rho - \rho_0) + \dots$, with $\chi = \pm 1$ is the sign of the nonlinearity, we have $F^{(1)} = \chi$ and Eq. (21) yields $c = \sqrt{-\rho\chi}$. In this case, whenever $\chi = -1$ (leading-order cubic nonlinearity of defocusing or repulsive type), the corresponding Riemann velocities $v_\pm = u \pm \sqrt{\rho}$ are real and coincides (at leading order), with those pertaining to the integrable (cubic) defocusing NLS system⁴¹. As a consequence we expect the system to undergo wave-breaking through the development of a gradient catastrophe (formation of vertical front in finite time) from smooth input conditions launched in the system. In the former case the consequent regularization of the dynamics, owing to the dispersive terms neglected in Eqs. (17), occurs in term of a dispersive shock^{40,41,43-51}. Conversely, such argument loses validity in the case $\chi = 1$ (focusing cubic nonlinearity), where $v_\pm = u \pm \sqrt{-\rho}$ would imply a catastrophe of the elliptic umbilic type (characteristic of systems with dispersionless limit of the elliptic type with complex conjugate eigenvelocities v_\pm), where $\rho - \rho_0$ increases in time. In such a case higher-order terms in Eq. (21) become important and no conclusive statements on the type of dynamics can be made from such an approach.

In the following section we show, with reference to a specific system, how an arbitrarily small mass perturbation of a solitary waves with non-zero background can lead, depending on its sign, to completely different decay scenarios implying either collapse or dispersive shock formation.

Application to cubic-quintic nonlinearity. In order to illustrate our theoretical findings on a realistic system of physical interest, we consider the CQNLS equation with competing nonlinearities. The CQNLS is a general model for superfluidity²⁶, nonlinear optics of saturable nonlinearities or cascading effects^{24,37}, and systems of quantum condensed gases with elastic two- and three-body interactions^{27,28}. The regime of interest here is a leading-order (cubic) nonlinearity of the defocusing type saturated by a focusing quintic nonlinearity, which is described by the nonlinear function $F(\rho) = -(\rho - \rho_0) + \alpha(\rho^2 - \rho_0^2)$, where α is a dimensionless parameter describing the strength of the quintic term (responsible for the collapse) over the defocusing cubic nonlinearity. We consider the following input field $\rho(x, t = 0)$:

$$\rho(x,0) = \left(\frac{\rho_a r}{r + \tanh^2 wx} - \rho_0 \right) (1 + \delta)^2 + \rho_0, \quad (22)$$

where the small parameter δ is introduced to measure a *mass perturbation* of a still (zero velocity) anti-dark soliton $\rho_{sol}(x) = \rho_a r / (r + \tanh^2 wx)$ [see profile in figure 1(a)], which is an exact solitary solution of the CQNLS provided $r = \rho_0 / (\rho_a - \rho_0)$, $w = 2\sqrt{\alpha(\rho_a - \rho_0)\rho_0/3}$, $\rho_a = (3 - 4\alpha\rho_0)/2\alpha$ ³⁵. Zero velocity anti-dark solitons of CQNLS are *always unstable* in the spirit of Ref. 38 while their background is stable³⁵, thereby being ideal candidate for




Figure 1 | (a) Density distribution $\rho(x, t = 0)$ at fixed perturbation $\delta = 0.2$, and (b) Reduced energy functional $H_r = H - M \frac{\partial F}{\partial \rho} \Big|_{\rho_0}$ (solid line) and energy H (dashed line) vs. perturbation parameter δ , for the perturbed antidark soliton of CQNLS equation, expressed by Eq. (22) with $\alpha = 0.1$ and $\rho_0 = 1$.

our theoretical investigation. We begin our analysis by calculating the correspondent reduced energy H_r , shown in figure 1(b). As seen, any positive mass perturbation $\delta > 0$ leads to $H_r < 0$ and, by virtue of Eq. (10), to collapse. In order to numerically ascertain this result, we perform a direct integration of the CQNLS via a fourth order finite difference scheme. In our simulations, we launch Eq. (22) at $t = 0$ and monitor its dynamical evolution in the system for a fixed combination of α, ρ and a varying δ , which defines the mass perturbation of the CQNLS soliton. Our integration scheme is developed within an adaptive fourth order Runge-Kutta method⁵², which automatically adjusts the time resolution in order to maintain good accuracy on the results. The relative error tolerance in the simulation results is 10^{-9} . The spatial resolution of the grid was $dx = 8 \cdot 10^{-3}$. During each time-step, we also monitored the conserved quantities (2), checking that their value remained practically constant. Figure 2 summarizes our results for instance for $\delta = 0.1$, $\alpha = 0.1$ and a background $\rho_0 = 1$ (similar dynamics is obtained for other choices of values). In perfect agreement with our theory, a positive mass variation leads to a catastrophic collapse of the wave function [see figure 2(a–c)], which is attained at a relatively short distance, as shown in figure 2(c). We also observe, in agreement with Eq. (12), that the background intensity ρ_0 does not change during the collapsing dynamics, remaining stable as also predicted by the linear stability analysis against the growth of periodic modulations³⁵.




Figure 2 | Collapse ruled by the defocusing-focusing CQNLS: (a) color level plot of $\rho = |\psi|^2$ in time-space (t, x) ; (b) snapshots of $\rho(x)$ at $t = 0$ (dashed line) and $t = 0.049$ (solid line); (c) peak intensity (or density) evolution along t . Here the quintic coefficient is $\alpha = 0.1$, the background $\rho_0 = 1$, and we employ a positive-mass perturbation $\delta = 0.1$.

When the initial soliton mass is conversely decreased, for $\delta < 0$, we have $H_r > 0$ [see figure 1(b)]. In the limit of a perturbative quintic nonlinearity $\alpha < 1$, from Eq. (14) we obtain

$$a(0) = 2 \frac{M\alpha}{I_0(3 - 4\alpha\rho_0)} \sim 2\alpha < 1, \tag{23}$$

which leads to a semiclassical behavior described by Eqs. (17) – (21) with $\alpha = \sqrt{\epsilon}$. This predicts the generation of dispersive shock waves when $\delta < 0$. Figure 3 reports numerical simulations for the case of a negative mass perturbation with $\delta = -0.01$ and the same initial parameters ($\alpha = 0.1, \rho_0 = 1$) as in figure 2. In complete agreement with Eq. (16), the intensity amplitude falls off as $\sim \frac{1}{t}$ [see figure 3(b)]. This process leads to wave breaking occurring through two symmetric gradient catastrophes (formation of symmetric steep fronts), followed by dispersive shock wave generation, as displayed in figure 3(c). It is worthwhile observing that for $\delta = -0.01$ the energy H of the field ψ is still negative, as indicated by figure 1(b). At variance with the case of bright localized waveform on a zero background, which under the condition $H < 0$, are always collapsing in NLS systems³, our analysis demonstrates that a suitable background intensity can arrest the blow-up, leading to a completely different instability scenario where dispersive shocks are generated.




Figure 3 | Dispersive shock generation in the defocusing-focusing CQNLS: (a) color level plot of $\rho = |\psi|^2$ in time-space (x, t) ; (b) peak intensity (or density) $\max_x(|\psi(x, t)|^2)$ versus t ; (c) snapshots of $\rho(x)$ at $t = 0$ (dashed line), $t = 0.4$ (solid thin line) and $t = 3$ (solid thick line). The parameters are as in figure 2 ($\alpha = 0.1, \rho_0 = 1$), except for a negative-mass perturbation $\delta = -0.01$.



Conclusions

In conclusion, we investigated wave instability in NLS systems in the presence of a non vanishing background. Through virial identities and semiclassical analysis, we have demonstrated a sufficient criterion for collapse and unveiled a novel instability dynamics characterized by the emission of dispersive shocks. Theoretical findings are verified against numerical simulations on the CQNLS equation, where we unveiled a set of solutions that exhibit a marginal crossover between the two aforementioned regimes. Besides the large interest of the physical community in wave collapse, these results show that a non vanishing background acts an affective mechanism for arresting collapse, leading to a new scenario characterized by the emission of shock waves.

- Kuznetsov, E. A., Rubenchik, A. & Zakharov, V. Soliton stability in plasmas and hydrodynamics. *Phys. Rep.* **142**, 103–165 (1986).
- Rasmussen, J. J. & Rypdal, K. Blow-up in nonlinear schrodinger equations-i a general review. *Physica Scripta* **33**, 481–497 (1986).
- Bergé, L. Wave collapse in physics: principles and applications to light and plasma waves. *Phys. Rep.* **303**, 259 (1998).
- Kelley, P. L. Self focusing of optical beams. *Phys. Rev. Lett.* **15**, 1005 (1965).
- Gaeta, A. L. Catastrophic collapse of ultrashort pulses. *Phys. Rev. Lett.* **84**, 3582–3585 (2000).
- Boyd, R. W., Lukishova, S. & Shen, Y. R. *Self focusing: past and present* (Springer, Topics in Applied Physics vol. 114, New York, 2009).
- Wong, A. Y. & Cheung, P. Y. Three-dimensional self-collapse of langmuir waves. *Phys. Rev. Lett.* **52**, 1222–1225 (1984).
- Gerton, J. M., Strekalov, M., Prodan, I. & Hulet, R. G. Direct observation of growth and collapse of a bose-einstein condensate with attractive potential. *Nature* **408**, 692–695 (2000).
- Lahaye, T. *et al.* *d*-wave collapse and explosion of a dipolar bose-einstein condensate. *Phys. Rev. Lett.* **101**, 080401 (2008).
- Zeff, B. W., Kleber, B., Fineberg, J. & Lathrop, D. P. Singularity dynamics in curvature collapse and jet eruption on a fluid surface. *Nature* **403**, 401–404 (2000).
- Bang, O., Christiansen, P. L., If, F., Rasmussen, K. O. & Gaididei, Y. B. Temperature effects in a nonlinear model of monolayer schein aggregates. *Phys. Rev. E* **49**, 4627–4636 (1994).
- Wang, X., Ablowitz, M. & Segur, H. Wave collapse and instability of solitary waves of a generalized kadmetsv-petviashvili equation. *Physica D* **78**, 241–265 (1994).
- Whalen, P., Moloney, J. V., Newell, A. C., Newell, K. & Kolesik, M. Optical shock and blow-up of ultrashort pulses in transparent media. *Phys. Rev. A* **86**, 033806 (2012).
- Merle, F. Lower bounds for the blowup rate of solutions of the zakharov equation in dimension two. *Communications on Pure and Applied Mathematics* **49**, 765–794 (1996).
- Bang, O., Rasmussen, J. J. & Christiansen, P. L. Subcritical localization in the discrete nonlinear schrodinger equation with arbitrary power nonlinearity. *Nonlinearity* **7**, 205–218 (1994).
- Bang, O., Krolkowski, W., Wyller, J. & Rasmussen, J. J. Collapse arrest and soliton stabilization in nonlocal nonlinear media. *Phys. Rev. E* **66**, 046619–(5) (2002).
- Skupin, S., Saffman, M. & Królkowski, W. Nonlocal stabilization of nonlinear beams in a self-focusing atomic vapor. *Phys. Rev. Lett.* **98**, 263902 (2007).
- Linzon, Y., Rutkowska, K. A., Malomed, B. A. & Morandotti, R. Magneto-optical control of light collapse in bulk kerr media. *Phys. Rev. Lett.* **103**, 053902 (2009).
- Desyatnikov, A. S., Buccoliero, D., Dennis, M. R. & Kivshar, Y. S. Suppression of collapse for spiraling elliptic solitons. *Phys. Rev. Lett.* **104**, 053902 (2010).
- Shim, B., Schrauth, S. E., Gaeta, A. L., Klein, M. & Fibich, G. Loss of phase of collapsing beams. *Phys. Rev. Lett.* **108**, 043902 (2012).
- Chen, Y.-F. *et al.* Measurement of fifth- and seventh-order nonlinearities of glasses. *J. Opt. Soc. Am. B* **23**, 347–352 (2005).
- Béjot, P. *et al.* Higher-order kerr terms allow ionization-free filamentation in gases. *Phys. Rev. Lett.* **104**, 103903 (2010).
- Béjot, P. *et al.* Transition from plasma-driven to kerr-driven laser filamentation. *Phys. Rev. Lett.* **106**, 243902 (2011).
- Dolgaleva, K., Shin, H. & Boyd, R. W. Observation of a microscopic cascaded contribution to the fifth-order nonlinear susceptibility. *Phys. Rev. Lett.* **103**, 113902 (2009).
- Greenberg, J. A. & Gauthier, D. J. Higher-order optical nonlinearity at low light levels. *Eur. Phys. Lett.* **98**, 24001 (2012).
- Josserand, C. Cavitation induced by explosion in an ideal fluid model. *Phys. Rev. E* **60**, 482–491 (1999).
- Abdullaev, F. K., Gammal, A., Tomio, L. & Frederico, T. Stability of trapped bose-einstein condensates. *Phys. Rev. A* **63**, 043604 (2001).
- Pieri, P. & Strinati, G. C. Derivation of the gross-pitaevskii equation for condensed bosons from the bogoliubovde gennes equations for superfluid fermions. *Phys. Rev. Lett.* **91**, 030401 (2003).
- Kuznetsov, E. & Turitsyn, S. Instability and collapse of solitons in media with a defocusing nonlinearity. *Sov. Phys. JETP* **67**, 1583–1588 (1988).
- Glassey, R. T. On the blowing up of solutions to the cauchy problem for nonlinear schrödinger equations. *J. Math. Phys.* **18**, 1794 (1977).
- Chung, Y. & Lushnikov, P. M. Strong collapse turbulence in a quintic nonlinear schrödinger equation. *Phys. Rev. E* **84**, 036602 (2011).
- Kivshar, Y. S. & Afanasjev, V. V. Dark optical solitons with reverse-sign amplitude. *Phys. Rev. A* **44**, R1446–R1449 (1991).
- Frantzeskakis, D., Hizanidis, K., Malomed, B. & Polymilis, C. Stable anti-dark light bullets supported by the third-order dispersion. *Physics Letters A* **248**, 203–207 (1998).
- Coskun, T. H. *et al.* Bright spatial solitons on a partially incoherent background. *Phys. Rev. Lett.* **84**, 2374–2377 (2000).
- Crosta, M., Fratolocchi, A. & Trillo, S. Bistability and instability of dark-antidark solitons in the cubic-quintic nonlinear schrödinger equation. *Phys. Rev. A* **84**, 063809 (2011).
- Zakharov, V. E. & Gelash, A. A. Nonlinear stage of modulation instability. *Phys. Rev. Lett.* **111**, 054101 (2013).
- Kivshar, Y. S. & Luther-Davies, B. Dark optical solitons: physics and applications. *Phys. Rep.* **298**, 81–197 (1998).
- Barashenkov, I. V. Stability criterion for dark solitons. *Phys. Rev. Lett.* **77**, 1193–1197 (1996).
- Kivshar, Y. S. & Agrawal, G. P. *Optical Solitons: from fibers to photonic crystals* (Academic Press, San Diego, 2003).
- Gurevich, A. V. & Pitaevskii, L. P. Nonstationary structure of a collisionless shock wave. *Sov. Phys. JETP* **38**, 291 (1973).
- Kamchatnov, A. M. *Nonlinear Periodic Waves and Their Modulations* (World Scientific, Singapore, 2000).
- Kamchatnov, A. M., Gammal, A. & Kraenkel, R. A. Dissipationless shock waves in bose-einstein condensates with repulsive interaction between atoms. *Phys. Rev. A* **69**, 063605 (2004).
- El, G. A. Resolution of a shock in hyperbolic systems modified by weak dispersion. *Chaos* **15**, 037103 (2005).
- Fratolocchi, A., Conti, C., Ruocco, G. & Trillo, S. Free-energy transition in a gas of noninteracting nonlinear wave particles. *Phys. Rev. Lett.* **101**, 044101 (2008).
- Dutton, Z., Budde, M., Slowe, C. & Hau, L. V. Observation of quantum shock waves created with ultra-compressed slow light pulses in a bose-einstein condensate. *Science* **27**, 293 (2001).
- Hofer, M. A. *et al.* Dispersive and classical shock waves in Bose-Einstein condensates and gas dynamics. *Phys. Rev. A* **74**, 023623 (2006).
- Meppelink, R. *et al.* Observation of shock waves in a large bose-einstein condensate. *Phys. Rev. A* **80**, 043606 (2009).
- Rothenberg, J. E. & Grischkowsky, D. Observation of the formation of an optical intensity shock and wave breaking in the nonlinear propagation of pulses in optical fibers. *Phys. Rev. Lett.* **62**, 531–534 (1989).
- Wan, W. J., Jia, S. & Fleischer, J. Dispersive superfluid-like shock waves in nonlinear optics. *Nature Physics* **3**, 46 (2007).
- Ghofraniha, N., Conti, C., Ruocco, G. & Trillo, S. Shocks in nonlocal media. *Phys. Rev. Lett.* **99**, 043903 (2007).
- Conti, C., Fratolocchi, A., Peccianti, M., Ruocco, G. & Trillo, S. Observation of a gradient catastrophe generating solitons. *Phys. Rev. Lett.* **102**, 083902 (2009).
- Shampine, L. F. (ed.) *Numerical Solution of Ordinary Differential Equations* (Chapman & Hall, New York, 1994).

Acknowledgments

We acknowledge funding from Italian Ministry of University and Research (MIUR, grant PRIN 2012BFNWZ2) and KAUST (Award No. CRG-1-2012-FRA-005).

Author contributions

A.F. and S.T. conceived the work and developed the theoretical analysis. J.S.T.G. performed the numerical simulations. A.F. and S.T. reviewed and wrote the manuscript.

Additional information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Trillo, S., Gongora, J.S.T. & Fratolocchi, A. Wave instabilities in the presence of non vanishing background in nonlinear Schrödinger systems. *Sci. Rep.* **4**, 7285; DOI:10.1038/srep07285 (2014).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder in order to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>