

Unsymmetrical, cyclic diborenes and thermal rearrangement to a borylborylene

Article (Accepted Version)

Stennett, Tom E, Mattock, James D, Vollert, Ivonne, Vargas, Alfredo and Braunschweig, Holger (2018) Unsymmetrical, cyclic diborenes and thermal rearrangement to a borylborylene. *Angewandte Chemie International Edition*, 57 (15). pp. 4098-4102. ISSN 1433-7851

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/73693/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Unsymmetrical, Cyclic Diborenes and Thermal Rearrangement to a Borylborylene

Tom E. Stennett,^[a] James D. Mattock,^[b] Ivonne Vollert,^[a] Alfredo Vargas^{*[b]} and Holger Braunschweig^{*[a]}


Abstract: Cyclic diboranes(4) based on a chelating monoanionic benzylphosphine linker were prepared by boron-silicon exchange between arylsilanes and B₂Br₄. Coordination of Lewis bases to the remaining sp² boron atom yielded unsymmetrical sp³-sp³ diboranes, which were reduced with KC₈ to their corresponding *trans*-diborenes. These compounds were studied by a combination of spectroscopic methods, X-ray diffraction and DFT calculations. PMe₃-stabilized diborene **6** was found to undergo thermal rearrangement to *gem*-diborene **8**. DFT calculations on **8** reveal a polar boron-boron bond, and indicate that the compound is best described as a borylborylene.

bridged sp²-sp³ diboranes by reaction of B₂Cl₄ with 2-silylamino- and 2-siloxypyridines, forming B-N and B-O bonds, respectively, by Me₃SiCl elimination (Scheme 1).^[12] The group of Siebert later succeeded in transferring a Cp* group to B₂Cl₄ via chlorosilane elimination, with a rearrangement giving a cyclopentadienylborylene-borane adduct as the final product.^[13] Inspired by these studies, we selected a readily prepared arylsilane containing a phosphine donor group, with the goal of generating new diboranes capable of being reduced to B-B multiply-bonded compounds. From this starting point, we present a rational synthesis of unsymmetrical diborenes and rearrangement of one of these compounds to its *gem* isomer.^[14]

Since the report of the first neutral diborene by Robinson and co-workers, which was prepared in low yields via a radical hydrogen scavenging process,^[1] the preparation of compounds bearing boron-boron double bonds^[2] has been extended by three rational and selective synthetic routes – the reductive coupling of two N-heterocyclic carbene (NHC)-stabilized arylidihaloboranes,^[2b, 3] the reduction of B₂X₂R₂ species in the presence of stabilizing phosphine ligands,^[4] and the stoichiometric reduction of carbene adducts of B₂Br₄.^[5] A very recent development in this field has been the realization of the first unsymmetrical diborenes,^[5a, 6] which promise to reveal more information about the fundamental nature of the boron-boron π-bond, and also allow investigation of the effect of a polarised B-B bond on the species' behaviour.

We are currently especially interested in the reactivity of phosphine-stabilized diborenes. Whereas carbenes typically remain bound to boron during the course of the reactions of diborenes with small molecules,^[3b, 5a, 7] phosphine donors can dissociate under UV light and their diborenes display markedly different reactivity.^[8] A drawback of the currently known examples of phosphine-stabilized diborenes is that they are mostly prepared from 1,2-dihalo-1,2-diarylboranes, B₂X₂R₂, the syntheses of which are limited in terms of substituent scope.^[9] With convenient routes now available to all of the tetrahalodiboranes(4),^[10] we are therefore keen to develop milder and better-yielding routes for the direct functionalization of B₂X₄ species.

Substituted monoboranes (RBX₂) can often be conveniently prepared by silicon-boron exchange of the corresponding R-SiMe₃ species with trihaloboranes,^[11] but equivalent reactions with diboranes(4) are limited to just a handful of examples. In the early 1980s, Haubold et al. prepared



Scheme 1. Formation of B-E bonds via silyl halide elimination


Reaction of 2-trimethylsilylbenzyl(dicyclohexyl)phosphine (**2a**), prepared in quantitative yield by treating the corresponding aryllithium compound, 2-LiC₆H₄CH₂PCy₂.Et₂O (**1**), with Me₃SiCl,^[15] with B₂Br₄ at 0 °C led to the formation of sp²-sp³ diborene **3a** in high yield (89%; Scheme 2). The ¹¹B NMR spectrum of **3a** shows the low- and high-field signals expected for the sp² (73.3 ppm) and sp³ (-10.5 ppm) boron atoms, respectively. A broad ³¹P NMR signal at 2.6 ppm is also indicative of coordination to boron. X-ray diffraction provided confirmation of the expected structure (Figure 1), with the B-B bond length of 1.678(9) Å slightly shorter than published sp²-sp³ diboranes,^[16] possibly as a result of steric constraints imposed by the chelating benzylphosphine group. Addition of a further equivalent of **2a** to **3a** resulted in no reaction at room temperature, while higher temperatures (80 °C) resulted in slow, unselective conversion to several species, none of which could be identified. The corresponding reaction with 2-trimethylsilylbenzyl(di-*tert*-butyl)phosphine (**2b**) also yields the equivalent sp²-sp³ diborene, **3b**. In this case, an intermediate species is observed when the reaction is performed at room temperature; observation of signals in the ¹¹B NMR spectrum at 68.6 and -5.7 ppm, a broad ³¹P signal at 21.6 ppm and a signal at 0.31 ppm in the ¹H NMR spectrum for the SiMe₃ group indicate formation of adduct **A**, which can be completely converted to **3b** by heating to 60 °C for 2 h. This observation explains the regioselectivity of the formation of **3a** and **3b**. Silicon-boron exchange reactions of haloboranes are, to the best of our knowledge, limited to sp²-hybridized boranes, hence the selective formation of the 1,2-isomer.^[17] The structure of **3b** was also determined by X-ray crystallography (Figure 1), with the B-B bond length (1.681(9) Å) and other parameters not differing significantly from those in **3a**.

[a] Dr. T. E. Stennett, I. Vollert, Prof. Dr. H. Braunschweig
Institute for Inorganic Chemistry
Julius-Maximilians-Universität Würzburg
Am Hubland, 97074 Würzburg (Germany)

and
Institute for Sustainable Chemistry & Catalysis with Boron
Julius-Maximilians-Universität Würzburg
Am Hubland, 97074 Würzburg (Germany)

E-mail: h.braunschweig@uni-wuerzburg.de

[b] J. D. Mattock, Dr. A. Vargas
Department of Chemistry, School of Life Sciences
University of Sussex
Brighton, BN1 9QJ, Sussex (UK)
E-mail: alfredo.vargas@sussex.ac.uk



Scheme 2. Synthesis of sp^2 - sp^3 diboranes **3a** and **3b** via Me_3SiBr elimination.






Figure 1. Crystal structures of **3a** (top), **3b** (middle) and **4** (bottom) with atomic displacement ellipsoids at the 50% probability level. Hydrogen atoms have been omitted for clarity. Selected bond lengths in Å: **3a**: B1-B2 1.678(9), P1-B1 1.954(8), C1-B2 1.54(1). **3b**: B1-B2 1.681(9), P1-B1 1.956(7), C1-B2 1.563(9). **4**: B1-B2 1.737(3), P1-B1 1.952(2), P2-B1 1.986(2), C1-B2 1.630(3).

Treatment of **3a** with small Lewis bases, L, led to their coordination to the remaining sp^2 boron atom and formation of sp^3 - sp^3 diboranes (L = PMe_3 (**4**), N,N-diisopropylimidazol-2-ylidene (*i*Pr) (**5**), Scheme 3), as indicated by a shift of the low-field resonance for the sp^2 boron atom to high field. The ^{31}P NMR spectrum of **4** shows two broad, mutually coupled doublets at -7.9 and -12.4 ppm ($^3J_{PP} \approx 40$ Hz), whereas the sole signal of **5** appears at -10.5 ppm. The structure of **4** was confirmed by X-



ray diffraction (Figure 1). The B-B bond length displays the expected slight elongation, to 1.737(3) Å, with the structural parameters being otherwise unremarkable.



Scheme 3. Synthesis of unsymmetrical diborenes **6** and **7**.

Reaction of **4** with KC_8 in benzene gave a stark color change to dark yellow and formation of a new species with a single, broad signal in its ^{11}B NMR spectrum at 20.4 ppm, within the range of known diborenes. After two days conversion was complete, and after workup diborene **6** was isolated as a yellow solid in 68% yield. Single crystals suitable for X-ray diffraction were grown from hexane, allowing confirmation of the formation of **6** (Figure 2, Table 1). At 1.563(3) Å, the B-B distance is in the expected range for a boron-boron double bond, while the P-B bonds are both contracted by roughly 0.07 Å compared to **4**. Interestingly, the benzylphosphine chelate appears to cause a distortion of the substituents from ideal double-bond geometry, as the P1-B1-B2-C1 unit has a torsion angle of 17.3°. A similar procedure resulted in the reduction of **5** to diborene **7**, isolated as an orange solid in 34% yield, which is the first reported non-allenic diborene stabilized by two different classes of Lewis base – a phosphine and an NHC. In this case, the ^{11}B NMR signals at 30.0 and 12.5 ppm are at the extreme ends of the range for previously reported diborenes, and therefore suggest a reasonable degree of polarity of the B-B bond. The structure of **7** was also confirmed by X-ray diffraction (Figure 2), and the B-B double bond (1.562(6) Å) found to be the same length as that in **6**, within error.

Figure 2. Crystal structures of **6** (left), **7** (middle) and **8** (right) with atomic displacement ellipsoids at the 50% probability level. Hydrogen atoms have been omitted for clarity. Key bond lengths are displayed in Table 1.

d/Å	6	7	8
B1-B2	1.563(3)	1.562(6)	1.558(7)
P1-B1	1.879(2)	1.863(5)	1.853(4)
P2-B	1.909(2)	-	1.869(4)
C1-B2	1.587(3)	1.573(5)	1.576(6)
C2-B2	-	1.582(6)	-

Table 1. Selected bond lengths (Å) for compounds **6-8**.

Calculations based on the Kohn-Sham Density Functional Theory (DFT) at the OLYP/TZ2P level were conducted to further understand the observed results and the underlying rationales. Figure 3 shows the frontier orbitals of **6** and **7** and Table 2 displays the calculated bond lengths, bond orders and partial charges of the two structures in the gas phase. While both species largely display diborene characteristics in terms of electronic structure, as shown by their frontier orbitals and the B-B bond length, the phosphine-stabilized compound **6** exhibits a slightly higher B-B bond order than **7**, and a less polarized B-B unit. In addition, this unequal charge distribution is inverted, i.e. in **6** the B1 centre carries the lesser negative charge, in contrast to the situation in **7**. These observations suggest that there is a higher C→B2→B1 charge flow compared to the P→B2→B1 path. The LUMO of both compounds has a large contribution from the π -system of the arene moiety, which is held close to the B=B plane by the chelate. This conjugation is noteworthy as homoarene substituents in previously reported diborenes sit exclusively perpendicular to the B=B bond. Both the HOMO and LUMO are found to be at lower energies in phosphine derivative **6** than in **7** due to the higher σ -donor strength of the NHC.

	6	7
d(BB)/Å	1.593	1.595
d(B2-C1)/Å	1.588	1.576
Bond order(BB)	1.52	1.48
q(B1)	-0.129	-0.149
q(B2)	-0.155	-0.113
q(Br)	-0.141	-0.153

Table 2. Calculated properties of compounds **6** and **7**.





Figure 3. Frontier molecular orbitals of compounds **6-8**.

Unsymmetrical diborenes remain very much a novelty, with the only reported examples coming from the group of Kinjo in the past year,^[5a, 6] and their behavior is as yet largely unexplored. We initially tested the thermal stability of **6** and **7**. Heating **6** to 80 °C in benzene led to a slow conversion to a new species displaying a broad signal at 52.2 ppm and a pseudo-triplet at -21.1 ppm in the ¹¹B NMR spectrum. Broad multiplets at 23.3 ppm and -11.9 ppm in the ³¹P NMR spectrum indicated that both phosphines remained coordinated to boron. After 2 d conversion was complete, and single crystals were grown by concentration of the reaction solution. X-ray diffraction revealed the product to be **8** (Scheme 4, Figure 2), in which an exchange of the diborene substituents had occurred, with both phosphines now bonded to the same boron atom. The B-B bond distance of 1.558(7) Å remains in the range for a boron-boron double bond, unchanged from **6**, while the B-P distances are slightly shorter than those in **6**. Heating compound **7** in benzene resulted in an even slower, unselective reaction, with no NMR signals comparable to those for **8** observed. As yet we have been unable to identify any of the products.



Scheme 4. Thermal rearrangement of **6** into **8**.

The nature of the boron-boron bond in **8** is not immediately clear; a classical π -bond would necessitate a formal negative charge on B2 and a positive charge on B1 (Scheme 4). Alternatively, the compound can be viewed as a boryl-substituted borylene, in which the monovalent B1 boron atom carries a lone pair of electrons. Although borylenes were formerly merely postulated as fleeting intermediates, isolable, metal-free borylene compounds stabilized by two (or one) neutral donors are now firmly established.^[18] A recent computational study by Ghorai and Jemmis predicted that a simplified diborene, B₂Br₂(PH₃)₂, would be more stable as its

gem isomer than as its *cis* and *trans* isomers (derivatives of both of which have been realized experimentally), while its NHC derivative favors the *trans* conformation.^[19] The authors proposed a borylborylene type structure, involving a large donor-acceptor contribution to the boron-boron bond, from the P₂B unit to the BBr₂ boron atom. Having isolated such a compound, we again turned to DFT for more insight into the formation and structure of **8**.

The calculated thermal conversion of **6** to **8** corresponds to a stabilization by 3.90 kcal/mol in terms of total electronic energy in the gas phase, and by 5.08 kcal/mol in toluene (within the conductor-like screening model). The more pronounced lowering in energy in the presence of a polarizable medium suggests a highly polarized B-B core in the product **8**. Table 3 shows the comparison of **6** and **8** in terms of selected parameters. The B-B bond order is somewhat lower in **8** (1.406) than in **6** (1.544), but nevertheless indicates significant double-bond character. Close inspection of the calculated partial charges ($q_{B1} = -0.243$, $q_{B2} = -0.038$) indicates a highly charge-separated state in **8**, in which the polarity is inverted with respect to **6** and the negative charge localized on B1. This analysis favors the borylborylene configuration proposed by Jemmis,^[19] where B1 carries a lone pair. Furthermore, the frontier molecular orbitals shown in Figure 3 indicate a marked difference in the HOMO compared to the *trans*-diborenes. While in both compounds **6** and **7**, the HOMO describes a typical (albeit slightly distorted) diborene π -system, in compound **8** the HOMO, though still bearing an overall π symmetry, is more centered on the P1-B1-P2 unit. The LUMO is of boron-boron π -antibonding character with significant delocalization into the arene π -system. While the HOMO-LUMO gaps of the compounds are of similar magnitude, both of these orbitals are stabilized by the rearrangement, the HOMO to a slightly greater extent. The HOMO-1 of all three compounds represents the B-B sigma bonding interaction. Compound **8** bears some resemblance to a diboraallene compound recently reported by Kinjo and co-workers;^[6] however, in that case, the HOMO is localized on the B=C double bond, with the lower-lying HOMO-1 describing the boron-boron π -interaction.

	8 (gas)	6 (gas)	8 (solv)	6 (solv)
$d(BB)$	1.618	1.593	1.612	1.593
Bond order(BB)	1.349	1.526	1.406	1.544
$q(B1)$	-0.243	-0.129	-0.240	-0.133
$q(B2)$	-0.019	-0.155	-0.038	-0.156
$q(Br)$	-0.131	-0.141	-0.169	-0.167

Table 3. Calculated properties of compounds **6** and **8**.




Figure 4. The first two deformation densities of compounds **6** and **8**, with negative charge flow depicted from red lobes to blue.

Additionally, the B-B interaction in **6** and **8** was estimated within the Energy Decomposition Analysis formalism (EDA) under the assumption of equal B1-P1 bond energies in **6** and **8**. Table 4 gives the values of the different bonding components. The total interaction energy is of slightly larger magnitude in **6**. Both the orbital interactions and repulsive Pauli components of the interaction are larger in **8** – this can be ascribed to borylene lone pair donation to the boryl centre, resulting in significant π - π interactions. The electrostatic component is larger in **8**, reflecting the more polar nature of the diboron unit. Furthermore, the deformation densities (Figure 4), which depict the charge flow upon interaction of B1 in **6** and **8** with the rest of the corresponding molecule, indicate significant similarities and differences between the two isomers. In both **6** and **8**, the main deformation density $\Delta\rho_1$ shows charge accumulation towards the boron atom stabilized by PMe₃. However, in **8**, one can see that charge flows to B1 not only from B1-P1 (dative) but also from B2 (covalent) – confirming the borylene-like nature of B1. The B2→B1 flow in **6** is only a secondary deformation density ($\Delta\rho_2$) and is of π -like symmetry, as expected for diborenes; in **8**, there is also some charge flow ($\Delta\rho_2$) towards the B2-B1 bond of σ -like character, further supporting the assignment of B1 as a doubly dative-stabilized borylene.

	8 (gas)	6 (gas)
E_{el}	-363.77	-282.71
E_{orb}	-612.09	-406.69
E_{Pauli}	810.15	498.72
E_{int}	-165.70	-190.68

Table 4. The interaction energy decomposition (kcal/mol) in **6** (fragment 1 = B1-Br) and **8** (fragment 1 = B1-PMe₃) with fragment 2 = rest of the molecule.

To summarize, we have used boron-silicon metathesis at B₂Br₄ to develop a route to unsymmetrical diborenes based upon a chelating benzylphosphine linker, with either a phosphine or an N-heterocyclic carbene as the second stabilizing ligand. DFT calculations confirm the slight polarity of these compounds. We have also reported thermodynamic rearrangement of one of these unsymmetrical *trans*-diborenes to an unprecedented *gem*-diborene containing a polar boron-boron bond. On the basis of DFT calculations, which reveal large electrostatic and orbital contributions to the bonding, we propose that the compound is best described as a borylborylene.

Acknowledgements

The authors gratefully acknowledge the European Research Council (ERC) under the European Union Horizon 2020 Research and Innovation Program (grant agreement no. 669054) for funding. A.V. thanks the University of Sussex for financial support.

Keywords: boron • borylene • multiple bonds • rearrangement • DFT calculations

- [1] Y. Wang, B. Quillian, P. Wei, C. S. Wannere, Y. Xie, R. B. King, H. F. Schaefer, III, P. v. R. Schleyer, G. H. Robinson, *J. Am. Chem. Soc.* **2007**, *129*, 12412-12413.
- [2] a) M. Arrowsmith, H. Braunschweig, T. E. Stennett, *Angew. Chem. Int. Ed.* **2017**, *56*, 96-115; b) P. Bissinger, H. Braunschweig, A. Damme, T. Kupfer, A. Vargas, *Angew. Chem. Int. Ed.* **2012**, *51*, 9931-9934; c) H. Braunschweig, R. D. Dewhurst, *Organometallics* **2014**, *33*, 6271-6277.
- [3] a) H. Braunschweig, I. Krummenacher, C. Lichtenberg, J. D. Mattock, M. Schafer, U. Schmidt, C. Schneider, T. Steffenhagen, S. Ullrich, A. Vargas, *Angew. Chem. Int. Ed.* **2017**, *56*, 889-892; b) H. Braunschweig, R. D. Dewhurst, C. Hoerl, A. K. Phukan, F. Pinzner, S. Ullrich, *Angew. Chem. Int. Ed.* **2014**, *53*, 3241-3244.
- [4] a) P. Bissinger, H. Braunschweig, M. A. Celik, C. Claes, R. D. Dewhurst, S. Endres, H. Kelch, T. Kramer, I. Krummenacher, C. Schneider, *Chem. Commun.* **2015**, *51*, 15917-15920; b) P. Bissinger, H. Braunschweig, A. Damme, T. Kupfer, I. Krummenacher, A. Vargas, *Angew. Chem. Int. Ed.* **2014**, *53*, 5689-5693.
- [5] a) W. Lu, Y. X. Li, R. Ganguly, R. Kinjo, *J. Am. Chem. Soc.* **2017**, *139*, 5047-5050; b) H. Braunschweig, R. D. Dewhurst, K. Hammond, J. Mies, K. Radacki, A. Vargas, *Science* **2012**, *336*, 1420-1422.
- [6] W. Lu, Y. X. Li, R. Ganguly, R. Kinjo, *Angew. Chem. Int. Ed.* **2017**, *56*, 9829-9832.
- [7] H. Braunschweig, T. Dellermann, W. C. Ewing, T. Kramer, C. Schneider, S. Ullrich, *Angew. Chem. Int. Ed.* **2015**, *54*, 10271-10275.
- [8] M. Arrowsmith, J. Böhnke, H. Braunschweig, M. Celik, C. Claes, W. Ewing, I. Krummenacher, K. Lubitz, C. Schneider, *Angew. Chem. Int. Ed.* **2016**, *55*, 11271-11275.
- [9] H. Klusik, A. Berndt, *J. Organomet. Chem.* **1982**, *234*, C17-C19.
- [10] a) M. Arrowsmith, J. Böhnke, H. Braunschweig, A. Deissenberger, R. D. Dewhurst, W. C. Ewing, C. Hörl, J. Mies, J. H. Muessig, *Chem. Commun.* **2017**, *53*, 8265-8267; b) H. Nöth, H. Pommerening, *Chem. Ber.-Recueil* **1981**, *114*, 398-399.
- [11] a) W. Haubold, J. Herdtle, W. Gollinger, W. Einholz, *J. Organomet. Chem.* **1986**, *315*, 1-8; b) D. Kaufmann, *Chem. Ber.-Recueil* **1987**, *120*, 853-854; c) D. Kaufmann, *Chem. Ber.-Recueil* **1987**, *120*, 901-905.
- [12] W. Haubold, J. Hrebicek, G. Sawitzki, *Z. Naturforsch. B.* **1984**, *39*, 1027-1031.
- [13] P. Greiwe, A. Bethäuser, H. Pritzkow, T. Kuhler, P. Jutzi, A. Siebert, *Eur. J. Inorg. Chem.* **2000**, 1927-1929.
- [14] After the preparation of this manuscript, Kinjo and co-workers published an example of a *gem*-diborene as a result of a hydrogen atom transfer from a diboraallene: W. Lu, Y. Li, R. Ganguly, R. Kinjo, *J. Am. Chem. Soc.* **2018**, *140*(4), 1255-1258.
- [15] H. P. Abicht, K. Issleib, *Z. Anorg. Allg. Chem.* **1976**, *422*, 237-242.
- [16] a) H. Braunschweig, A. Damme, J. O. C. Jiménez-Halla, T. Kupfer, K. Radacki, *Angew. Chem. Int. Ed.* **2012**, *51*, 6267-6271; b) R. D. Dewhurst, E. C. Neeve, H. Braunschweig, T. B. Marder, *Chem. Commun.* **2015**, *51*, 9594-9607; c) S. Pietsch, E. C. Neeve, D. C. Apperley, R. Bertermann, F. Y. Mo, D. Qiu, M. S. Cheung, L. Dang, J. B. Wang, U. Radius, Z. Y. Lin, C. Kleeberg, T. B. Marder, *Chem. Eur. J.* **2015**, *21*, 7082-7098.
- [17] a) D. A. Singleton, S. W. Leung, *J. Organomet. Chem.* **1997**, *544*, 157-161; b) A. Boussonnière, X. C. Pan, S. J. Geib, D. P. Curran, *Organometallics* **2013**, *32*, 7445-7450.
- [18] a) R. Kinjo, B. Donnadiou, M. A. Celik, G. Frenking, G. Bertrand, *Science* **2011**, *333*, 610-613; b) M. Soleilhavoup, G. Bertrand, *Angew. Chem. Int. Ed.* **2017**, *56*, 10282-10292; c) H. Braunschweig, R. D. Dewhurst, F. Hupp, M. Nutz, K. Radacki, C. W. Tate, A. Vargas, Q. Ye, *Nature* **2015**, *522*, 327-330; d) M. Arrowsmith, D. Auerhammer, R. Bertermann, H. Braunschweig, G. Bringmann, M. A. Celik, R. D. Dewhurst, M. Finze, M. Grune, M. Hallmann, T. Hertle, I. Krummenacher, *Angew. Chem. Int. Ed.* **2016**, *55*, 14462-14466.
- [19] S. Ghorai, E. D. Jemmis, *Chem. Eur. J.* **2017**, *23*, 9746-9751.

Entry for the Table of Contents (Please choose one layout)

Layout 1:

COMMUNICATION

Text for Table of Contents

((Insert TOC Graphic here))

Author(s), Corresponding Author(s)*

Page No. – Page No.

Title


Layout 2:

COMMUNICATION

Tom E. Stennett, James D. Mattock,
Ivonne Vollert, Alfredo Vargas* and
Holger Braunschweig*

Page No. – Page No.

Title



Unsymmetrical diborenes were prepared by reduction of cyclic diborenes(4) containing a benzylphosphine bridge. A doubly phosphine-stabilized diborene underwent an unprecedented, thermally induced rearrangement to form a borylborylene.