

Evaluation of performance and perceived utility of mental healthcare indicators in routine health information systems in five low- and middle-income countries

Article (Published Version)

Jordans, Mark, Chisholm, Dan, Semrau, Maya, Gurung, Dristy, Abdulmalik, Jibril, Ahuja, Shalini, Mugisha, James, Mntambo, Ntokozo, Kigozi, Fred, Petersen, Inge, Shidhaye, Rahul, Upadhaya, Nawaraj, Lund, Crick, Thornicroft, Graham and Gureje, Oye (2019) Evaluation of performance and perceived utility of mental healthcare indicators in routine health information systems in five low- and middle-income countries. *BJPsych Open*, 5 (5). a70 1-9. ISSN 2056-4724

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/82623/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Evaluation of performance and perceived utility of mental healthcare indicators in routine health information systems in five low- and middle-income countries

Mark Jordans, Dan Chisholm, Maya Semrau, Dristy Gurung, Jibril Abdulmalik, Shalini Ahuja, James Mugisha, Ntokozo Mntambo, Fred Kigozi, Inge Petersen, Rahul Shidhaye, Nawaraj Upadhaya, Crick Lund, Graham Thornicroft and Oye Gureje

Background

In most low- and middle-income countries (LMIC), routine mental health information is unavailable or unreliable, making monitoring of mental healthcare coverage difficult. This study aims to evaluate a new set of mental health indicators introduced in primary healthcare settings in five LMIC.

Method

A survey was conducted among primary healthcare workers ($n = 272$) to assess the acceptability and feasibility of eight new indicators monitoring mental healthcare needs, utilisation, quality and payments. Also, primary health facility case records ($n = 583$) were reviewed by trained research assistants to assess the level of completion (yes/no) for each of the indicators and subsequently the level of correctness of completion (correct/incorrect – with incorrect defined as illogical, missing or illegible information) of the indicators used by health workers. Assessments were conducted within 1 month of the introduction of the indicators, as well as 6–9 months afterwards.

Results

Across both time points and across all indicators, 78% of the measurements of indicators were complete. Among the best performing indicators (diagnosis, severity and treatment), this was significantly higher. With regards to correctness, 87% of all completed indicators were correctly completed. There was a trend towards improvement over time. Health workers'

perceptions on feasibility and utility, across sites and over time, indicated a positive attitude in 81% of all measurements.

Conclusion

This study demonstrates high levels of performance and perceived utility for a set of indicators that could ultimately be used to monitor coverage of mental healthcare in primary healthcare settings in LMIC. We recommend that these indicators are incorporated into existing health information systems and adopted within the World Health Organization Mental Health Gap Action Programme implementation strategy.

Declaration of interest

None.

Keywords

Mental healthcare; indicators; primary healthcare; low- and middle-income settings; health information system.

Copyright and usage

© The Royal College of Psychiatrists 2019. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

Globally, and especially in low- and middle-income countries (LMIC), there is a large treatment gap for mental disorders, as indicated by the lack of service for a great majority of people in need of mental healthcare.¹ Integrating mental healthcare into primary healthcare has been recommended as a way of bridging this treatment gap by offering more accessible, holistic and less stigmatising services to people in need.² The World Health Organization (WHO) launched the Mental Health Gap Action Programme Intervention Guide (mhGAP-IG) in 2010 (and a second version in 2016), to provide evidence-based clinical guidance for the assessment and management of priority mental, neurological and substance use disorders in LMIC non-specialist settings.³ Building on this development, and in order to facilitate the process of integration, a broader health system strengthening approach involving policy changes, sustainable financing mechanisms and workforce development is required.

One such system requirement is a well-functioning routine health information system that ensures the production, analysis, dissemination and use of reliable and timely information related to service delivery.⁴ Moreover, in order for the much-needed investments in scaling up of mental health services to be efficiently used, comprehensive and accurate information on the availability, utilisation and equity of such services is required. However, health

information systems to support actions in mental healthcare are weak in LMIC. Most routine information systems do not have any, or have very limited, indicators related to mental healthcare. When information is collected, it often produces unreliable data.⁵ The aim of this study was to evaluate the performance and perceived utility of a new set of indicators for routine monitoring of mental healthcare within primary healthcare settings in five LMIC.

Method

Setting

The study was conducted in five LMIC in Africa and Asia participating in the Emerald programme (India, Nepal, Nigeria, South Africa and Uganda). The Emerald program aimed to improve mental health outcomes by generating evidence and capacity to enhance health system performance in LMIC.⁶

Emerald programme

As a way of responding to the lack of indicators to monitor mental healthcare, the Emerald programme developed, through a Delphi study, a set of indicators that can be used to appraise mental

healthcare needs, utilisation, outcomes and costs.⁷ From an initial set of 52 generated items, the eight most highly prioritised indicators were selected and operationalised through consultative workshops in each country setting convened to develop data collection formats suitable for use by frontline health workers. The eight indicators included were: diagnosis (as multiple-choice response options, which included depression, alcohol use disorder, schizophrenia and epilepsy, as well as other disorders depending on priorities per country team), exact diagnosis (as open response format), severity, functioning, administered treatment, referral, follow-up and payment for services (the latter six items all had multiple response options, including none or not relevant). These formats were subsequently integrated into district-level mental healthcare programmes, as part of ongoing research programmes evaluating the integration of mental healthcare into primary healthcare.^{8,9} (See supplementary File 1, available at <https://doi.org/10.1192/bjp.2019.22>, for the standard form, which was adapted in each setting.)

Instruments

First, a structured questionnaire was developed for the purpose of this study, to assess health workers' perceptions of the utility (i.e. acceptability and feasibility) of using the data collection format for the new indicators. The questionnaire consisted of 14 questions, which were administered through interviews conducted by trained research assistants. Second, a health information case record review form was developed to assess the performance of the newly introduced indicators based on observation of the completed formats. A trained research assistant checked all case records, first assessing whether the completion of each of the indicators (i.e. level of completion (response options: 'yes/no')). Subsequently, for each completed indicator, the research assistant assessed the level to which completed information was correctly or incorrectly completed (i.e. level of correctness of completion (response options: 'correct/incorrect'), with the latter defined as 'illogical completion/missing information/illegible information'). The form consisted of eight items and the ratings of its completion and correctness of completion were made by trained research assistants.

Sample

See Table 1 for an overview of the number of interviews and case record reviews across sites and time-points. The structured interviews were administered to 272 health workers who had been trained to provide services and to use the new data collection forms. The sample was selected from primary healthcare facilities (all countries) and a few district hospitals (India and Uganda). Selection was done either randomly (in Nepal, among the 205 trained health workers, 177 were still in post, among whom 50 were randomly selected; in Nigeria, two health workers were randomly selected among the 4–6 trained health workers per facility) or purposively (in South Africa, staff were selected based on availability and willingness until the target sample was reached; in Uganda and India, the health facility in charge approached health workers involved in record keeping or mental health services, respectively). The health information records ($n = 586$) assessed for completion and correctness were selected randomly in all of

the countries. Randomisation of the case records was done by: (a) randomly selecting a specified number of facilities and (b) randomly selecting a number of case records within the selected facility.

Procedure

The interviews with health workers and the health information record reviews were performed 2–4 weeks after the new forms were introduced and had started to be used in the clinics. We chose the period of 2–4 weeks to allow some time for proper adoption of the procedure post training, including the printing and distribution of the required registers/forms. This time frame was to allow the staff some time to establish familiarity with the new forms before the interviews/assessments were conducted at baseline. A second wave of interviews and record reviews were conducted with the same health workers and in the same health facilities 6–9 months after the introduction of the forms, to assess whether there was improvement, deterioration or constancy in the scores over time. Interviews and record reviews were conducted by research assistants who had received an average of 5 days training at each of the participating sites.

Data analysis and management

Data from questionnaires and record review forms (for both time points; T1 and T2) were entered into SPSS and analysed using descriptive statistics. The calculation of the proportion of completion of indicators was based on the total number of cases; correctness was calculated based on the completed indicators and was determined by whether the information entered in the data form was assessed by the research assistant to be correct. Thus, for example, if the indicator slot for 'diagnosis' was filled, a rating was made to indicate that the information was indeed collected (completion) and whether the information filled into the slot was indeed a relevant diagnosis (correctness). Data collection took place between August 2015 and October 2017.

Ethics

Ethical approval for this study was obtained from King's College London, the World Health Organization and the institutional review boards of each of the participating sites. Subsequently, permission was obtained from the appropriate Department/Ministry of Health officials, as well as from the healthcare workers and health facility managers. Written informed consent was sought and obtained from all study participants. The facility managers were approached by research assistants for their participation in the study and for permission to access the health information records.

Results

Across the five sites, 272 health workers were interviewed and 586 case records were reviewed. Table 2 shows the performance ratings of the new indicators with regard to levels of completion and correctness of completion. Overall, high levels of completion and of correctness were achieved for most of the indicators at

Table 1 Overview of sample

	India (N)	Nepal (N)	Nigeria (N)	South Africa (N)	Uganda (N)	Total
T1 surveys of health workers	16	50	36	24	26	152
T2 surveys of health workers	9	45	36	18	12	120
T1 record reviews	61	95	45	32	66	299
T2 record reviews	74	58	43	40	69	284

T1 refers to 2–4 weeks after introduction of the new indicators within primary healthcare settings; T2 refers to 6–9 months after the introduction.

Table 2 Levels of completion and correctness of completion of mental health indicators in routine health information systems across five LMICs

			Diagnosis N (%)	Exact diagnosis N (%)	Severity N (%)	Functioning N (%)	Treatment administered N (%)	Referral N (%)	Follow-up N (%)	Payment N (%)
India	Completion	T1 N = 61	61 (100)	49 (80.0)	61 (100)	n/a	60 (98.0)	2 (100) ^a	41 (100) ^a	n/a
		T2 N = 74	65 (87.8)	63 (85.1)	29 (39.1)	n/a	65 (87.8)	4 (100) ^a	3 (100) ^a	n/a
	Correctness ^b	T1	60 (98.0)	46 (94.0)	60 (98.0)	n/a	42 (70.0)	2 (100) ^a	29 (100) ^a	n/a
		T2	64 (98.4)	55 (87.3)	29 (100)	n/a	64 (98.4)	4 (100) ^a	3 (100) ^a	n/a
Nepal	Completion	T1 N = 95	82 (86.3)	27 (28.4)	81 (85.3)	45 (47.4)	76 (80.0)	16 (16.8)	55 (57.9)	35 (36.8)
		T2 N = 58	57 (98.3)	43 (74.1)	56 (96.6)	38 (65.5)	55 (94.8)	32 (55.2)	51 (87.9)	37 (63.8)
	Correctness ^b	T1	79 (96.3)	24 (88.9)	81 (100)	28 (62.2)	71 (93.4)	15 (93.7)	52 (94.5)	29 (82.8)
		T2	57 (100)	43 (100)	56 (100)	38 (100)	55 (100)	32 (100)	51 (100)	37 (100)
Nigeria	Completion	T1 N = 45	36 (80)	n/a	34 (75.6)	34 (75.6)	45 (100)	45 (100)	27 (60)	39 (86.7)
		T2 N = 43	43 (100)	n/a	42 (97.7)	37 (86.0)	42 (97.7)	42 (97.7)	43 (100)	43 (100)
	Correctness ^b	T1	25 (69.4)	n/a	32 (94.1)	32 (94.1)	44 (97.8)	42 (93.3)	22 (81.5)	39 (100)
		T2	43 (100)	n/a	41 (97.6)	36 (97.3)	41 (97.6)	39 (92.9)	43 (100)	43 (100)
South Africa	Completion	T1 N = 32	28 (87.5)	31 (96.9)	30 (93.8)	n/a	20 (62.5)	31 (96.9)	32 (100)	n/a
		T2 N = 40	40 (100)	40 (100)	40 (100)	n/a	40 (100)	40 (100)	40 (100)	n/a
	Correctness ^b	T1	28 (100)	31 (100)	30 (100)	n/a	20 (100)	31 (100)	31 (96.9)	n/a
		T2	40 (100)	40 (100)	40 (100)	n/a	40 (100)	40 (100)	40 (100)	n/a
Uganda	Completion	T1 N = 66	65 (98.5)	43 (65.2)	n/a	64 (97.0)	62 (95.4)	22 (33.9)	59 (89.4)	42 (63.6)
		T2 N = 69	69 (100)	62 (89.9)	n/a	68 (98.6)	69 (100)	27 (39.1)	63 (91.3)	22 (31.9)
	Correctness ^b	T1	62 (95.4)	23 (53.5)	n/a	58 (95.1)	32 (50.0)	7 (35.0)	18 (31.0)	0
		T2	61 (88.4)	37 (59.7)	n/a	45 (66.2)	52 (75.4)	9 (33.3)	45 (71.4)	8 (36.4)

a. The response options and the denominators for these items were different from the other countries; these are therefore not included in the summaries of results.

b. Correctness refers to level of completed indicators that was correctly completed (as opposed to 'incorrect completion').

both time points. The combined completion and correctness ratings for both time points were especially high for diagnosis (94 and 95%, respectively), severity (83 and 99%, respectively) and treatment (92 and 86%, respectively). These scores were slightly lower for exact diagnosis (72 and 84%, respectively), functioning (76 and 83%, respectively) and follow-up (83 and 82%, respectively). To give an indication of the samples of cases for which records were reviewed, the breakdown of diagnoses recorded was as follows (for T1 data): AUD (18% in Nepal, 3% in Uganda, 20% in India, 22% in Nigeria); depression (28% in Nepal, 100% in South Africa, 14% in Uganda, 46% in India, 60% in Nigeria); epilepsy (13% in Nepal, 68% in Uganda, 5% in Nigeria); psychosis (32% in Nepal, 5% in Uganda, 1% in India, 13% in Nigeria); and others (such as anxiety disorders, dementia, development disorders; 9% in Nepal, 10% in Uganda, 33% in India).

The combined completion and correctness for both time points was lowest for referral (60 and 84%, respectively) and payment (58 and 72%, respectively). The lower scores for referral and payments were all clustered within one country setting (i.e. Uganda), where 7/8 measurements points showed results that were <50% positive. The only other trend for such lower scores were for the completion of indicators at T1 in Nepal (4/8 scoring 50% positive). In South Africa scores for completion and correctness were high across indicators and time points with 18/24 measurements showing 100% positives. Furthermore, the T2 correctness measurements for Nepal and Nigeria were all over 90%. These contributed to a cross-country trend for improvement of scores between T1 and T2 for completion of indicators (7/8 showed improvement over time) and for correctness of completion (6/8 showed improvement). There was only a cross-country trend for deterioration over time for completion of the severity indicator.

The utility of the new mental health indicators was assessed through health workers' perceptions of the feasibility and acceptability of using the indicators in routine healthcare settings across the five countries (Table 3). Overall, there were moderate-to-high scores on perceived utility. All but two items had over 80% positive responses (positive responses were defined as the combination of 'agree' and 'strongly agree') across time and countries. These were especially high for perceived importance of routine mental health

indicators (96% positive) motivation to use these indicators (95%), as well as perceived relevance (91%), integration in health workers' daily work (90%), ease of use (86%), confidence to use (84%) and overall satisfaction with the new forms (82%). Perceived usefulness of the indicators in daily practice fell a bit behind, with 74% positive responses. Having insufficient time was the only item where only a minority had a positive response (33%). Indeed, a combined average of 27% of the reports showed that more than 10 min additional time was spent on recording.

Over time, there was an overall reduction of perceived utility. In 15 of the T1–T2 comparisons, there was a reduction of 20% or greater. These were especially seen in South Africa. In only one instance was there an increase of more than 20% in positive evaluation for an item from T1 to T2; this was for the level of confidence in using the formats in India. Finally, Nepal and Nigeria were the two sites with the most positive assessments of feasibility and acceptability by respondents (i.e. both scoring over 80% positive on 14/18 items).

Discussion

A functioning routine mental health information system is required to guide the process of reducing the treatment gap for mental illness globally.¹⁰ The poor availability of data is a hindrance for the achievement of that goal in LMIC.¹¹ A mental health information system aims to improve the effectiveness and efficiency of the mental health service and ensure more equitable delivery. Information that enables managers and service providers to make more informed decisions could aid the delivery of quality care.¹² Given that most LMIC do not have any, or only very few, indicators to monitor mental healthcare as part of their existing Health Management Information System (HMIS), this study aimed to assess the performance and utility of a new set of indicators in five LMIC (India, Nepal, Nigeria, South Africa and Uganda). This is in line with WHO's Mental Health Action Plan 2013–2020, which sets a target of 80% of countries to routinely collect and report on a core set of mental health indicators within their national information systems by 2020.¹³ The new set of indicators go beyond

Table 3 Health workers' perceptions on using mental health indicators in the routine health information system across five LMICs

Item	Response	India		Nepal		Nigeria		South Africa		Uganda	
		T1	T2	T1	T2	T1	T2	T1	T2	T1	T2
		N = 16 N (%)	N = 9 N (%)	N = 50 N (%)	N = 45 N (%)	N = 36 N (%)	N = 36 N (%)	N = 24 N (%)	N = 18 N (%)	N = 26 N (%)	N = 12 N (%)
Currently involved in routine data collection?	Yes	9	7 (77.8)	n/a	n/a	36 (100)	36 (100)	21 (87.5)	13 (72.2)	n/a	n/a
		7	2 (22.2)			0	0	3 (12.5)	5 (27.8)		
Received training on routine data collection?	Yes	16	9 (100)	43 (86.0)	43 (95.6)	18 (50)	32 (88.9)	8 (33.3)	7 (38.9)	9 (34.6)	5 (41.7)
	No	0	0	7 (14.0)	2 (4.4)	18 (50)	4 (11.1)	16 (66.7)	11 (61.1)	17 (65.4)	7 (58.3)
How many mental illness patients do you see on average in a week?	1 or fewer	3 (18.8)	3 (42.9)	35 (70.0)	18 (40.0)	31 (86.1)	34 (94.4)	7 (29.2)	5 (27.8)	3 (11.5)	0
	2–4 per week	8 (50.0)	3 (42.9)	12 (24.0)	23 (51.1)	5 (13.9)	2 (5.6)	7 (29.2)	10 (55.6)	14 (53.9)	8 (66.7)
	5–7 per week	3 (18.8)	0	2 (4.0)	2 (4.4)	0	0	2 (8.3)	2 (11.1)	4 (15.4)	4 (33.3)
	8–10 per week	1 (6.3)	0	0	1 (2.2)	0	0	4 (16.7)	0	3 (11.5)	0
	>10 per week	1 (6)	1 (14.3)	1 (2.0)	1 (2.2)	0	0	4 (16.7)	1 (5.6)	2 (7.7)	0
How much time do you think you spend, on average, with each patient who has mental health problems?	<5 min	0	1 (14.3)	0	0	7 (19.4)	14 (38.9)	0	0	3 (11.5)	0
	5–10 min	2 (12.5)	1 (14.3)	6 (12.0)	12 (26.7)	9 (25)	7 (19.4)	4 (16.7)	2 (11.1)	10 (38.5)	4 (33.3)
	11–20 min	0	1 (14.3)	18 (36.0)	16 (35.6)	6 (16.7)	6 (16.7)	10 (41.7)	5 (27.8)	10 (38.5)	5 (41.7)
	21–30 min	2 (12.5)	0	13 (26.0)	10 (22.2)	7 (19.4)	5 (13.8)	9 (37.5)	7 (38.9)	2 (7.7)	3 (25)
	>30 min	12 (75.0)	4 (57.1)	13 (26.0)	7 (15.6)	7 (19.4)	4 (11.1)	1 (4.2)	4 (22.2)	1 (3.9)	0
How much time do you spend on average recording information for any patient?	<5 min	1 (6.3)	0	21 (42.0)	24 (53.3)	10 (27.8)	11 (30.6)	0	1 (5.6)	7 (28.0)	5 (41.7)
	5–10 min	2 (12.5)	4 (57.1)	26 (52.0)	15 (33.3)	16 (44.4)	14 (38.9)	7 (29.2)	9 (50.0)	14 (56)	5 (41.7)
	11–20 min	7 (43.8)	1 (14.3)	3 (6.0)	5 (11.1)	9 (25)	7 (19.4)	13 (54.2)	5 (27.8)	3 (12.0)	2 (16.7)
	>20 min	6 (37.5)	2 (28.6)	0	1 (2.2)	1 (2.8)	4 (11.1)	4 (16.7)	3 (16.7)	1 (4.0)	0
How much additional time was needed using the new format?	<5 min	0	1 (14.3)	13 (26.0)	8 (17.8)	13 (36.1)	9 (25)	12 (50.0)	2 (11.1)	14 (53.9)	4 (33.3)
	5–10 min	6 (37.5)	4 (57.1)	24 (48.0)	30 (66.7)	8 (22.2)	15 (41.7)	11 (45.8)	11 (61.1)	10 (38.5)	4 (33.3)
	11–20 min	3 (18.8)	0	10 (20.0)	5 (11.1)	6 (16.7)	8 (22.2)	0	1 (5.6)	2 (7.7)	3 (25.0)
	>20 min	7 (43.8)	2 (28.6)	3 (6.0)	2 (4.4)	9 (25)	4 (11.1)	1 (4.2)	4 (22.2)	0	1 (8.3)
How satisfied are you with the new format?	Very dissatisfied	0	0	1 (2.0)	1 (2.2)	1 (2.8)	1 (2.8)	0	1 (5.6)	2 (7.7)	0
	A little dissatisfied	1 (6.3)	2 (28.6)	1 (2.0)	4 (8.9)	0	1 (2.8)	0	1 (5.6)	1 (3.9)	2 (16.7)
	Neutral	3 (18.8)	0	1 (2.0)	0 (0.0)	2 (5.6)	8 (22.2)	1 (4.2)	7 (38.8)	4 (15.4)	0
	A little satisfied	8 (50.0)	2 (28.6)	36 (72.2)	28 (62.2)	11 (30.6)	3 (8.3)	5 (20.8)	4 (22.2)	11 (42.3)	5 (41.7)
	Very satisfied	4 (25.0)	3 (42.9)	11 (22.0)	12 (26.7)	22 (61.1)	23 (63.9)	18 (75.0)	5 (27.8)	8 (30.8)	5 (41.7)
It is possible to use the mental health indicators in routine practice	Strongly disagree	0	0	4 (8.0)	2 (4.4)	1 (2.8)	2 (5.6)	0	2 (11.1)	0	0
	Disagree	1 (6.3)	4 (44.4)	14 (28.0)	3 (6.7)	3 (8.3)	9 (25)	3 (12.5)	0	4 (15.4)	4 (33.3)
	Neutral	0	0	0	3 (6.7)	2 (5.6)	5 (13.9)	0	5 (27.8)	1 (3.9)	0
	Agree	8 (50.0)	3 (33.3)	27 (54.0)	29 (64.4)	18 (50)	15 (41.6)	20 (83.3)	6 (33.3)	19 (73.1)	5 (41.7)
	Strongly agree	7 (43.8)	2 (22.2)	5 (10.0)	8 (17.8)	12 (33.3)	5 (13.9)	1 (4.2)	5 (27.8)	2 (7.7)	3 (25.0)
I do not have enough time to complete additional questions for routine data collection	Strongly disagree	0	0	8 (16.0)	12 (26.7)	10 (27.8)	4 (11.1)	1 (4.2)	3 (16.7)	4 (15.4)	3 (25.0)
	Disagree	1 (96.3)	3 (42.9)	24 (48.0)	20 (44.4)	17 (47.2)	21 (58.3)	13 (54.2)	3 (16.7)	16 (61.5)	4 (33.3)
	Neutral	2 (12.5)	1 (14.3)	1 (2.0)	3 (6.7)	1 (2.8)	0	1 (4.2)	4 (22.2)	1 (3.9)	0
	Agree	10 (62.5)	3 (42.9)	16 (32.0)	7 (15.6)	7 (19.4)	10 (27.8)	9 (37.5)	7 (38.9)	4 (15.4)	5 (41.7)
	Strongly agree	3 (18.8)	0	1 (2.0)	3 (6.7)	1 (2.8)	1 (2.8)	0	1 (5.6)	1 (3.9)	0
Routine mental health data collection is important	Strongly disagree	0	0	0	0	1 (2.8)	1 (2.8)	0	0	0	2 (16.7)
	Disagree	0	1 (11.1)	1 (2.0)	0	1 (2.8)	0	0	0	0	0
	Neutral	0	1 (11.1)	0	0	1 (2.8)	1 (2.8)	0	1 (5.6)	0	0
	Agree	1 (6.3)	6 (66.7)	15 (30.0)	20 (44.4)	23 (63.9)	16 (44.4)	14 (58.3)	10 (55.6)	12 (46.2)	4 (33.3)
	Strongly agree	15 (93.8)	1 (11.1)	34 (68.0)	25 (55.6)	10 (27.8)	18 (50)	10 (41.7)	7 (38.8)	(53.8)	6 (50.0)

use. Still, we saw low ratings on a number of questions, indicating that critical perspectives were also voiced. Second, not all countries used the same or the full set of indicator ratings. The variance between sites is a result of a process of in-country consultations and decision-making with regards to selection and formulation of the indicators. Third, this study did not include the assessment of (a) the quality of information that is reported using these new indicators (i.e. it did not include any validation of the completed information, for example, assessing the accuracy of the diagnosis given), or (b) the use of the information subsequent to collection (i.e. uptake within the larger HMIS system). The assessment of correctness of completion therefore only refers to whether and how the newly introduced formats were completed – a first threshold for adequate use in practice. Fourth, different countries utilised different selection methods for recruiting respondents (random, purposive and convenience), which may detract from the uniformity of the findings. Finally, this study focused exclusively on health facility level indicators directly related to service provision. Other health system indicators (e.g. number of trained personnel, availability of medications) are equally important for adequate monitoring of mental health services.⁷

This study has several implications. First, efforts to close the mental health treatment gap should be matched with efforts to close the information gap associated with mental healthcare in LMIC.¹⁰ This is especially pertinent, as meaningful planning for scaling up mental health services cannot take place in the absence of reliable mental health information to serve as a reliable benchmark of service utilisation and trends over time. Based on the results of this study, a limited set of indicators can be recommended for adoption into routine mental healthcare surveillance, which will be particularly useful when rolling out mhGAP-IG programmes.

Second, in addition to improved planning, the integration of a set of mental health indicators in HMIS will be an important ingredient for maintaining or improving the quality of care. Health workers and/or their supervisors, can use the information to better monitor the adequacy of initiated treatments following diagnosis, treatment adherence or drop-out, as well as patients' improvements over time, or lack thereof.




Third, if the set of indicators that were part of this study were integrated in HMIS and adequately implemented to collect data within routine primary healthcare settings, they could potentially be used to monitor coverage for mental healthcare through the following combination of indicators; *diagnosis*, (*change in*) *severity*, (*change in*) *functioning*, (*frequency of*) *treatment utilisation*, and also in low-resource health systems. This is especially salient as most mental health services information is limited to tertiary care settings¹⁷ and most focuses on health system inputs rather than processes or outcomes.¹² Monitoring the level of financial protection may require a different method, for example, household surveys. Moreover, for calculation of actual coverage, reliable prevalence data are needed (i.e. the denominator), which may be unavailable in many settings.

Fourth, perceived utility is especially high shortly after training. Even though across the board introduction after training seems sufficient to achieve good performance, ongoing supervision and booster trainings may be needed to sustain positive attitudes towards use.¹⁵

Fifth, the next steps should involve the evaluation of using this set of indicators outside of a research programme. While the study took place in a routine care setting, a non-research setting will more closely reflect real-life implementation. Also, future research should evaluate the use of mental healthcare indicators within the HMIS process subsequent to data collection, i.e. processing, analyses, dissemination and use at facility levels, as well as beyond.

In conclusion, this study set out to explore the feasibility of monitoring mental healthcare when provided at scale by non-specialist

health workers. The study demonstrates high levels of performance and utility across time for a set of indicators that could ultimately be used to monitor the coverage of mental healthcare in primary health-care settings in five LMIC and contribute to maintaining or improving the quality of care. We recommend that these indicators be included in existing health information systems, and adopted as part of systems strengthening interventions necessary to facilitate the rolling out of the mhGAP-IG package, and to allow for evidence-based and data-driven decision-making and planning.

Mark Jordans, PhD , Reader, Centre for Global Mental Health, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK; **Dan Chisholm**, PhD, Programme Manager for Mental Health, Regional Office for Europe, World Health Organization, Switzerland; **Maya Semrau**, PhD, Research Fellow, Centre for Global Mental Health, Institute of Psychiatry, Psychology and Neuroscience, King's College London; and Global Health and Infection Department, Brighton & Sussex Medical School, Brighton, UK; **Dristy Gurung**, MA , Research Coordinator, Transcultural Psychosocial Organization, Nepal; **Jibril Abdulmalik**, MD, Senior Lecturer, Department of Psychiatry, University of Ibadan, Nigeria; **Shalini Ahuja**, PhD, Researcher, Public Health Foundation of India & Centre for Global Mental Health, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK; **James Mugisha**, MD, Senior Lecturer, Kyambogo University, Kampala, Uganda & Butabika National Referral and Teaching Mental Hospital, Uganda; **Ntokozi Mtambo**, MA, Researcher, University of Kwazulu-Natal, South Africa; **Fred Kigozi**, PhD, Senior Consultant Psychiatrist and Researcher, Butabika National Referral and Teaching Mental Hospital, Uganda; **Inge Petersen**, PhD, Research Professor and Director, Centre for Rural Health, School of Nursing and Public Health, University of Kwazulu-Natal, South Africa; **Rahul Shidhaye**, MD, Clinical Psychiatrist, Public Health Foundation of India, India; and CAPHRI School for Public Health and Primary Care, Maastricht University, the Netherlands; **Nawaraj Upadhaya**, MA, MSc, Researcher, Transcultural Psychosocial Organization, Nepal; **Crick Lund**, PhD , Professor of Public Mental Health and Professor of Global Mental Health and Development, Alan J Flisher Centre for Public Mental Health, Department of Psychiatry and Mental Health, University of Cape Town, South Africa, and Centre for Global Mental Health, Health Service and Population Research Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK; **Graham Thornicroft**, PhD, Professor of Community Psychiatry, Centre for Global Mental Health and Centre for Implementation Science, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK; **Oye Gureje**, PhD, MA, Professor of Psychiatry and Director, WHO Collaborating Centre for Research and Training in Mental Health, Neurosciences and Substance Abuse, Department of Psychiatry, University of Ibadan, Nigeria; and Professor Extraordinary, Department of Psychiatry, Stellenbosch University, South Africa

Correspondence: Mark Jordans, King's College London, Institute of Psychiatry, Psychology and Neuroscience, 16 De Crespigny Park, Camberwell, London SE5 8AF, UK. Email: mark.jordans@kcl.ac.uk

First received 28 Aug 2018, final revision 26 Feb 2019, accepted 8 Mar 2019

Supplementary material

Supplementary material is available online at <https://doi.org/10.1192/bjpo.2019.22>.

Funding

This work was supported by the European Union's Seventh Framework Programme (FP7/2007–2013) under grant agreement no. 305968, and by the PRIME Research Programme Consortium, funded by the UK Department of International Development for the benefit of developing countries. G.T. is supported by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care (CLAHRC) South London and by the NIHR Applied Research Centre (ARC) at King's College London NHS Foundation Trust, and the NIHR Applied Research and the NIHR Asset Global Health Unit award. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. G.T. receives support from the National Institute of Mental Health of the National Institutes of Health under award number R01MH100470 (Cobalt study). G.T. is supported by the UK Medical Research Council in relation the Emilia (MR/S001255/1) and Indigo Partnership (MR/R023697/1). awards. I.P., C.L. and G.T. are funded by the NIHR Global Health Research Unit on Health System Strengthening in Sub-Saharan Africa, King's College London (GHRU 16/136/54) using UK aid from the UK Government. The views expressed in this publication are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care. No funding bodies had any role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The views expressed in this publication are not necessarily those of the funders.

Acknowledgements

We thank Drs Charlotte Hanlon and Atalay Alem and the TPO Nepal Emerald team for their work on this study. We also thank Grace Ryan (London School of Hygiene and Tropical Medicine) for her contributions to the development of the indicators and study design. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care. D.C. is a staff member of the World Health Organization. The authors alone are responsible for the views expressed in this publication and they do not necessarily represent the decisions, policy or views of the World Health Organization.

References

- 1 Thornicroft G, Chatterji S, Evans-Lacko S, Gruber M, Sampson N, Aguilar-Gaxiola S, et al. Undertreatment of people with major depressive disorder in 21 countries. *Br J Psychiatry* 2017; **210**(2): 119–24.
- 2 World Health Organization. *Mental Health Gap Action Programme: Scaling up Care for Mental, Neurological and Substance use Disorders*. WHO, 2008.
- 3 World Health Organization. *mhGAP Intervention Guide for Mental, Neurological and Substance use Disorders in Non-Specialized Health Settings (Version 2.0)*. WHO, 2016.
- 4 Aqil A, Lippeveld T, Hozumi D. PRISM framework: a paradigm shift for designing, strengthening and evaluating routine health information systems. *Health Policy Plann* 2009; **24**(3): 217–28.
- 5 Upadhaya N, Jordans MJD, Lund C, Alem A, Hanlon C, Kigozi F, et al. Mental health information systems in low and middle-income countries: cross country situation analysis. *Int J Ment Health Syst* 2016; **10**(60): 1–11.
- 6 Semrau M, Evans-Lacko S, Alem A, Ayuso-Mateos JL, Chisholm D, Gureje O, et al. Strengthening mental health systems in low and middle-income countries: the EMERALD programme. *BMC Med* 2015; **13**(79): 1–9.
- 7 Jordans MJD, Chisholm D, Semrau M, Upadhaya N, Abdulmalik J, Ahuja S, et al. Indicators for routine monitoring of effective mental healthcare coverage in low-and middle-income settings: a Delphi study. *Health Policy Plann* 2016; **31**(8): 1100–6.
- 8 Lund C, Tomlinson M, De Silva M, Fekadu A, Shidhaye R, Jordans MJ, et al. PRIME: a programme to reduce the treatment gap for mental disorders in five low- and middle-income countries. *PLoS Med* 2012; **9**(12): e1001359.
- 9 Eaton J, Gureje O, Silva M, Sheikh TL, Ekpe E, Abdulaziz M, et al. A structured approach to integrating mental health services into primary care: development of the Mental Health Scale Up Nigeria intervention (mhSUN). *Int J Ment Health Syst* 2018; **12**(1): 11.
- 10 Ryan G, De Silva M, Terver JS, Ochi OP, Eaton J. Information systems for global mental health. *Lancet Psychiatry* 2015; **2**(5): 372–3.
- 11 Eaton J, McCay L, Semrau M, Chatterjee S, Baingana F, Araya R, et al. Scale up of services for mental health in low-income and middle-income countries. *Lancet* 2011; **378**(9802): 1592–603.
- 12 World Health Organization. *Mental Health Information System*. WHO, 2005.
- 13 World Health Organization. *Mental Health Action Plan 2013-2020*. WHO, 2013.
- 14 Lauriks S, Buster MCA, de Wit MAS, Arah OA, Klazinga NS. Performance indicators for public mental healthcare: a systematic international inventory. *BMC Public Health* 2012; **12**(1): 214–40.
- 15 Ahuja S. Evaluation of mental health information systems in low- and middle-income countries: - experiences from a cross-country qualitative study. *Br J Psychiatry* (in press).
- 16 De Silva M, Lee L, Fuhr D, Rathod S, Chisholm D, Schellenberg J, et al. Estimating the coverage of mental health programmes: a systematic review. *Int J Epidemiol* 2014; **43**: 341–53.
- 17 Ahuja S, Shidhaye R, Semrau M, Thornicroft G, Jordans MJD. Mental health information systems in resource-challenged countries: experiences from India. *BJPsych Int* 2018; **15**(2): 43–6.

