Selective photoinduced reduction of nitroarenes to n-arylhydroxylamines

Article (Accepted Version)

Kallitsakis, Michael G, Ioannou, Dimitris I, Terzidis, Michael A, Kostakis, George E and Lykakis, Ioannis N (2020) Selective photoinduced reduction of nitroarenes to n-arylhydroxylamines. *Organic Letters*, 22 (11). pp. 4339-4343. ISSN 1523-7060

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/91350/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Selective Photoinduced Reduction of Nitroarenes to *N*-Arylhydroxylamines

Michael G. Kallitsakis,^{*,†} Dimitris I. Ioannou,[†] Michael A. Terzidis,[‡] George E. Kostakis,[£] and Ioannis N. Lykakis,^{*,†}

[†] Department of Chemistry, Aristotle University of Thessaloniki, University Campus, Thessaloniki 54124, Greece

[‡] Department of Nutritional Sciences & Dietetics, International Hellenic University, P.O. 141 Sindos, 57400, Thessaloniki, Greece

[£] Department of Chemistry, School of Life Sciences, University of Sussex, Brighton BN1 9QJ, UK

ABSTRACT: We report the selective photoinduced reduction of nitroarenes to *N*-arylhydroxylamines. The present methodology facilitates this transformation in the absence of catalyst or additives and uses only light and methylhydrazine. This non-catalytic photoinduced transformation proceeds with a broad scope, excellent functional-group tolerance and high yields. The potential of this protocol reflects on the selective and straightforward conversion of two general antibiotic molecules, azomycin and chloramphenicol, to the bio-active hydroxylamine species.

N-arylhydroxylamines (N-AHA) are versatile organic molecules with useful applications; valuable intermediates in the synthesis of biologically active molecules, inhibitors of polymerization and precursors for the synthesis of other organic fragments.¹ The selective reduction of nitroarenes is the most convenient protocol to obtain these attractive organic scaffolds; this transformation is promoted either by zinc dust in aqueous NH_4Cl , aluminum-amalgam or even enzymatic biocatalysts.¹ However, these reagents are used in stoichiometric amounts, a significant drawback; thus, alternative methodologies have been developed. These protocols involve heterogeneous catalytic systems (with Pt, Ru, Ni, Pd, Rh, Ir) and either molecular hydrogen^{1,2} or hydrogen donor molecules including borohydrides^{1,3a,b} or hydrazine hydrate^{1,3c-h} (Scheme 1). The former protocol requires high pressures and temperatures, while the latter promotes this selective transformation under milder reaction conditions, yet the use of additives is necessary.

Light is the most abundant renewable energy source, and therefore, photochemistry has emerged as an efficient, green, eco-friendly approach to promote chemical transformations, especially organic reductive processes.⁴ In this direction, the photoinduced reduction of nitroarenes has been achieved via photocatalytic hydrogenation or transfer hydrogenation processes.⁵ To date, metal-based protocols, including transition metal nanoparticles or complexes supported on active surfaces, promote

the transformation of nitroarenes to the corresponding arylamines^{6,7} or the corresponding azoxy and azo-compounds (Scheme 1A).⁸ Therefore, the selective photochemical synthesis of N-AHA from nitroarenes remains a significant challenge.

Given the importance of N-AHA in organic chemistry and inspired from metal-based^{9,10} or photochemical¹¹ methodologies developed in our laboratory that facilitate the synthesis of interesting organic scaffolds,¹⁰ we reasoned that it should be possible to develop a direct photoinduced reduction of nitroarenes to N-AHA with the use of methylhydrazine as hydrogen donor molecule (Scheme 1B). To the best of our knowledge, only specific examples with limited applicability on the photochemical reduction of 4-nitropyridine-1-oxide^{12a} and *para*-X-substituted rich nitroarenes (X=H, Me, MeO)^{12b-d} into the corresponding hydroxylamines in acidic alcoholic solvents have been reported.

In order to evaluate the photochemical reduction conditions, we used 4-nitrotoluene **1** as a probe molecule with different catalysts, hydrazine or methyl hydrazine and MeOH or MeCN as solvents (Table S1). Inspired by metal-based selective reduction protocols of nitroarenes developed in our laboratory,⁹ we initially studied the photo-transformation of **1** in the presence of Cu(II) salts or complexes.^{10a} In these cases a mixture of the corresponding *N*-(*p*-tolyl)hydroxylamine **1a**, toluidine **1b** and azoxy arene **1c** was obtained (Table S1, entries 1 and 2); while under thermal conditions, we observed no reaction takes place

(Table S1, entry 3). Similar results were also observed in the reactions with Co(II) salts and complexes, as catalysts, in MeOH or MeCN (Table S1, entries 4-6) and 9,10-dicyanoanthracene and Rose Bengal, as conventional photosensitizers, (Table S1, entries 7 and 8). The reaction in the absence of any catalyst but in the presence of hydrazine hydrate (Table S1, entries 9 and 10) yields a mixture of products, and amine **1b** is the major.

Scheme 1. Photo- and thermal-catalytic reductive processes of nitroarenes

Surprisingly, when we incorporated methylhydrazine in the absence of any catalyst, **1a** was selectively formed, in quantitative yield and the product could be isolated by a simple filtration over a short pad of silica, thus omitting chromatographic purification (Table 1, entry 1). To get a better insight and rationalize this unexpected result, we performed a set of reactions. We found that phenylhydrazine also promotes the selective formation of hydroxylamine **1a**, but at a lower isolated yield (Table 1, entry 2); 1,1-dimethylhydrazine performs poorly (Table 1, entry 3), whereas the use of 1,2-diphenylhydrazine, 4-nitrophenylhydrazine, *p*-toluenesulfonylhydrazine or hydrazine hydrate suppress the formation of **1a** (Table 1, entries 4-7). The addition of water (20 μ L, 0.2% V/V) with methylhydrazine, found not affect the transformation of **1** to **1a** (Table 1, entry 8), whereas under H₂ atmosphere (balloon) no photochemical conversion of **1** is accomplished (Table 1, entry 9). Control experiments revealed that the reaction does not proceed in the absence of methylhydrazine or light (Scheme S1), with the use of a cut-off filter at 420 nm or upon white LEDs irradiation (Table 1, entries 10-13). Moreover, less amount of methylhydrazine (3 or 4 mol-excess) or time (15 or 30 min) leads to the formation of **1a** in moderate yield (Table 1, entries 14-17). On the contrary, the use of methyl hydrazine, in 10 mol-excess, or prolonged reactions afforded **1a** in quantitative yields (Table 1, entries 18

and 19). The last set of reactions included the use of other various hydrogen donor molecules, i.e. hydrosilanes, boranes, ammonium formate or formic acid and isopropyl alcohol, which did not facilitate the selective reduction (Table S2).

Table 1. Hydrazine evaluation in the photoinduced reduction of **1.**

A/A	Hydrazines ^a	Relative yields ^b			
		1 (%)	1a (%)	1b (%)	1c (%)
1	MeNHNH ₂	-	>99	-	-
2	PhNHNH ₂	32	68	-	-
3	Me ₂ NNH ₂	5	32	-	63
4	PhNHNHPh	100	-	-	-
5	4-NO ₂ PhNHNH ₂	100	-	-	-
6	TsNHNH ₂	100	-	-	-
7	NH ₂ NH ₂ •H ₂ O	-	23	48	29
8 ^c	MeNHNH ₂	-	95	5	-
9	H ₂ bubbling	100	-	-	-
10	-	100	-	-	-
11 ^d	MeNHNH ₂	100	-	-	-
12 ^e	MeNHNH ₂	100	-	-	-
13 ^f	MeNHNH ₂	100	-	-	-
14	MeNHNH ₂ (3 mol-excess)	26	65	-	9
15	MeNHNH ₂ (4 mol-excess)	15	85	-	-
16	MeNHNH ₂ (10 mol-excess)	-	95	5	-
17	MeNHNH ₂ (15 min)	45	55	-	-
18	MeNHNH ₂ (30min)	16	84	-	-
19	MeNHNH ₂ (300min)	-	100	-	-

^aConditions: **1** (0.2 mmol), hydrazine source (1mmol), MeCN (1 mL), 60min. ^bRelative yields of products measured by the appropriate proton signals from the ¹H NMR of the crude reaction mixture. ^c20 μ L(0.2% V/V) of H₂O was added in the reaction mixture. ^dIn the absence of light at 28 °C and 70 °C for 1h, respectively. ^eIrradiation with a cut-off filter at 420 nm, t=1h. ^fIrradiation using white LED, 11W, for 24h.

Interestingly, the reaction, except acetonitrile, takes place in a variety of solvents including ethanol, isopropylalcohol, dimethylcarbonate, tetrahydrofuran, 1,2-dichloroethane and toluene (Table S3, entries 1-8). The reaction in ethyl acetate showed lower selectivity (Table S3, entry 9) with significant amounts of toluidine **1b** and azoxy-arene **1c**. No reaction occurs in water (Table S3, entry 10), but the use of a solvent mixture H₂O/CH₃CN (1:1) promotes the reduction and together with the formation of **1b** (Table S3, entry 11).¹³

Scheme 2. Photoinduced synthesis of various *N*-AHA using methylhydrazine.

After establishing the optimum conditions, we assessed the scope of the protocol towards various nitroarenes and focused on different variations. Remarkably, this photoinduced protocol shows a broad functional group tolerance and a series of nitroarenes (**1-22**) yielded the desired corresponding N-AHA (**1a-22a**) in excellent isolated yields (Scheme 2). The chemoselective reduction of the nitro group, in the presence of other reducible functional groups, is a very challenging task; however, this protocol leads to the selective and sole reduction of the nitro group to the corresponding hydroxylamine in the presence of carboxyl, cyano and carbonyl groups (**10a-14a**). Interestingly, under the present protocol, no reductive dehalogenated process occurs;¹⁴ and the corresponding halogenated aromatic N-AHA **15a-18a** species are formed in excellent yields. To further identify the limitations of the present light-driven procedure, heterocyclic substrates such as 6-nitrophthalide (**21**) and 6-nitro-1*H*-indazole (**22**) was reduced, giving quantitatively the corresponding hydroxylamines **21a** and **22a**, respectively. On the other hand, no reduction was observed with 1-nitro-hexane

even after 120 min of irradiation, a result that supports the necessity of an aromatic ring to facilitate this transformation. A lab-scale reaction was also performed using **1** (1 mmol) and MeNHNH₂ (5 mmol) in 3 mL of CH₃CN, and the corresponding N-AHA **1a** was isolated after 2 h irradiation time in 87% yield.

Given the simplicity of the present protocol, when compared with already known methodologies,^{2,3} we reasoned that the approach would be applicable for the synthesis of a series of biologically interesting molecules. Thus, aiming to expand the scope of this protocol in pharmaceutical relevant molecules, we attempted a direct photoinduced reduction of two antibiotic molecules; azomycin (**23**) and chloramphenicol (**24**).¹⁵ The hydroxylamine and/or nitroso derivatives of **23** were found to exhibit high potential for the treatment of latent tuberculosis (i.e. imidazole derivatives are also designed to intercalate with intracellular nucleophiles including DNA), and treatment of several bacterial infections. At the same time, the corresponding derivatives of **24** were used as an eye ointment to treat conjunctivitis.^{15a} To the best of our knowledge, the already known protocols that yield **23a** and **24a** involve metal-based, electrochemical and biosynthetic processes;¹⁶ with significant drawbacks such as low yields, difficulty in handling and separation of the mixture of several unstable products. Notably, the present simple methodology provides easy, clean and straightforward access to the corresponding N-AHA **23a** and **24a** in high isolated yields 96% and 90%, respectively, within 90 min reaction time (Scheme 3).

Scheme 3. Application of the present methodology to the selective photo-reduction of the antibiotics azomycin **23 and chloramphenicol **24**.**

Regarding the mechanism of this photochemical transformation, it is essential to note that:

The addition of an equimolar amount of 9,10-dicyanoanthracene (9,10-DCA, 0.2 mmol) retards the reaction process (**1a** in 45% and **1c** in 22%); however, the presence of 0.2 mmol of 1,3,5-trimethoxybenzene (TMB) does not affect the reaction conversion (**1a** in 90%) significantly (Scheme S3). In the same context, the addition of the radical inhibitor TEMPO in an equimolar amount did not suppress the reaction, which suggests that a free radical pathway is not predominate in this photochemical transformation (Scheme S3). These results support possible a proton-coupled electron transfer (PCET) process between the excited form of the nitroarene and the methylhydrazine; however, a hydrogen atom transfer (HAT) cannot be excluded.¹⁷

No formation of the N-AHA **5a** was observed during the photochemical reaction of the corresponding azoxy- (**5c**), azo- (**5e**) and hydrazobenzene (**5d**) under the same conditions and in the absence or in the presence of 0.2 mmol of H₂O as additive

(Scheme S4). This result supports the presence of a direct route of the reduction process.

Besides, the photoinduced reduction progress of the reaction between **5** and methylhydrazine was directly monitored by UV-Vis (Figure S1); the formation of the corresponding *N*-AHA **5a** was observed as the only product.

Of note is that the reaction of nitrosobenzene **5f** with five, three, two or even one equivalent of methylhydrazine at 28°C in CD₃CN yields, within 15 min yields to **5a** in >85% (Table S9), proved by direct ¹H NMR monitoring. This result supports the, possible, *in situ* formation of a nitrosoarene intermediate, which is transformed into the corresponding hydroxylamine in the presence of methylhydrazine through a non-photochemical pathway; although an electrochemical reductive pathway¹⁸ can also take part during the photo-reaction.

It is well-known that hydrazine hydrate produces harmless side-products such as nitrogen and water under photo-irradiation.¹⁹ Thus, for examining the possibility of decomposition, we performed a set of photo-reductions using methylhydrazine and phenylhydrazine, accordingly. In the former, a significant amount of methane was measured using a gas chromatography-thermal conductivity detector (GC-TCD) (Scheme S5), however, in the presence of phenylhydrazine, a significant amount of benzene was verified by ¹H-NMR spectroscopy, during the photochemical reduction of **1** into **1a** (Scheme S6). Moreover, the corresponding hydrazone of formaldehyde, a decomposition product of methylhydrazine was identified in the reaction mixture by ¹H-NMR spectroscopy (Figure S2).^{19a} These results indicate the presence of a radical degradation process of the methylhydrazine in the presence or absence of the nitroarene,^{17,20} and support further the necessity of methylhydrazine mol-excess based on nitroarene amount.

Based on these experimental data, we envisage a plausible mechanistic pathway shown in Scheme 4. The first step involves the excitation of nitrobenzene (PhNO₂^{*}), followed by a PCET or HAT pathway, between PhNO₂^{*} and methylhydrazine, to give PhN(OH) and PhN(OH)[•] radicals in an equilibrium form. Preliminarily kinetic studies performed in CD₃CN and monitored directly by ¹H NMR, showed a faster reduction for the electron-withdrawing substituted nitroarene (4-MeOOC, **10**) against the electron-donating substituted nitroarenes (4-Me, **1** and 4-MeO, **6**) (Figures S3-S5). This result may support the formation of the above-proposed radicals. Then, the PhN(OH)₂, through a second PCET or HAT process with methylhydrazine, transforms to *N*-phenyldihydroxylamine (PhN(OH)₂) and subsequently to the nitrosobenzene intermediate, after the loss of a water molecule, which can be directly reduced to the desired *N*-phenylhydroxylamine (PhNHOH) through a non-photochemical process (Scheme 4).^{19,20} Although this is a plausible mechanistic pathway, theoretical calculations and electrochemical studies are in progress to support this hypothesis.

In conclusion, we present an efficient methodology that selectively reduces nitroarenes to the corresponding *N*-AHA. The cooperation of nitroarenes and methylhydrazine orchestrate, a light-driven, *in situ* generation of nitrosoarenes which yield *N*-AHA in a fast and clean manner. The protocol has a broad scope, excellent functional-group tolerance, absence of by-products, while other reducible functional groups remain intact. The photoinduced reduction methodology is simple and applies to the bioactive molecules azomycin and chloramphenicol, giv-

ing rise to a plausible approach for a series of pro-drug molecules, thus opening new research avenues with high synthetic and biological impact.

Scheme 4. A plausible mechanism for the photoinduced synthesis of *N*-phenylhydroxylamine from nitrobenzene

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

General procedures, mechanistic studies, and NMR spectra

AUTHOR INFORMATION

Corresponding Authors

Michael G. Kallitsakis - Department of Chemistry, Aristotle University of Thessaloniki, University Campus, Thessaloniki 54124, Greece. kallitsos29@gmail.com

Ioannis N. Lykakis - Department of Chemistry, Aristotle University of Thessaloniki, University Campus, Thessaloniki 54124, Greece. Email: lykakis@chem.auth.gr

Authors

Dimitris I. Ioannou - Department of Chemistry, Aristotle University of Thessaloniki, University Campus, Thessaloniki 54124, Greece.

Michael A. Terzidis - Department of Nutritional Sciences & Dietetics, International Hellenic University, P.O. 141 Sindos, 57400, Thessaloniki, Greece.

George E. Kostakis - Department of Chemistry, School of Life Sciences, University of Sussex, Brighton BN1 9QJ, UK.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

MGK acknowledges the State Scholarships Foundation (IKY) for the financial support through the Operational Program «Human Resources Development, Education and Lifelong Learning» in the context of the project «Reinforcement of Postdoctoral Researchers

- 2nd Cycle” (MIS-5033021). MAT acknowledges financial support from the Hellenic Foundation for Research and Innovation (HFRI) and the General Secretariat for Research and Technology (GSRT) under the grant agreement No[776] “PhotoDaLu” (KA97507). We thank Prof. K. Triantafyllidis (Department of Chemistry, AUTH) for performing the GC-TCD experiments. We thank Dr. C. Gabriel (HERACLES Research Center, KEDEK, Laboratory of Environmental Engineering (EnvE-Lab), Department of Chemical Engineering, AUTH, Greece) for performing and analyzing the HRMS experiments. INL acknowledges the Empirikeion Foundation for financial support of the photo-apparatus.

REFERENCES

- (1) *The chemistry of Hydroxylamines, Oximes and Hydroxamic Acids* (Eds. Rappoport, Z. and Liebman, J. F. **2009**, John Wiley & Sons, Ltd.
- (2) (a) Boymans, E. H.; Witte, P. T.; Vogt, D. *Catal. Sci. Technol.* **2015**, *5*, 176-183. (b) Rong, Z.; Du, W.; Wang, Y.; Lu, L. *Chem. Commun.* **2010**, *46*, 1559-1561. (c) Takenaka, Y.; Kiyosu, T.; Choi, J.-C.; Sakakura, T.; Yasuda, H. *Green Chem.* **2009**, *11*, 1385-1390.
- (3) (a) Doherty, S.; Knight, J. G.; Backhouse, T.; Summers, R. J.; Abood, E.; Simpson, W.; Paget, W.; Bourne, R. A.; Chamberlain, T. W.; Stones, R.; Lovelock, K. R. J.; Seymour, J. M.; Isaacs, M. A.; Hardacre, C.; Daly, H.; Rees, N. H. *ACS Catal.* **2019**, *9*, 4777-4791. (b) Prakash, P.; De Masi, D.; Geertsen, V.; Misserque, F.; Li, H.-Namboothiri, I. N. N.; Gravel, E.; Doris, E. *Chemistry Select* **2017**, *2*, 5891-5894. (c) Jawale, D. V.; Gravel, E.; Boudet, C.; Shah, N.; Geertsen, V.; Li, H.; Namboothiri, I. N. N.; Doris, E. *Chem. Commun.* **2015**, *51*, 1739-1742. (d) Hojczyk, K. N.; Feng, P.; Zhan, C.; Ngai, M.-Y. *Angew. Chem. Int. Ed.* **2014**, *53*, 14559-14563. (e) Shil, A. K.; Das, P. *Green Chem.* **2013**, *15*, 3421-3428.
- (4) (a) Kou, J.; Lu, C.; Wang, J.; Chen, Y.; Xu, Z.; Varma, R. S. *Chem. Rev.* **2017**, *117*, 1445-1514. (b) Capaldo, L.; Ravelli, D. *Eur. J. Org. Chem.* **2017**, 2056-2071; (c) Friedmann, D.; Hakkı, A.; Kim, H.; Choi, W.; Bahnemann, D. *Green Chem.* **2016**, *18*, 5391-5411.
- (5) (a) Nabid, M. R.; Nazaru, N.; Asadi, S.; Heravi, M. M.; Sedghi, R. *Cur. Org. Chem.* **2016**, *20*, 696-734. (b) Kadama, H. K.; Tilvea, S. G. *RSC Adv.* **2015**, *5*, 83391-83407.
- (6) Recent photocatalytic hydrogenation processes: (a) Hao, C.-H.; Guo, X.-N.; Sankar, M.; Yang, H.; Ma, B.; Zhang, Y.; Tong, X.; Jin, G.; Guo, X.-Y. *ACS Appl. Mater. Interfaces*, **2018**, *10*, 23029-23036. (b) Yu, Z.; Chen, Z.; Chen, Y.; Peng, Q.; Lin, R.; Wang, Y.; Shen, R.; Cao, X.; Zhuang, Z.; Li, Y. *Nanoresearch* **2018**, *11*, 3730-3738.
- (7) Recent photocatalytic transfer hydrogenation processes: (a) Amanchi, S. R.; Kumar, K. V. A.; Lakshminarayana, B.; Satyanarayana, G.; Subrahmanyam, C. *New J. Chem.* **2019**, *43*, 748-754. (b) Todorov, A. R.; Aikonen, S.; Muuronen, M.; Helaja, J. *Org. Lett.* **2019**, *21*, 3764-3768. (c) Piggott, E. K.; Hope, T. O.; Crabbe, B. W.; Jalbert, P.-M.; Orlova, G.; Hallett-Tapley, G. L. *Catal. Sci. Technol.* **2017**, *7*, 5758-5765. (d) Xiao, Q.; Sarina, S.; Waclawik, E. R.; Jia, J.; Chang, J.; Riches, J. D.; Wu, H.; Zheng, Z.; Zhu, H. *ACS Catal.* **2016**, *6*, 1744-1753. (e) Tsutsumi, K.; Uchikawa, F.; Sakai, K.; Tabata, K. *ACS Catal.* **2016**, *6*, 4394-4398. (f) Yang, X.-Y.; Chen, B.; Zheng, L.-Q.; Wu, L.-Z.; Tung, C.-H. *Green Chem.* **2014**, *16*, 1082-1086.
- (8) (a) Chen, G.-J.; Xin, W.-L.; Wang, J.-S.; Cheng, J.-Y.; Dong, Y.-B. *Chem. Commun.* **2019**, *55*, 3586-3589. (b) Mondal, B.; Mukherjee, P. S. *J. Am. Chem. Soc.* **2018**, *140*, 12592-12601. (c) Chen, X. Y.; Fu, W.-F. *ACS Omega* **2018**, *3*, 1904-1911. (d) Chaiseeda, K.; Nishimura, S.; Ebitani, K. *ACS Omega* **2017**, *2*, 7066-7070 (e) Guo, X.; Hao, C.; Jin, G.; Zhu, H.-Y.; Guo, X.-Y. *Angew. Chem. Int. Ed.* **2014**, *53*, 1973-1977. (f) Zhu, H.; Ke, X.; Yang, X.; Sarina, S.; Liu, H. *Angew. Chem. Int. Ed.* **2010**, *49*, 9657-9661.
- (9) (a) Iordanidou, D.; Zarganes-Tzitzikas, T.; Neochoritis, C. G.; Dömling, A.; Lykakis, I. N. *ACS Omega*, **2018**, *3*, 16005-16013. (b) Papadas, I. T.; Fountoulaki, S.; Lykakis, I. N.; Armatas, G. S. *Chem. Eur. J.*, **2016**, *22*, 4600-4607. (c) Andreou, D.; Iordanidou, D.; Tamiolakis, I.; Armatas, G. S.; Lykakis, I. N.; *Nanomaterials*, **2016**, *6*, 54-65. (d) Fountoulaki, S.; Daikopoulou, V.; Gkizis, P. L.; Tamiolakis, I.; Armatas, G. S.; Lykakis, I. N. *ACS Catal.* **2014**, *4*, 3504-3511. (e) Gkizis, P. L.; Stratakis, M.; Lykakis, I. N. *Catal. Commun.* **2013**, *36*, 48-51.
- (10) (a) Kallitsakis, M.; Loukopoulos, E.; Abdul-Sada, A.; Tizzard, G. J.; Coles, S. J.; Kostakis, G. E.; Lykakis, I. N. *Adv. Synth. Catal.* **2017**, *359*, 138-145. (b) Andreou, D.; Kallitsakis, M. G.; Loukopoulos, E.; Gabriel, C.; Kostakis, G. E.; Lykakis, I. N. *J. Org. Chem.* **2018**, *83*, 2104-2113.
- (11) (a) Tzirakis, M. D.; Lykakis, I. N.; Orfanopoulos, M. *Chem. Soc. Rev.* **2009**, *38*, 2609-2621. (b) Symeonidis, T. S.; Athanasoulis, A.; Ishii, R.; Uozumi, Y.; Yamada, Y. M. A.; Lykakis, I. N. *ChemPhotoChem* **2017**, *1*, 479-484.
- (12) (a) Kanedo, C.; Yamada, S.; Yokoe, I. *Tet. Lett.* **1966**, 4729-4733. (b) Hashimoto, S.; Sunamoto, J.; Fujii, H.; Kano, K. *Bull. Chem. Soc. Jpn.* **1968**, *41*, 1249-1251. (c) Hashimoto, S.; Kano, K. *Tet. Lett.* **1970**, *11*, 3509-3512. (d) Hashimoto, S.; Kano, K. *Bull. Chem. Soc. Jpn.* **1972**, *45*, 549-553.
- (13) A further set of experiments with electron-withdrawing and electron-donating substituted nitroarenes were performed in MeOH and MeCN, and the results are summarized in Tables S4-S8.
- (14) (a) Marin, M.; Miranda, M. A.; Marin, M. L. *Catal. Sci. Technol.* **2017**, *7*, 4852-4858. (b) Sakamoto, H.; Imai, J.; Shiraishi, Y.; Tanaka, S.; Ichikawa, S.; Hirai, T. *ACS Catal.* **2017**, *7*, 8, 5194-5201.
- (15) (a) Kumar, P.; *Pharmacology and Therapeutics for Dentistry (Seventh Edition)*, **2017**, 457-487. (b) Qu, Y.; Spain, J. C. *Environ. Microbiol.* **2011**, *13*, 1010-1017. (c) Singh, R.; Manjunatha, U.; Boshoff, H. I.; Ha, Y. H.; Niyomrattanakit, P.; Ledwidge, R.; Dowd, C. S.; Lee, I. Y.; Kim, P.; Zhang, L.; Kang, S.; Keller, T. H.; Jiricek, J.; Barry, C. E. *Science* **2008**, *322*, 1392-1395.
- (16) (a) McClelland, R. A.; Panicucci, R.; Rautht, A. M. *J. Am. Chem. Soc.* **1987**, *109*, 4308. (b) Corbett, M. D.; Chipko, B. R. *Antimicrob. Agent Chemother.* **1978**, *13*, 193-198. (c) Lu, H.; Chanco, E.; Zhao, H. *Tetrahedron* **2012**, *68*, 7651-7654. (d) Jakubec, P.; Urbanova, V.; Medrikova, Z.; Zboril, R. *Chem. Eur. J.* **2016**, *22*, 14279-14284.
- (17) Selected reports on the photochemistry of nitroarenes: (a) Wubbers, G. G.; Snyder, E. J.; Coughlin, E. B. *J. Am. Chem. Soc.* **1988**, *110*, 2543-2548. (b) D. Döpp, D. *Topics Curr. Chem.* **1975**, *55*, 49-85. (c) Frolov, A. N.; Kuznetsova, N. A.; Eltsöv, A. V. *Russ. Chem. Rev.* **1976**, *45*, 1024-1034. (d) Cu, A.; Testa, A. C. *J. Am. Chem. Soc.* **1974**, *96*, 1963-1965. (e) Barltrop, J. A.; Bunce, N. J.; Thomson, A. J. *Chem. Soc.* **1968**, 1467-1474. (f) Hurley, R.; Testa, A. C. *J. Am. Chem. Soc.* **1966**, *88*, 4330-4332.
- (18) (a) Steudel, E.; Posdorfer, J.; Schindler, R. N. *Electrochim. Acta* **1995**, *40*, 1587-1594. (b) Huang, Y.; Lessard, J. *Electroanalysis* **2016**, *28*, 2716-2727.
- (19) Selected reports on the photochemical decomposition of hydrazine (a) Lambert, C. E.; Shank, R. C. *Carcinogenesis* **1988**, *9*, 65-70. (d) Hawkinst, W. G.; Houston, P. L. *J. Phys. Chem.* **1982**, *86*, 704-709. (g) Arvis, M.; Devillers, C.; Gillois, M.; Curtat, M. *J. Phys. Chem.* **1974**, *78*, 1356-1360. (h) Stief, L. J.; DeCarlo, V. J.; *J. Chem. Phys.* **1966**, *44*, 4638-4639.
- (20) King, D. M.; Bard, A. J. *J. Am. Chem. Soc.* **1965**, *87*, 419-423