Optiflow vs Vapotherm as extended weaning mode from nasal continuous positive airway pressure in preterm infants ≤ 28 weeks gestational age

Article (Accepted Version)

Fernandez-Alvarez, J Ramon, Mahoney, Liam, Gandhi, Rashmi and Rabe, Heike (2020) Optiflow vs Vapotherm as extended weaning mode from nasal continuous positive airway pressure in preterm infants ≤ 28 weeks gestational age. *Pediatric Pulmonology*. ISSN 8755-6863

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/91949/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Cover page

Comparison of MRI and Neurosonogram 1- and 2-Dimensional Morphological Measurements of the Newborn Corpus Callosum

Michael Mills¹ *, Vincent Pelling¹, Lisa M Harris¹, Joely Smith², Neil Aiton³, Heike Rabe^{3,4}, Jose Ramon Fernandez-Alvarez^{3,4}

¹Department of Radiological Sciences, Brighton & Sussex University Hospitals NHS Trust, Brighton, UK

² Radiological Sciences Unit, Imperial College Healthcare NHS Trust, London, UK

³Brighton and Sussex Medical School, Brighton, UK

⁴Department of Neonatology, Brighton & Sussex University Hospitals NHS Trust, Brighton, UK

* Address for correspondence:

Michael Mills

Office G8

St Mary's Hall,

Eastern Road,

Brighton, UK BN2 5JF

Tel: +44 01273 523473 Ext. 63471

Email: Michael.Mills@bsuh.nhs.uk

No financial assistance was received in support of the study

No financial ties or conflict exist for any author

Study category: Clinical Research Article

Abstract

Background:

Developmental abnormalities of the corpus callosum (CC) are linked to multiple neuro-developmental disorders, for which neonatal neuroimaging may allow earlier diagnosis and intervention. MRI is often considered the most sensitive imaging modality to white matter changes, while neurosonogram (NS) remains the clinical staple. This study assess the correlation between MRI and US measurements of the neonatal CC using a protocol derived from established methodologies

Methods: MR and NS images from an existing cohort of term infants (≥ 37 weeks gestational age) were studied. Length and area measurements of the CC made with linear (LUS) and phased array US (PUS) data were compared to those from MRI. Intra-observer reliabilities were estimated.

Results: Moderate to strong correlation strengths were observed for length measurements and the total area of the CC. Sectional area measurements showed poorer correlations. Bland-Altman plots support improved correspondence of length and total area measurements. LUS data appeared to correspond closer to MRI. All three modalities showed comparable repeatability.

Conclusion: NS correlates well with some MRI measurements of the CC and shows similar levels of repeatability, making them possibly interchangeable. Use of LUS, a technique rarely used for NS, may be preferable to the standard approach for morphological studies.

Introduction

The Corpus Callosum (CC) is a bundle of hundreds of millions of myelinated and non-myelinated nerve fibres connecting the two hemispheres of the brain ¹. It has been shown to be involved in functions such as sound localisation ² and stereoscopic vision ³, which has led to the hypothesis that its evolution occurred to facilitate integration of sensory signals between the brain's two hemispheres ^{4,5}.

Developmental abnormalities of the CC are linked to multiple neuro-developmental disorders such as: attention deficit hyperactivity disorder, autism, dyslexia and foetal alcohol spectrum disorder ⁶⁻⁸. Early detection of the malformation or underdevelopment of the CC, which are associated with these disorders, allows for earlier interventions on the part of health care professionals and local public sector providers. Although contradictory results can be found in the literature, outcomes are often improved as a result of intervention programmes for children with known neurological disorders or learning disabilities ⁹⁻¹¹.

Foetal and neonatal neuroimaging using brain MRI or neurosonogram (NS), including the CC, is in common clinical practice ¹². Cranial US is often considered the preferred imaging modality for studies of the neonatal brain as it can be performed at the bedside, is non-ionising, requires little disturbance of the neonate and is relatively inexpensive compared to other imaging modalities ¹³. MRI however is often considered to be the most sensitive imaging modality to changes in white matter structures and measurements of brain growth ¹⁴⁻¹⁶, and has been used in many studies which correlate measurements of the CC with neurological outcomes. MRI is also not limited to specific imaging orientation and locations, while NS may only be performed through specific viewing windows; the neonatal fontanelles. However,

MRI is time consuming, expensive, often requires patient sedation and is acoustically noisy.

Recent publications show no significant differences on comparison of linear measurements of some neuroanatomic structures when obtained using MR and US: width, height, length and depth of the corpus callosum, cerebellum, pons, cavum septum pellucidum, fourth ventricle, basal ganglia and caudate head ¹⁷. Measurements of the ventricular index and diameter of the posterior horn of the lateral ventricle, extra-cerebral space, inter-hemispheric fissure and cortex of the cingulate gyrus did show significant differences however ¹⁷. Strong correlations have also been reported between ventricular volumes for paediatric patients as measured on MR and US images¹⁸.

In this study we aimed to investigate the level of correspondence between measurements of the neonatal CC as measured from MRI and US datasets using a phased array ultrasound (PUS) and linear array ultrasound (LUS) transducer, and established methodologies for these techniques, in a group of neonates undergoing cooling therapy after a diagnosis of hypoxic ischemic encephalopathy (HIE). We also aimed to assess the intra-rater reliability of these measurements.

Materials and Methods

We performed a retrospective cohort study of 39 term neonates, diagnosed with HIE, admitted to a tertiary surgical neonatal department from January 2014 to November 2015, undergoing total body cooling therapy, and imaged as part of the normal clinical pathway.

All imaging was performed on a 1.5T Siemens Symphony MRI scanner (Siemens, Erlangen, Germany), and a Philips HD11 XE US scanner (Philips, Eindhoven, The Netherlands). US images were obtained with either a 4-12MHz PUS transducer or a 5-12MHz LUS transducer, using presets designed for neonatal cranial use. MR Images were acquired using the standard clinical protocol: the T1 weighted sagittal and T2 weighted turbo inversion recovery magnitude (TIRM) coronal images of the clinical protocol were used within this study (summary MRI details can be found in Table.1).

-Insert Table 1 Here-

Multiple datasets were available for US images and MRI. To minimise selection bias only images from datasets acquired within a seven day period between all three imaging sessions were selected for analysis. Datasets were then chosen based on the identification of anatomical landmarks and ensuring the image captured the entire extent of the CC in the mid-sagittal plane. For MR, the mid-sagittal slice was identified as that which clearly showed the patient's nose and for which the cerebellar vermis was visualised and the spinal cord well resolved. For US, the superior border of the CC had to be clearly identifiable against its surroundings and,

the cavum septum pellucidum, the cerebellar vermis and the aqueduct had to all be visible on one image. Coronal slices on which to assess the thickness of the corpus callosum were initially selected based on a clear visualisation of the interhemispheric fissure, cavum septum pellucidum and the foramina of Monro.

-Insert Figure 1 Here-

All measurements on NS and MRI were performed as described previously for newborn infants by a single reviewer (MM), a medical physicist with experience of viewing medical images, under the guidance of a Neonatologist with expertise in neuroimaging (RF) ^{17,19,20}. Four lengths and eight areas were determined on the mid-sagittal images, while a single length was measured on the coronal images as follows. The long axis diameter of the CC was defined as the greatest length measured by drawing a line from the anterior edge of the most anterior portion to the posterior edge of the most posterior portion of the corpus callosum. From the long axis line all sectional areas of the CC were then defined. Seven regions were defined by drawing perpendicular lines to the long axis midpoint, at one third and at one fifth from the anterior and posterior end. A final line was drawn perpendicular to the long axis line at the inner border of the anterior notch of the CC. These divisions defined the rostrum, genu, rostral body, anterior midbody, posterior midbody, isthmus and splenium as defined by Witelson (Figure 2) ²⁰. The body width of the CC was defined as the thickness of the CC at its midpoint, perpendicular to the long axis. The genu width and splenium width were then defined as the greatest diameter when drawing a perpendicular line from the most anterior edge to the most posterior edge of the boundaries of the genu or splenium sections (as defined above). The area of each section of the CC was summed to yield its total area. Figure 2 shows a schematic of the sagittal measurements and Figure 3 shows this process in four stages on an

example anonymised ultrasound image from one of the included subjects. A single length measurement in the coronal plane, the coronal width (described by Liu et.al. as the antero-posterior diameter), was also obtained by measuring the thickness of the CC above the cavum septum pellucidum and below the inter-hemispheric fissure, perpendicular to its edges ¹⁹.

Prior to all measurements, a histogram normalisation of the image window and level was performed. This was performed to improve the consistency of viewing conditions. Measurements were then obtained using the length calliper and freehand area measurement tools of the local PACS viewer (Philips IntelliSpace PACS 4.4 Enterprise). Image zooming by 400% was also performed to ensure the CC encompassed as much of the viewing monitor as possible. No additional image processing occurred.

Intra-observer variability was assessed by performing ten repeat measurements on three randomly selected neonate's images, for which LUS and PUS transducer and MRI data were present. The reviewer was blinded to previous results. Intra-observer error was estimated using the coefficient of variation (COV):

$$COV = \frac{\sigma}{\bar{x}} \times 100\%$$

where σ and $\bar{x}$ are the standard deviation and mean recorded values. The results across the three patients' images were then pooled.

All analysis was performed using SPSS23 (IBM, Armonk, NY). Descriptive statistics, median measurement and interquartile range (IQR), and correlation strength from comparisons of measurements from different modalities are shown in Table 1.

Statistical analysis of correlations was performed using the Spearman's correlation coefficient, assuming non-normal data distribution: P value < 0.001 was taken as statistically significant. Correlation by Spearman's coefficient was graded: strong ($p > 0.7$), moderate ($0.4 \geq p \leq 0.7$), weak ($p < 0.4$). Additionally Bland-Altman plots were constructed for each measurement performed comparing the MRI to the US technique. Sets of 95% confidence intervals were constructed for these plots as:

$$CI_{\pm} = \bar{d} \pm 1.96s$$

where $\bar{d}$ is the mean of the calculated differences between imaging methods (for example the difference in measurements of the long axis length as measured on MRI and LUS images), and s the standard deviation of the differences. IRB approval was not required.

-Insert Figure 2 Here-

-Insert Figure 3 Here-

Results

Thirty two neonates who underwent MRI and US provided useable data: seven were excluded due to missing data or the entire CC not residing within the image. All included studies were of term babies (53% male), with a median birth weight of 3415g (IQR 762g).

Statistically significant correlations (all moderate to strong) were observed for the long axis, genu and sectional areas 2 and 7 as well as for the total area, for both modality comparisons. The LUS to MRI comparison also yielded significant correlations (strong) for the splenium length and coronal width measurements. Moderate correlation strengths were seen for all the other measurements except for the area of regions 3, 4, 5 and 6 (rostral body, anterior midbody, posterior midbody and isthmus). The correlation strength generally increased with the measurement scale. Stronger correlations to MRI measurements were observed for LUS than PUS images.

The Bland-Altman plots' 95% confidence intervals were of the order of the size of the measurement values, or larger, for sectional area measurements of MRI compared to LUS and PUS (Supplemental Figure S1 and S2). Confidence intervals were found to be proportionately smaller for the length and total area measurements. Comparable performance was observed for both PUS and LUS compared with MRI datasets.

The lowest intra-observer error, estimated by the COV, for MRI, PUS and LUS was seen for measurements of the long axis length and the total cross-sectional area, while measurements of the area of the rostrum and genu show the largest COV, approaching 20%. A comparable level of performance could be seen in each of the

modalities (Table 3). However the LUS results show the lowest COV for the majority of measurements.

-Insert Table 2 Here-

-Insert Table 3 Here-

Discussion

Our study shows that measurements made from NS are comparable with measurements made from MR images for many structural measurements of the corpus callosum and may hence be reliably used to characterise the structure of the CC. Length and total area measurements showed higher correlation strengths, the majority of which were significant at the P value < 0.001 level, compared to the sectional areas of the Witelson segmentation: sectional area measurements showed a wide spread of correlation strengths, half of which were poor ($p < 0.4$). Bland-Altman analysis also showed that the tightest 95% confidence intervals, relative to the measurement magnitude, were seen for the length and total area measurements, suggesting they were more strongly related. The clinical significance of the limits of agreement is unknown.

The higher correlation strengths, smaller confidence intervals, and greater repeatability for the length measurements, especially the long axis length, and total cross-sectional area, make these the most suitable candidates for future study, especially considering their potential medical implications^{19,21–24}. This improved performance may in part be due to the small size of many of the regional areas which will have a larger proportional measurement uncertainty. Correlation strength does not simply improve consistently with increases in area however (see Table.2). All regional areas of the CC as defined here are defined in relation to the long axis line, which is manually defined²⁰. If the placement of this line varies over the three imaging modalities the measured areas will change. The long axis length measurement is perhaps the most robust measurement explored here; however even small changes in its angulation may have large implications to the regional areas which result.

The repeatability of any measurement is important to ensure the conclusions drawn from a set of results can be applied to multiple patients. The results of the intra-observer error testing are encouraging with all pooled COVs below 20%, with comparable levels of repeatability observed across all imaging techniques. The long axis length and total cross-sectional area show the greatest level of repeatability. Greater repeatability was seen compared to that reported by Davies et.al ²⁵ who assessed inter- and intra-observer variability of measurements of the width of the anterior horn, thalamo-occipital distance of the lateral ventricles and widths of the third and fourth ventricles assessed via NS in neonates. While these are neither the same cranial structures nor a matched cohort, they are of similar scales to those measured here. Future research should include a robust assessment of intra- and inter-observer error of measurements obtained from clinical members of staff more accustomed to reviewing neonatal neuroimages.

It had been hypothesized that the LUS data would yield closer correspondence to their MRI counterparts given the improved resolution possible with a less-divergent beam and, in this instance, marginally increased central probe frequency. This study appears to add weight to this hypothesis as stronger correlations between length measurements (except for genu width) and total area, the most reliable measurements investigated, were observed when the LUS data was compared to MRI. Further testing may establish if this apparent improvement is sustained when confounding variables, such as differences in probe frequency, are controlled for.

Throughout this study all segmentation and measurement of the CC has been performed manually. This is a lengthy and difficult process, which may introduce increased measurement variability and uncertainty. The use of automatic measurement and segmentation techniques, of which recent research demonstrates

the viability for some features of the fetal or neonatal brain ^{26–28}, should be explored in future work: the results of which may make measurement of CC features more feasible in the clinic.

This study was entirely retrospective and hence there was no control of imaging protocol. Future work would benefit from bespoke imaging protocols to ensure effective imaging of the CC specifically is achieved. This may include altering imaging parameters such as the use of a central probe frequency of 12MHz for both LUS and PUS imaging, the use of adaptive image processing to enhance boundary detection, and spatial compounding to reduce visual clutter which may obscure the edges of the CC, for US images. The neonatal brain has a much higher water content than the adult and the myelination of neurones is far from complete, both of which can affect image contrast ^{29,30}. In order to overcome the *T1* and *T2* lengthening associated with the increased water content, and hence to recover better contrast in the brain, increases in the *TE* and *TR* values used in the MRI studies may also result in improved measurement precision ^{29,30}. Increased field strength may be beneficial, however the corresponding safety implications must be considered.

It should be noted that the patient cohort, term neonates with a diagnosis of HIE, are not a cohort representative of the entire neonatal population. However, the increased risk of neurological impairment in this study group make them a reasonable initial study group ³¹.

Conclusion

The development of the corpus callosum is associated with multiple neurological impairments. The ability to identify patients as being at increased risk of specific neurological disorders would allow the care pathway for these patients to be optimised. The use of a bedside, cost-effective, non-invasive and non-ionising imaging modality would be ideal for early diagnostic/prognostic assessment.

This report is novel as data from both linear and phased ultrasound arrays is compared with MRI datasets. In this study we demonstrated that some measurements of the neonatal CC obtained US images, especially those with a linear array probe, correlate well with those from MRI. Long axis length and total area appear the best correlated and most precise measurements in this study. These results suggest that some measurements are possibly interchangeable between MRI and US, and hence the biological correlates currently found in studies utilising MRI may be expected from US studies. The feasibility and quality of large scale research may therefore be increased and improved by focussing on linear measurements of the CC long axis. The use of imaging protocols designed to optimize the definition and contrast of the CC and employing automatic segmentation and analysis should also be explored.

References

1. Waxman SG. Clinical Neuroanatomy. 26th ed. McGraw Hill; 2010.
2. Gazzaniga MS. Cerebral specialization and interhemispheric communication Does the corpus callosum enable the human condition ? Brain 2000;123(7):1293–326.
3. Aboitiz F, López J, Montiel J. Long distance communication in the human brain : timing constraints for inter-hemispheric synchrony and the origin of brain lateralization. Biol Res 2003;36(1):89–99.
4. Aboitiz F, Montiel J. One hundred million years of interhemispheric communication : the history of the corpus callosum. Brazilian J Med Biol Res 2003;36(4):409–20.
5. Mhrshahi R. The Corpus Callosum as an Evolutionary Innovation. J Exp Zool Part B Mol Dev Evol 2006;306(1):8–17.
6. Paul LK. Developmental malformation of the corpus callosum: A review of typical callosal development and examples of developmental disorders with callosal involvement. J Neurodev Disord 2011;3(1):3–27.
7. Xia Z, Hoeft F, Zhang L, Shu H. Neuroanatomical anomalies of dyslexia: Disambiguating the effects of disorder, performance, and maturation. Neuropsychologia 2016;81:68–78.
8. Just MA, Cherkassky VL, Keller T a., Kana RK, Minshew NJ. Functional and anatomical cortical underconnectivity in autism: Evidence from an fmri study of an executive function task and corpus callosum morphometry. Cereb Cortex 2007;17(4):951–61.
9. Arnold LE, Hodgkins P, Kahle J, Madhoo M, Kewley G. Long-Term Outcomes of ADHD : Academic Achievement and Performance. J Atten Disord 2015;19(1):[Epub ahead of print].
10. Mathes PG, Denton CA, Fletcher JM, Anthony JL, Francis DJ. The effects of theoretically different instruction and student characteristics on the skills of struggling readers. Read Res Q 2005;40(2):148–82.
11. Mcgoey KE, Eckert TL, Dupaul GJ, Hyperactivity AT. Early Intervention for Preschool-Age Children with ADHD : A Literature Review. J Emot Behav Disord 2002;10(1):14–28.
12. Wezel-meijler G Van, Steggerda SJ, Leijser LM. Cranial Ultrasonography in Neonates : Role and Limitations. Semin Perinatol 2010;34(1):28–38.
13. Meijler G. Neonatal Cranial Ultrasonography. 2nd ed. Springer; 2012.

14. Miller SP, Cozzio CC, Goldstein RB, et al. Comparing the Diagnosis of White Matter Injury in Premature Newborns with Serial MR Imaging and Transfontanel Ultrasonography Findings. *Am J Neuroradiol* 2003;24(8):1661–9.
15. Maalouf EF, Duggan PJ, Counsell SJ, et al. Comparison of Findings on Cranial Ultrasound and Magnetic Resonance Imaging in Preterm Infants. *Pediatrics* 2001;107(4):719–27.
16. Rademaker KJ, Uiterwaal CS, Beek FJ, et al. Neonatal cranial ultrasound versus MRI and neurodevelopmental outcome at school age in children born preterm. *Arch Dis Childhood-Fetal Neonatal Ed* 2005;90(6):F489–94.
17. Leijser LM, Srinivasan L, Rutherford M a., Counsell SJ, Allsop JM, Cowan FM. Structural linear measurements in the newborn brain: Accuracy of cranial ultrasound compared to MRI. *Pediatr Radiol* 2007;37(7):640–8.
18. Gilmore JH, Gerig G, Specter B, et al. Infant cerebral ventricle volume: a comparison of 3-D ultrasound and magnetic resonance imaging. *Ultrasound in Medicine and Biology* 2001;27(8):1143–6.
19. Liu F, Cao S, Liu J, Du Z, Guo Z, Ren C. Ultrasound measurement of the corpus callosum and neural development of premature infants *. *Neural Regen Res* 2013;8(26):2432.
20. Witelson S. Hand and sex differences in the isthmus and genu of the human corpus callosum: a postmortem morphological study. *Brain* 1989;112(3):799–835.
21. Thompson DK, Lee KJ, Bijnen L Van, et al. Accelerated Corpus Callosum Development in Prematurity Predicts Improved Outcome. *Hum Brain Mapp* 2015;36(10):3733–48.
22. Anderson NG, Laurent I, Cook N, Woodward L, Inder TE. Growth rate of corpus callosum in very premature infants. *Am J Neuroradiol* 2005;26(10):2685–90.
23. Ghassabian A, Herba CM, Roza SJ, et al. Infant brain structures, executive function, and attention deficit/hyperactivity problems at preschool age. A prospective study. *J Child Psychol Psychiatry* 2013;54(1):96–104.
24. Handel LLEVAN, Uiterwaal CSPM, Groenendaal F, et al. Corpus Callosum Size in Relation to Motor Performance in 9- to 10-Year-Old Children with Neonatal Encephalopathy. *Pediatr Res* 2008;63(1):103.
25. Davies MW, Swaminathan M, Chuang SL, Betheras FR. Reference ranges for the linear dimensions of the intracranial ventricles in preterm neonates. *Arch Dis Childhood-Fetal Neonatal Ed* 2000;82(3):F218–23.

26. Huang R, Namburete A, Noble A. Learning to segment key clinical anatomical structures in fetal neurosonography informed by a region-based descriptor. *J Med Imaging* 2018;5(1):014007.
27. Beare RJ, Chen J, Kelly CE, et al. Neonatal Brain Tissue Classification with Morphological Adaptation and Unified Segmentation. *Front Neuroinform* 2016;10:12.
28. Qiu W, Chen Y, Kishimoto J, et al. Automatic segmentation approach to extracting neonatal cerebral ventricles from 3D ultrasound images. *Med Image Anal* 2017;35:181–91.
29. Saunders DE, Thompson C, Gunny R, Jones R, Cox T. Magnetic resonance imaging protocols for paediatric neuroradiology. *Pediatr Radiol* 2007;37(8):789–97.
30. Rutherford MA. MRI of the Neonatal Brain [Internet]. 2002. Available from: <http://www.mrineonatalbrain.com>
31. Jacobs SE, Berg M, Hunt R, Tarnow-Mordi WO, Inder TE, Davis PG. Cochrne Review: Cooling for newborns with hypoxic ischaemic encephalopathy. *Evidence-Based Child Heal* 2010;5(1):474–531.

Figure Legends

Figure 1. Example of midsagittal images of a single patient from MRI (A), Linear Array US (B) and Phased Array US (C).

Figure 2. Schematic of the Corpus Callosum showing the divisions of the Witelson scheme. Clockwise these regions correspond to the rostrum, genu, rostral body, anterior midbody, posterior midbody, isthmus and splenium. The total area is the sum of these seven areas. The four sagittal length measurements are also displayed. Adapted from ²⁰.

Figure 3. Example Linear Array US image of the corpus callosum demonstrating the measurement procedure for measurements obtained in the midsagittal plane. The midsagittal image is first identified (A). The longest axial diameter is then defined as a line from the most anterior to the most posterior boundary of the corpus callosum. Its seven regions, outlined in Figure 2, are drawn perpendicular to this line (B). These perpendicular bisectors of the long axis were defined by drawing perpendicular lines to the long axis midpoint, at one third and at one fifth from the anterior and posterior end. A final line is drawn perpendicular to the long axis line at the inner border of the anterior notch of the CC. The seven regions of the corpus callosum are then drawn manually (C). The body width was defined as the thickness of the CC at its midpoint, perpendicular to the long axis. The genu width and splenium width were then defined as the greatest diameter when drawing a

perpendicular line from the most anterior edge to the most posterior edge of the boundaries of the genu or splenium sections (D).

Figure 1

Figure 2

Figure 3