An overview of the innate and adaptive immune system in atherosclerosis

Article (Accepted Version)

Mohammad-Rezaei, Mina, Arefnezhad, Reza, Ahmadi, Reza, Abdollahpour-Alitappeh, Meghdad, Mirzaei, Yousef, Arjmand, Mohammad-Hassan, Ferns, Gordan A, Bashash, Davood and Bagheri, Nader (2021) An overview of the innate and adaptive immune system in atherosclerosis. IUBMB Life, 73 (1). pp. 64-91. ISSN 1521-6543

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/96636/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

An Overview of the Innate and adaptive immune System in Atherosclerosis

Mina Mohammad-Rezaei¹, Reza Arefnezhad^{2,3}, Reza Ahmadi⁴, Meghdad Abdollahpour-Alitappeh⁵, Yousef Mirzaei⁶, Mohammad-Hassan Arjmand^{1,7}, Gordon A Ferns⁸, Davood Bashash⁹, Nader Bagheri^{a*}

¹ Cellular and Molecular Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

² Halal Research Center of IRI, FDA, Tehran, Iran.

³ Department of Anatomy, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran

⁴ Clinical Biochemistry Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

⁵ Cellular and Molecular Biology Research Center, Larestan University of Medical Sciences, Larestan, Iran.

⁶ Department of Biogeosciences, Scientific Research Center, Soran University, Soran, Kurdistan Region, Iraq.

⁷ Cancer Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

⁸ Brighton & Sussex Medical School, Division of Medical Education, Falmer, Brighton, Sussex BN1 9PH, UK

⁹ Department of Hematology and Blood Banking, School of Allied Medical Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

***Corresponding author:** Cellular and Molecular Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran. Tel: +98 9181731073; Fax: +98-3813330709. E-mail addresses: n.bagheri1985@gmail.com

Abbreviations

APCs: Antigen-presenting cells

ApoB100: Apolipoprotein B100

APPs: Acute-phase proteins

Arg1: Arginase 1

Bcl6: B cell lymphoma 6

CAD: Coronary artery disease

CRP: C-reactive protein

CVD: Cardiovascular disease

DAMPs: Damage-associated molecular patterns

DCs: Dendritic cells

FKN: Fractalkine

FOXP3: Forkhead box P3

GM-CSF: Granulocyte-macrophage colony-stimulating factor

HLA-DR: Human leukocyte antigen

HSPs: Heat-shock proteins

HSP60: Heat shock protein 60

HSP70: Heat shock protein 70

ICAM -1: Intercellular adhesion molecule 1

ICOS: Inducible T cell co-stimulator

ICOSL: Inducible T cell co-stimulator ligand

IDO: Indoleamine 2,3-dioxygenase

IFN: Interferon

IFN γ : Interferon-gamma,

IL: Interleukin

IL-17R: IL-17 receptor

iNKT: Invariant NKT

IRF4: Interferon regulatory factor 4

iTregs: Induced Tregs

JAK1: Janus kinase 1

JAK3: Janus kinase 3

LGL: Large granular lymphocytes

LPS: Lipopolysaccharide

LXR: Liver x receptor

MCP-1: Monocyte chemoattractant protein-1

MerTK: Mer receptor kinase

MHC: Major histocompatibility complex

MMP: Matrix metalloproteinase

MPO: Myeloperoxidases

MR: Mannose receptor

mDC: Myeloid DC

NET: Neutrophil extracellular traps

NK: Natural killer

NKT: Natural killer T

Nrf2: Nuclear erythroid-2 related factor

nTregs: Natural Tregs

ox-LDL: Oxidized low-density lipoprotein

PCR: Polymerase chain reaction

pDC: Plasmacytoid DC

PECAM -1: Platelets, endothelial cell adhesion molecule 1

PF4: Platelet factor 4

PPAR γ : Peroxisome proliferator-activated receptor γ

PTX3: Pentraxin-3

RNS: Reactive nitrogen specie

ROS: Reactive oxygen species

ROR γ t: Retinoic acid-related orphan receptor gamma t

RUNX1: Runt-related transcription factor 1

SMC: Smooth muscle cells

STAT3: Signal transducer and activator of transcription 3

STAT-4: Signal transducer and activator of transcriptoion-4

STAT6: signal transducer and activator of transcription 6,

TBX21: T-box transcription factor,

TFH: T follicular helper,

TGF- β : transforming growth factor beta,

Th: T-helper,

TLR4: toll-like receptor 4,

TNF- α : tumor necrosis factor-alpha,

TRAIL: tumor necrosis factor-related apoptosis-inducing ligand,

Tregs: regulatory T cells,

VCAM3: vascular cell adhesion molecule 3,

VEGF: vascular endothelial growth factor,

VSMCs: vascular smooth muscle cells.

Abstract

Cardiovascular disease is the leading cause of death globally. Coronary artery disease (CAD) is a chronic inflammatory disease usually caused by atherosclerosis, in which the coronary arteries become narrowed by atheromatous plaque. Plaques in atherosclerosis are formed through the accumulation of lipids and various immune cells. Both adaptive and innate immune systems are involved in the pathogenesis of atherosclerosis and facilitate plaque formation and disease progression. Almost all immune system cells, including neutrophils, B cells, T cells monocytes, macrophages, foam cells, and dendritic cells (DCs), play a vital role in atherosclerotic plaque. Atherogenesis, the normal function of the endothelium, is initially disrupted and, then, cells of the immune system are recruited to the endothelium following increased expression of cell adhesion molecules. Accumulation of immune cells and lipids leads to the formation of a necrotic nucleus. As the disease progresses, smooth muscle cells form fibrous layers, whose rupture results in exposing the necrotic nucleus and thrombosis. Accordingly, the present review was conducted to determine the role of different cells in innate and adaptive immune systems in inhibition and progression of atherosclerosis.

Keywords: atherosclerosis; innate immune system; adaptive immune systems

1 | Introduction

Coronary artery diseases (CAD) is a complex clinical disease which have a variety of serious complications (1). Plaques in atherosclerosis are formed through the accumulation of lipids and various types of immune cells (2). Common symptoms of this disease include chest pain or discomfort as well as sudden manifestations of the disease comprising myocardial infarction and death (3, 4). Underlying mechanisms vary depending on the disease. Atherosclerosis is the main mechanism of CAD. Atherosclerosis can be caused by smoking, high blood pressure, high blood cholesterol, poor diet, obesity, the lack of exercise, and diabetes (5). Atherosclerosis begins with dysfunction of endothelial cells, subendothelial deposition, and altered lipoproteins serving as damage-associated molecular patterns (DAMPs) that activate immune receptors, increase proliferation of immune cells, and induce vascular inflammation (6). It is now established that atherosclerosis is a chronic inflammatory disease occurring in the arterial wall. The oxidation of low-density lipoprotein (LDL) in sub-endothelial play a crucial role in atherosclerotic plaque formation. Oxidized LDL (ox-LDL) is taken up by endothelial cells, smooth muscle cells, and macrophages mainly through several receptors and activated other immune cells (7). In response to endogenous altered structures (e.g., ox-LDL) stimulated both specific and innate immune responses (8). The pathogenesis of atherosclerosis depends on the amount of lipids in the blood because lipid accumulation causes the absorption and activation of immune cells (9). This study aims to describe the function of different cells of adaptive and innate immune systems in progression and suppression of atherosclerosis.

2 | The Role of Innate Immune System in Atherosclerosis

In atherosclerosis, the expression of adhesion molecules (including vascular cell adhesion molecule 3 (VCAM3), intercellular adhesion molecule 1 (ICAM -1), and platelets endothelial cell adhesion molecule 1 (PECAM -1) increase on the surface of endothelial cells. The increased expression of these cells recruits immune system cells towards endothelial cells (10). Various cells from the innate immune system are activated in response to extracellular signals, such as lipopolysaccharides, and intracellular signals, such as ox-LDL. These can lead to progression of atherosclerosis by producing inflammatory mediators (11). In the following, we addressed the role of innate immune system cells in pathogenesis of atherosclerosis.

2.1 | The role of Macrophages in Atherosclerosis

Macrophages, in a process known as phagocytosis, ingest microbes, cancer cells, foreign substances, cellular debris, and anything else not having the type of proteins specific to normal body cells on its surface (12). Macrophages are originated from monocyte cells and located in various body tissues (13). During the course of atherosclerosis, macrophages located under the endothelium vessels take up ox-LDL, and cholesterol derived from ox-LDL is accumulated in these cells. The accumulated cholesterol is pumped from macrophages by the process of cholesterol efflux. If there is a defect in this process, macrophages are transformed into foam cells (14-16). One way to form atherosclerotic plaque is necrosis of macrophages and cholesterol release in the environment (17-19). Ox-LDL activates macrophages through toll-like receptor 4 (TLR4) and increases secretion of tumor necrosis factor- α (TNF- α) from them (20). Macrophages have remarkable flexibility and change their phenotype in response to environmental alteration (21, 22). Overall, the imbalance between macrophage subsets seem to play an essential role in the progress of atherosclerosis (23).

2.1.1 | Classic or M1 Macrophages

Classic or M1 macrophages are activated in response to interferon-gamma (IFN- γ) produced by Th1 lymphocytes. M1 macrophages produce low and high levels of interleukin (IL)-10 and IL-12, respectively (24). This type of macrophages secrete factors, including inflammatory cytokines TNF- α and IL-1 beta (IL-1 β), that increase the differentiation of naive T cells into Th1; Studies have shown that production of acute-phase proteins (APPs), such as C-reactive protein (CRP), leads to formation of M1 macrophages and progression of atherosclerotic plaque (25). According to *in-vitro* investigations, ox-LDL elevates IL-8 expression in M1 macrophages (26). Additionally, M1 macrophages in atherosclerotic plaque cause destruction of the fibrous layer and availability of necrotic nuclei through production of metalloproteinase (MMP). In the human beings, M1 macrophages cause destruction of fibrous layer, plaque rupture, and, eventually, myocardial infarction through production of collagenase (MMP1) and stromelysin-2 (MMP-10) (27).

2.1.2 | Alternative or M2 Macrophages

M2 macrophages support the regulation/suppression of inflammatory responses and wound healing (28). IL-13 and IL-4 cytokines produced by Th2 lymphocytes can induce the production of alternative or M2 macrophages (29). M2 macrophages produce low and high levels of IL-12 and IL-10, respectively, and lead to the increased differentiation of naive T cell lymphocytes into Th2 (30, 31). Although M2 macrophages predominate in the early stages of atherosclerosis, M1 macrophages become predominated as the disease progresses (21). An increased number of M2 macrophages in the initial stages of atherosclerosis may play an atheroprotective role by creating small atherosclerotic lesions. In contrast, M1 macrophages are predominated with development of atherosclerosis, leading to plaque development by expressed NO synthase (iNOS) (32). Cholesterol efflux is impaired and accumulation of cytoplasmic lipids increases in M2 macrophages because of low expression of cholesterol efflux gene (LXR-a). However, M2 macrophages, as compared with M1 macrophages, have less potential to become foam cells (33). In a study, *Isa et al.* showed that ox-LDL activates M1 macrophages while having a lethal effect on M2 macrophages (34, 35). M2 macrophages are categorized into four subclasses, including M2a, M2b, M2c, and M2d.

2.1.3 | M2a Macrophages

M2a phenotype, in which “a” stands for alternative, is produced in response to IL-4 and IL-13. IL-4 (a Th2-produced cytokine) and IL-13 bind to receptor IL-4R α , improving M2 polarization through several pathways, typically Janus kinase 3 (JAK3) and Janus kinase 1(JAK1) signaling, further causing activation and translocation of signal transducer and activator of transcription 6 (STAT6). Other transcription factors involved include interferon regulatory factor 4 (IRF4) and peroxisome proliferator-activated receptor γ (PPAR γ). STAT6, IRF4, and PPAR γ regulate many of the genes associated with mouse M2 macrophages, such as arginase 1 (Arg1), CD206 (or macrophage mannose receptor 1, Mrc1), resistin-like- α (Retnla or Fizz1), and Ym1 (chitins 3-like 3; Chi3l3) (36). M2a macrophages strongly express mannose receptor (MR/CD206), decoy receptor IL-1RII, and IL-1 receptor antagonist. M2a macrophages exhibit an anti-inflammatory phenotype and inhibit production of TNF- α , IL-1 β , granulocyte-macrophage colony-stimulating factor (GM-CSF), IL-10, IL-6, and IFN- γ . M2a macrophages are less involved in phagocytosis and contribute to deposition of extracellular matrix by producing polyamines, collagen, and

transforming growth factor beta (TGF- β). Therefore, M2a macrophages may also be defined as "tissue-repairing" macrophages (37).

2.1.4 | M2b Macrophages

M2b macrophages, although sharing the same characteristics as normal M2 macrophages by exposing to immune complexes, TLR antagonists, and IL-1 receptor ligands, express pro-inflammatory cytokines such as TNF- α , IL-1 β , and IL-6 (38).

2.1.5 | M2c Macrophages

The inhibitory cytokine IL-10 and glucocorticoids induce production of M2c macrophages. Expression of pentraxin-3 (PTX3) and production of anti-inflammatory cytokines, including TGF- β and IL-10, increase in M2c macrophages. Actually, PTX3 is an innate immune receptor which regulates inflammation. Besides, PTX3 is considered as a marker of vascular pathology (such as myocardial infarction and heart failure) (39, 40). M2c macrophages are the main source of PTX3 production (41). There is accumulating evidence suggesting that PTX3 produced by M2c macrophages has a protective role in mice with atherosclerosis (40). M2c macrophages express high levels of Mer receptor kinase (MerTK) that increases effective activity of these macrophages (38, 42). M2b and M2c, known as "regulator macrophages", can improve atherosclerosis by modulating tissue damage caused by long-term activation of M1 macrophages (43). In spite of this, Fleming *et al.*, showed that M2b and M2c macrophages lack the ability to participate in tissue repair because they, unlike M2a macrophages, are unable to synthesize extracellular matrix (44).

2.1.6 | M2d Macrophages

M2d macrophages are induced through TLR antagonists and induce the production of the adenosine A2A receptor. Adenosine inhibits production of TNF- α , IL-1, and IFN- γ through binding to its receptor. Adenosine signaling also induces pro-angiogenic features in macrophages through expressing vascular endothelial growth factor (VEGF) and IL-10 (45, 46).

2.1.7 | Other Macrophages

It has been reported that macrophages have a high degree of flexibility, and other types of macrophages in atherosclerosis plaque include M4, Mox, M(Hb), and Mhem (47-49). Oxidized phospholipids can result in a type of macrophages known as Mox. Nuclear erythroid-2 related factor (Nrf2), a transcription factor, can polarize Mox macrophages. These macrophages exhibit different morphological structure and a combination of biological functions of M2 and M1 macrophages. In fact, Mox macrophages have weak activity in chemotaxis and phagocytosis, and has been shown to constitute approximately 30% of the all macrophages in atherosclerotic lesions established in mice. Nevertheless, the presence of Mox macrophages in human atherosclerotic lesions has not been described until (50). M(Hb) or Hemoglobin-stimulated macrophages are another subset of macrophages present in atherosclerosis (51). Expression of CD163 increases in these macrophages. The Hb/Hp complex stimulates human monocytes to increase CD163 expression and induces the M (Hb) phenotype in atherosclerotic plaques (52). The Hb / Hp complex secretes anti-inflammatory cytokines such as IL-10 (53). Besides, M(Hb) macrophages increase cholesterol efflux by increasing LXRA activity, and prevent formation of foam cells. Raised activity of LXRA additionally stimulates expression of ferroportin, an iron exporter (which is an iron exporter) and decreases the amount of intracellular iron. Decreased levels of intracellular iron lead to the decreased ROS production by M(Hb) macrophages. Finn *et al.*, approved the effect of the decreased ROS production by M(Hb) macrophages in atherosclerotic plaque in humans (51). Platelet-derived chemokine CXCL4 induces the production of M4 cells (54). CXCL4 chemokine (or platelet factor 4 (PF4)) is expressed in numerous cells implicated in atherosclerosis, including monocytes, endothelial cells, and macrophages (55, 56). CXCL4 expression in human atherosclerotic plaques is directly related to severity of the disease (57). M4 and Mhem macrophages have pro-atherogenic and atheroprotective roles through decreased and increased expression of CD163, respectively (58).

2.2 | The Role of Monocytes in Atherosclerosis

Monocytes may differentiate into macrophages and myeloid DCs (59). Monocytes originate from the myeloid lineage in the bone marrow and enter the blood stream, showing a relatively brief half-life in the blood (60). Monocytes have various phenotypes in humans and mice. Mouse monocytes can be identified by expression of CD115, CD11b, CCR2, and CX3CR1 (61, 62). Based on this, two subgroups of mouse monocytes have been identified, the first and second

phenotypes are $\text{Gr1}^+\text{Ly6C}^{\text{high}}\text{CCR2}^+\text{CX3CR1}^{\text{low}}$ and $\text{Gr1}^-\text{Ly6C}^{\text{low}}\text{CCR2}^-\text{CX3CR1}^{\text{high}}$, respectively. Ly6C^{hi} (inflammatory) monocytes are discharged from the blood in a CCR2-CCl2-dependent manner, and are vital in the case of inflammation. Ly6C^{low} (patrolling) monocytes migrate into the bloodstream via CX3CR1-CX3CL1 signaling. Elevated blood cholesterol during atherosclerosis plays a very important role in expansion of $\text{Ly6C}^{\text{high}}$ monocytes. In this regard, investigations performed on mice have indicated that the frequency of $\text{Ly6C}^{\text{high}}$ monocytes is raised in the first days following acute myocardial infarction. However, the frequency of Ly6C^{low} monocytes is raised following acute myocardial infarction and can boost wound healing and regeneration of blood vessels (63-66). Entry of monocytes into atherosclerotic plaques depends on various molecules, of which chemokines and their receptors play a crucial role. CCR2 deficiency in mice has been indicated to dramatically decrease growth of atherosclerotic plaque (63). Deficiency in common ligand of E-selectin and P-selectin on leukocytes has been found to reduce the transfer of $\text{Ly6C}^{\text{high}}$ monocytes toward atherosclerotic plaque in mice (67). Deficiency or blockage of VCAM-1, PECAM-1, ICAM-1, or p-selectin has been shown to decrease adhesion and monocyte rolling and also progression of atherosclerotic plaque in mice (68-70). Filtration of monocyte subtypes to atherosclerotic plaques indicates that these cells can differentiate into diverse subtypes of macrophages. Ly6C^{low} and $\text{Ly6C}^{\text{high}}$ monocytes become macrophages M2 and M1, respectively. Therefore, both monocyte subtypes may play a vital role in inhibition and progression of atherosclerosis (71, 72). In the humans, $\text{CD14}^{\text{high}}$ monocytes are the major subtype of circulating monocytes, and CD14^{low} monocytes are the rare subtype of circulating monocytes. CD14^+ monocytes are determined by expression of CD14 and CD16 (or Fc γ RIII receptor, which has a pivotal role in identification of immune complexes) (73, 74). According to expression of Fc γ RIII (CD16) receptor, monocytes in the humans are classified into three subgroups, including non-classical ($\text{CD14}^+ \text{CD16}^{++}$), intermediate ($\text{CD14}^{++} \text{CD16}^+$), and classical ($\text{CD14}^{++} \text{CD16}^-$) (75-78). $\text{CD14}^{\text{high}}\text{CD16}^-$ monocytes account for about 80-90% of circulating monocytes, and these cells express high and low levels of chemokine receptor CCR2 and CX3CR1, respectively. Nevertheless, $\text{CD14}^+\text{CD16}^{\text{high}}$ monocytes are $\text{CX3CR1}^{\text{high}}$ and CCR2^{low} in terms of expression (73, 79). $\text{CD14}^{\text{high}}\text{CD16}^-$ monocytes produce low levels of TNF- α and high levels of IL-10, whereas $\text{CD14}^+\text{CD16}^{\text{high}}$ monocytes produce low levels of anti-inflammatory cytokines (e.g. IL-10), and high levels of pro-inflammatory cytokines (e.g. TNF- α) (80). $\text{CD14}^+\text{CD16}^{\text{high}}$ monocytes promote atherosclerosis by increased production of TNF- α (74).

The number of classical monocytes is raised in the first days following acute myocardial infarction and decreases after seven days. Additionally, the number of non-classical monocytes is raised through a decrease in the number of classical monocytes (81). Non-classical monocytes are expanded in the patients with CAD, and the number of this subset is directly associated with CAD severity (82), revealing phenotypic complexity of monocytes in the patients with CVD (82, 83).

2.3 | The Role of Neutrophils in Atherosclerosis

Neutrophils, also known as neutrocytes or heterophils, are the most abundant type of granulocytes and makeup 40-70% of all white blood cells in the human body (84). Neutrophils, as the most frequent population of circulating white blood cells (85), are among the first cells to eliminate microbial pathogens by phagocytosis, production of ROS, myeloperoxidase, and various proteolytic enzymes (86). Most neutrophil functions occur in response to signals transmitted through receptors, such as RIG-I-like receptors, NOD-like receptors, C-type lectin receptors, and TLRs (87, 88). Bone marrow-derived neutrophils express high and low levels of CXCR4 and CXCR2, respectively. The CXCL12 chemokine binds to CXCR4 ligand to transmit signals in order to maintain neutrophils in the bone marrow and induces return of old neutrophils to the bone marrow. Elevated cholesterol levels increase the expression of CXCR2 and its ligand CXCL1 and release of neutrophils from the bone marrow. Impaired expression of CXCL12-CXCR4 stimulates neutrophils and increases the size of atherosclerotic plaque (89). Studies on mouse models of atherosclerosis have shown a direct relationship between the number of peripheral neutrophils and size of the atherosclerotic lesion. Importantly, it is predicted that regulating factors of neutrophil homeostasis are involved in progression of atherosclerotic disease by influencing the number of peripheral neutrophils (89). During inflammatory processes, the number of neutrophils increases in the bloodstream, and the IL-8 cytokine attracts these cells to necrotic nucleus and exerts its effective function (90). An increase in neutrophils influx and inflammatory activity of these cells in atherosclerotic plaques may play a vital role in progression of atherosclerosis through secretion of various mediators, including PTX3, ROS, neutrophil elastases, MPO, leukotriene B4, neutrophil extracellular traps (NET), connexin, IFN- γ , and MMPs (91). For example, leukotriene B4 and its receptor BLT1 activate neutrophils and recruit them to atherosclerotic plaques (92). Neutrophils in human atherosclerotic lesions

produce PTX3 in response to inflammatory signals (e.g., bacterial products, TNF- α , and IL-1) (41). The PTX3 protein also increases inflammation, chemotaxis, phagocytosis, tissue damage, and, finally, progression of atherosclerosis by activating classical complement pathway (93, 94). Primary granules in neutrophils contain a good source of MPO enzyme (95), which increase the production of ox-LDL (96). Circulating neutrophils have lower MPO, while elevated MPO levels of plasma in the patients with hyperlipidemia reveal depletion of neutrophil granules (97). MPO can also result in the accumulation of cholesteryl ester in macrophages, and the formation of foam cells can develop atherosclerosis. Moreover, MPO may attract neutrophils to necrotic nucleus in atherosclerosis (98). Several studies have shown an increase in the circulating levels of PTX3 and MPO in the patients with atherosclerosis, and an increase in these two markers has been mentioned as prognostic markers for acute and chronic heart failure, which may be a major risk factor for CVD in the humans (99, 100). Secondary and tertiary granules of neutrophil containing large amounts of MMP-2 and MMP-9 destroy endothelial cells by influencing collagen type IV, which is a pivotal agent in the endothelial cell membrane (101). NETs are among novel mechanisms of neutrophils against microorganisms. NETs include core components, DNA, primary granules (elastase and MPO), secondary granules (PTX3 and lactoferrin), and third granules (MMP-9) (90). Inflammation in atherosclerosis can cause the release of NETs from neutrophils (102). NETs lead to thrombosis due to deposition of fibrin protein (103). Furthermore, NETs cause apoptosis of endothelial cells and destabilization of atherosclerotic plaque (104).

Neutrophils are also involved in development of atherosclerosis by secreting azurocidin, proteinase 3, and defensin. Azurocidins can lead to activation of protein kinase c and a raise in the expression of VCAM-1 and ICAM-1, followed by recruitment and attracting the monocytes (105). Besides, proteinase 3 is able to increase expression of cell adhesion molecules and adhesion of neutrophils and monocytes to endothelial cells (106). Defensins not only intensify expression of chemokines and cell adhesion molecules, but also impair endothelial function through increased production of oxygen free radicals (107, 108).

2.4 | The Role of Mast Cells in Atherosclerosis

Mast cells, also known as a mastocyte, is a migrant cell in connective tissue containing many granules rich in heparin and histamine. Notably, Mast cells are a type of granulocyte derived

from the myeloid stem cell that is a part of neuroimmune and immune systems (109). Mast cells, as a type of innate immune cell agents, play a fundamental role in allergic patients. Mast cells are detected not only in all the vascular tissues, nerves, near blood vessels, mucus-producing glands, smooth muscle cells, and hair follicles (110), but also in carotid arteries of healthy individuals as well as in advanced and early atherosclerotic plaques(111). The presence of mast cells in atherosclerotic plaques indicates their potential role in the disease (111). *In-vivo* studies revealed that mast cell deficiency significantly decreases atherosclerosis in mice (112). Activation of mast cells by allergens or microorganisms releases inflammatory mediators from the granules. Classic mast cells are activated by binding FcεRI receptors to immunoglobulin E (IgE) with high affinity (113). Reduced expression of FcεR1a has been found to lead to a dramatic decrease in the lipid deposition in aortic arch of mice, which might be due to inactivation of mast cells by FcεR1a and severe decline in release of inflammatory mediators, which can decrease the number of inflammatory cells such as macrophages and T cells in atherosclerotic plaque (114, 115). Mast cells are also activated by different mechanisms, such as IgG or immune complexes (116). In animal models of atherosclerosis, the immune complex of oxLDL-IgG has been observed in plaques, and the presence of these complexes may be implicated in activation of mast cells (117).

In-vitro studies demonstrated that the oxLDL-IgG immune complex secretes inflammatory cytokines (such as IL-8 and TNF-α) and some other mediators (e.g., histamine and tryptase) from human mast cells (118). TLRs are among other pathways related to activation of mast cells. Previous studies have demonstrated that TLR4 antagonist decrease activity of mast cells, presumably indicating that mast cells are activated by TLR4. Additionally, activation of mast cells by TLR4 induces apoptosis in vascular smooth muscle cells of atherosclerotic plaques (119, 120). Furthermore, the receptor of complement system components C3a and C5a can be activated mast cells in the plaque (121). Mast cells can be involved in onset of atherosclerosis through diverse mediators, such as histamine (which raises vascular permeability) (122). Likewise, mast cells can play a considerable role in the progress of atherosclerotic plaques by secreting different cytokines. Based on studies conducted on a mouse model of atherosclerosis, deficiency in IFN-γ or IL-6 produced by mast cells is associated with a decrease in plaque size (112). Secretion of inflammatory mediators by mast cells induces expression of adhesive molecules, VCAM-1, E-selectin, P-selectin, and ICAM-1 in endothelial cells and enhances recruitment and adhesion of leukocytes to vascular endothelium. This occurrence may indicate

that function of mast cells in atherosclerosis is dramatically related to secretion of cytokines (123). Histamine is another compound found in the granules of mast cells. Histamine levels are considerably raised in the patients with atherosclerosis compared to healthy individuals, which may indicate the role of histamine in progression of atherosclerosis (124). *In-vitro* investigations have implicated that secretion of histamine and protease mediators by mast cells could induce apoptosis of macrophages. Furthermore, protease secretion can lead to apoptosis of endothelial cells and smooth muscle cells of the vascular wall (125). *In-vivo* experiments in mice have shown that a noticeable increase in the secretion of protease tryptase raises formation of carotid plaque and makes the carotid artery to become narrow, whereas a decrease in the secretion of tryptase decreases the size of carotid plaques (126). Studies on mouse models of atherosclerosis have indicated that tryptase increases the expression of IL-8 chemokines, which have an ability to attract monocytes and neutrophils to endothelial cells, and monocyte chemoattractant protein-1(MCP-1) and may have a striking role in the development of atherosclerosis (127). In the human beings, tryptase promotes the formation of foam cells by repressing activation of nuclear receptor liver x receptor (LXR) – α , and reverse transport of cholesterol (128, 129). Moreover, tryptase could boost angiogenesis through degradation of collagen type IV, vitronectin, and fibronectin (130). Inhibition of protease chymase in mice has been shown to diminish plaque development and size of necrotic nucleus (131). Chymase destabilizes the plaques by stimulating apoptosis of smooth muscle cells (SMC) and degradation of vitronectin and fibronectin (132). Heparin is another compound present in mast cell granules that may increase the formation of foam cells (133). Therefore, mast cells cause plaque instability and exposing of necrotic nucleus through secretion of various mediators, such as histamine, heparin, proteases (tryptase and chymase), and multiple cytokines (such as IL-6, IL-8, MCP1, TNF- α , and IFN- γ) (134).

2.5 | The Role of Natural Killer Cells (NKs) in Atherosclerosis

Natural killer cells (NK cells) or large granular lymphocytes (LGL) are a type of cytotoxic lymphocytes which are vital to the innate immune system (109). NKs recognize pathological changes in the tissues through inhibitory and activating receptors. Decreased and increased expression of MHC-I molecules leads to utilization of activating receptors and inhibitory receptors, respectively (135, 136). The interplay of NK cells with the infected cells can trigger activating receptors, secreting perforin, granzyme, and anti-inflammatory cytokines (e.g., IL-1 β ,

TNF- α , and IFN- γ), and exciting inflammation and direct destruction of target cells (135-137). In the humans, NK cells have been demonstrated to exist in atherosclerosis lesions (138). NK cells are leading sources for production of IFN- γ . Moreover, various studies have shown that a decrease in the IFN- γ levels significantly reduces atherosclerotic plaques in mice, which may indicate the role of NK cells in atherosclerosis (139, 140). In atherosclerotic plaques, there are different chemokines for attracting innate immune cells. For example, CX3CL1 and MCP-1 chemokines play a vital role in attracting NK cells to necrotic nucleus (141, 142). Overall, the function of NK cells in atherosclerosis has not been identified.

2.6 | The Role of DCs in Atherosclerosis

DCs are antigen-presenting cells (APCs) in the mammalian immune system. Their chief function is antigen processing and its presentation on the cell surface to T cells of the immune system. They act as messengers among innate and adaptive immune systems (143). DCs are derived from common precursor CD34⁺ in the bone marrow and placed in various tissues of the body (144). DCs have a pivotal role in both innate and adaptive immune systems (145). Various receptors (integrins, TLRs, scavenger receptors, Fc receptors, and C-type lectins) are expressed to uptake antigens at the surface of DCs. For example, TLR4 plays a vital role in the onset and development of atherosclerosis (146). Lipopolysaccharide (LPS) from Gram-negative bacteria, ox-LDL, and heat-shock proteins (HSPs) are detected by TLR4, leading to activation of signaling transduction (147, 148). However, there are studies representing that ox-LDL may inhibit TLR4 signaling (149). Activation of TLR4 at DC levels via *Chlamydia pneumonia* may be involved in the development of atherosclerosis (150). Identification of nucleic acids in infectious pathogens by TLR7, TLR8, and TLR9 may be involved in the destabilization of atherosclerotic plaques (151, 152). Antigens processed by DCs, accompanied by MHC molecules, are presented to T cells (153). Expression of chemokine receptors, MHC molecules, cytokines (TNF- α and IL-12), and costimulatory molecules increases during DC maturation, resulting in the expansion of different CD4⁺ T cell subtypes, such as T helper (Th) cells (Th17, Th1, and Th1) (145, 154). Secretion of CCL12 and CCL5 chemokines and expression of adhesive molecules, E-selectin, P-selectin, and integrins induce DC recruitment to atherosclerotic plaques (155, 156). The CX3CL1 chemokine seems to be an important agent for infiltrating DCs into atherosclerotic plaques, because deficiency of Fractalkine (FKN) receptor

leads to a decrease in the number of DCs and atherosclerosis (157). Studies performed on mouse models manifested that the decreased number of DCs reduces lipid accumulation and the number of foam cells, and ultimately reduces the size of atherosclerotic plaques (158). *In-vitro* investigations revealed that ox-LDL increases expression of CD83, HLA-DR, CD36, and CD86 molecules through activation and maturation of human DCs (159, 160). Ox-LDL also diminishes the expression of chemokine receptors CCR7 and CCL21 in DCs, which can decrease the migration of these cells (161). Interactions of DCs with NK cells enhance production of IL-12 and IFN- γ and may have a role in the formation and progression of atherosclerosis plaques (139). The presence of classical myeloid DCs (mDCs) and plasmacytoid DCs (pDCs) has been approved in atherosclerosis plaques. pDCs are activated by the help of the TLR9 ligand, and increase the expression of tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) at the level of CD4⁺ T cells, apoptosis of vascular smooth muscle cells (VSMCs), and, finally, instability of atherosclerotic plaque, by secreting type I interferon (IFN) (162). Clinical studies demonstrated a decrease in the population of pDCs and mDCs in the peripheral blood from the patients with atherosclerosis compared with the control group (163).

This decrease in the number of DCs derived from blood circulation may be in light of the increased presence of DCs in atherosclerotic plaques, showing the possible role of these cells in the development of atherosclerotic plaques (163, 164). Activation of DCs destabilizes atherosclerotic plaques in two following ways; in the first way, DCs can differentiate naive T cells into different subsets of effector T cells. Activation of cytotoxic T cells causes rupture and exposing of necrotic nucleus of plaques. In the second way, DCs destroy extracellular matrix connections and rupture plaques by producing some proteases, such as metalloproteinases. The interaction between mDC, pDC, and other immune cells activates other immune cells, rupture of plaques, eventually leading to atherosclerosis. DCs under certain physiological or pathological conditions may express anti-inflammatory cytokines and inhibitory receptors and cause tolerogenic immune responses. Different environmental stimuli help DCs suppress immune responses by producing regulatory and anergic T cells. Regulatory DCs can present antigens but show a different expression profile, containing lower levels of stimulatory molecules and inflammatory cytokines and higher levels of anti-inflammatory cytokines. In addition, regulatory DCs are resistant to maturation signals (165). Regulatory DCs enhance immune tolerance by decreasing expression of stimulatory molecules, enhancing expression of inhibitory molecules

(e.g. indoleamine 2,3-dioxygenase (IDO)), promoting induction of anti-inflammatory cytokines (e.g., TGF- β and IL-10), and inhibiting production of inflammatory cytokines (e.g. TNF α and IL-12) (166). Regulatory DCs interfere with anergy and depletion of inflammatory T cells as well as production and proliferation of atheroprotective Tregs (167, 168). Accumulation of apolipoprotein B100 (ApoB100), LDLs, and ox-LDLs in the blood vessel wall attracts immune cells and gives rise to chronic inflammation. Loading DCs with ApoB100 and ox-LDL reduces production of non-inflammatory cytokines (169). DCs may be useful in inducing tolerance against autoantigens in the walls of arteries and in creating new treatment methods for atherosclerosis by forming a cell population of CD4⁺ Foxp3⁺ Tregs. Tregs can inhibit atherosclerosis by secreting inhibitory cytokines, TGF- β and IL-10 (170). Overall, DCs may play a dual role in the disease.

3 | The Role of Specific Immunity in Atherosclerosis

As mentioned above, atherosclerosis, characterized by inflammation, lipid accumulation, cell death, and fibrosis, is a chronic inflammatory disease, which causes arterial plaque. Identification of T cells in human atherosclerotic plaques has indicated the possible role of the specific immune system in the disease. Recently, a wide variety of researches have confirmed the role of specific immunity in atherosclerosis. In the following, we represented a summary regarding the role of specific immunity in atherosclerosis.

3.1 | The Role of T Lymphocytes in Atherosclerosis

T cells are a lymphocyte, derived from the thymus gland which play a pivotal role in the immune response. T cells can be discriminated from other lymphocytes by the presence of a T-cell receptor on cell surface (171). APCs such as macrophages, DCs, and B cells trigger T lymphocyte responses. T cell activation begins with binding of TCR to the MHC molecule and binding of costimulatory molecules such as CD28 and CD40L (172). T cells act as a source of inflammatory cytokines such as IL-6, IL-12, and TNF- α .

Ox-LDL is produced in the arterial wall by increased oxidative stress, and may activate T cells in atherosclerotic plaques. Specific T cells, induced against ox-LDL autoantigens, are formed over the process of atherosclerosis (173), and the presence of these cells is essential for production of

specific IgG autoantibodies against ox-LDL (174, 175). Moreover, transfer of these specific T cells can exacerbate atherosclerosis (176). Therefore, immune responses by B and T cells play a major role in inflammation and formation of atherosclerotic plaques. The β 2-glycoprotein I is another autoantigen involved in atherosclerosis. Indeed, β 2-glycoprotein I is a phospholipid-binding protein found in the human atherosclerotic plaques (177). Immunization of mice with defective low-density lipoprotein receptor (LDLR) has been found to accelerate the formation of atherosclerotic plaque (178).

3.1.1 | CD4⁺ T Cell Subsets in Atherosclerosis

CD4⁺ T cells have different subtypes, including Th1, Th2, Treg, Th9, Th22, Th17, and CD28^{null} T cell. T helper cells, with high flexibility, are converted into different subsets of CD4⁺ T cells depending on the environmental conditions.

3.1.1.1 | The Role of Th1 Cells in Atherosclerosis

Th cells have a vital role in adaptive immunity. Conversion of T lymphocytes into Th1 depends on neighboring inflammatory cytokines, those existing around T lymphocytes. IL-12 activates the signal transducer and activator of T-box transcription factor (TBX21) (or T-bet) and transcription-4 (STAT-4) transcription factors, which are necessary for the formation of Th1 cells. Th1 cells activate other cells of the immune system by secretion of different cytokines especially IFN- γ . Th1 cells are the most abundant population of T cells present in atherosclerotic plaques (173, 179). The absence of IFN- γ or IFN- γ receptors in mouse models suggests the decreased susceptibility of atherosclerosis (180-182). Furthermore, injection of recombinant IFN- γ causes an increase in the size of atherosclerotic plaques (183). Studies in mice with defective LDLR have shown that the formation of atherosclerotic plaques in the mice receiving bone marrow transplantation from donor mice with a failure in IFN- γ is more severe compared with those receiving from healthy donor mice (184). In addition to Th1 cells, IFN- γ is released by NKT cells, NK cells, macrophages, and muscle cells (185). IFN- γ prompts the formation of atherosclerotic plaques through lipid uptake by macrophages and the formation of foam cells, attraction of macrophages and T cells, the increased expression of MHC-II, as well as increased activation of APC and release of Th1 cytokine (186, 187). Moreover, IFN- γ may cause rupture of atherosclerotic plaques by decreasing proliferation of SMCs, decreasing synthesis of collagen,

and increasing production of extracellular matrix-degrading proteins (188-190). IFN- γ may have an anti-atherosclerotic effect by decreasing concentrations of serum cholesterol, scavenger receptor A, suppression of LDLR-related protein, MMP9, CD36, and inhibition of lipoprotein lipases (190). A disorder in transcription factor T-bet in mouse models has been shown to decrease the atherosclerosis severity (191). Insufficiency of MHC class II-associated invariant chain (CD74) has been reported to cause a decrease in the activation of T cells and atherosclerosis in mice (192). Interplay between the CD40L on activated T cells and the CD40 receptor on APCs results in initiation of inflammation (193). Inhibition of the activation pathway of CD40-CD40L in mouse models has been suggested to have the potential to decrease the size of atherosclerotic plaques (194-196). There are some studies demonstrating the role of IL-18 in atherosclerosis. IL-18 deficiency in mice can lead to a decrease in the activity of Th1 cells and size of atherosclerotic plaque (197). IL-18, IFN- γ , and IL-12 are Th1 cell response-related cytokines, which can accelerate atherosclerosis, showing their important potential as appropriate targets for the treatment of atherosclerosis.

3.1.1.2 | The Role of Th2 Cells in Atherosclerosis

Th2 cells are implicated in allergic diseases (e.g., asthma and atopy), and their role in atherosclerosis depends on the stage and place of disease and release of cytokines. In addition, the role of Th2 cells in atherosclerosis has not been exactly determined, and its differentiation is initiated with IL-4. This cytokine can activate STAT6 and, subsequently, induce the GATA3 transcription factor. GATA3 can inhibit production of IFN- γ and increase production of IL-5 and IL-4. Th2 cells also produce IL-25, IL-5, IL-9, and IL-13, leading to the production of antibodies by B cells. In atherosclerosis-prone mice, the response of TH2 cells has been shown to have a protective role against development of early atherosclerotic plaques (198). It seems that Th2 cells have a protective role against atherosclerosis because they suppress Th1 cells (having a pro-atherogenic role), while atherosclerosis severity has been found to decrease in IL-4^{-/-} ApoE^{-/-} mice (199, 200). IL-4 has pro-atherogenic effects, including activation of mast cells, apoptosis of SMCs, plaque rupture, increased protease production, and decreased collagen production (201). Furthermore, it can stimulate elastase (MMP-12), that can damage arterial walls and increase the formation of the aneurysm (202). The role of IL-13 in atherosclerosis has not been well understood, while IL-5 has an opposite effect in comparison with IL-4 and has a protective role

in atherosclerosis. Receiving bone marrow transplantation from mice with a disorder in producing IL-5 has been reported to lead to plaque formation (203). Anti-atherogenic characteristics of IL-5 may be due to the production of protective antibodies (204). IL-33 is a member of IL1-family cytokine that also incorporates IL-18 and IL-1 β . IL-33 was first identified to have a relationship with Th2 cell differentiation in 2005 (205). It has been declared that the injection of IL-33 to mice can elevate the concentration of Th2 cytokines in lymph nodes and serum as well as the production of atheroprotective IgM-type anti-oxLDL antibodies, leading to a decrease in the expansion of atherosclerotic plaque (206). Accordingly, the increase in the Th2 cell response cannot be assumed as an appropriate method for the limitation of vascular inflammation and a decrease in the severity of atherosclerosis.

3.1.1.3 | The Role of Th17 Cells in Atherosclerosis

Th17 helper cells are a subtype of T helper cells which are different from Th2 and Th1 subtypes in terms of growth and development. Th17 cells have a special role in fighting against extracellular pathogens and fungal infections. Th17 cells are effective in defending against bacterial and fungal pathogens and produce IL-17 (207, 208). TGF- β along with IL-4 causes the formation of Th9 but TGF- β along with IL-6 causes the formation of Th17. IL-6 activates the signal transducer and activator of transcription 3 (STAT3), which is necessary for the expression and function of retinoic acid-related orphan receptor gamma t (ROR γ t) (209). While unable to directly induce differentiation of Th17, TGF- β suppresses transcription factors needed for differentiation of Th1 and Th2. Production of both Th17 and Th9 subtypes is suppressed through induction of CD4⁺ Foxp3⁺ T cells, which have a protective role against atherosclerosis. Results from a study indicated that IL-17 is generated by T cells in human vessels and IL-17 along with IFN- γ results in an inflammatory response in VSMCs (207). Th17 cells play a role in autoimmune diseases, but their role in atherosclerosis is not completely understood (210). There are some studies performed on the role of IL-17 in atherosclerosis, showing that a failure in signaling of IL-17 receptor (IL-17R) in mice causes a decrease in the size of atherosclerosis plaque and serum concentration of anti-oxLDL IgG antibodies (211). *Erbel et al.* in a study showed that mRNA expression of the IL-17 cytokine increases in symptomatic carotid plaques, unlike asymptomatic plaques; they concluded that mRNA expression of the IL-17 cytokine is related to lipid-rich, complicated, and unstable plaques (212). The nuclear receptor of ROR- γ t is

necessary for differentiation of Th0 to Th17 (213). Other crucial transcription factors for Th17 differentiation include Aryl hydrocarbon receptor, runt-related transcription factor 1 (RUNX1), STAT3, and ROR- α (214, 215). IL-23 and IL-21 are required for the maintenance and proliferation of Th17, respectively. IL-17A, IL-21, IL-22, and IL-17F are produced by Th17 cells. Both IL-4 and IFN- γ have been shown to suppress differentiation of Th17 in different mice models (207).

3.1.1.4 | The Role of Th9 Cells in Atherosclerosis

Th9 cells are a subset of T cells, which are commonly along with the Th2 phenotype. These cells are the main source of IL-9. The IL-9 production in Th9 cells is induced by TGF- β and IL-4 and repressed by IFN- γ (216). According to the study, the plasma level of IL-9 is higher in patients with atherosclerosis than healthy individuals. Other studies have shown that the IL-9 level is higher in human atherosclerotic plaques than vascular samples from healthy individuals (217). Besides, other studies suggested that the plasma levels of IL-9 are raised in the patients with acute coronary syndrome compared with healthy individuals, although there are presently no reports about change in the amount of Th9 cells in the blood (218). It is not yet known whether Th9 cells contribute to progression or inhibition of atherosclerosis.

3.1.1.5 | The Role of Th22 Cells in Atherosclerosis

Th22 cells have a unique property as they express IL-22 and the major transcription factor of the aryl hydrocarbon receptor, but not IL-17 and IFN- γ (219). The cytokines responsible for stimulation and differentiation of Th22 cells are unknown (220). In addition, it is not known whether Th22 cells have an anti-atherogenic or pro-atherogenic role. The level of Th22 cells in the blood from patients with acute coronary syndrome was reported to increase compared with the healthy control group (218, 221). Other studies revealed that the plasma level of IL-22 is higher in patients with symptomatic atherosclerosis than those with asymptomatic atherosclerosis (218, 221, 222). Results from a study on the mice susceptible to atherosclerosis with IL-22 deficiency indicated that IL-22 decreases atherosclerosis by suppressing pro-atherogenic intestinal microbes (223).

3.1.1.6 | The Role of T_{FH} Cells in Atherosclerosis

T follicular helper (TFH) cells are a subset of CD4 T cells migrating to B-cell follicles after activating and promoting formation of germinal center and switching of B-cell isotype, which requires B cell lymphoma 6 (Bcl6) transcription factor (224). Moreover, TFH cells are possible to have a pro-atherogenic profile. According to the evidence, atherogenic media increases autoimmune responses of CXCR3⁺ T_{FH} cells in the mice susceptible to atherosclerosis (225). Nevertheless, in mice fed a high-cholesterol diet, B cells of marginal zone repressed the T_{FH} cell response, and thereby, limited progression of atherosclerosis (226). Furthermore, in some mice, TFH cells having pro-atherogenic activity in germinal centers could be regulated by a subset of CD8⁺ T cells that have a regulatory function (227). The pathway related to inducible T cell co-stimulator (ICOS) and its ligand (ICOSL) is vital for differentiation and sustaining of TFH cells. It has been reported that blocking the ICOS-ICOSL signaling pathway in Apoe^{-/-} mice decreases the progression of atherosclerosis and leads to a reduction in the number of T_{FH} cells in secondary lymphoid organs (228). Interestingly, TFH cells can be derived from Tregs. These cells have an pro-atherogenic profile and can increase the severity of atherosclerosis in mice. In the patients with CAD, the plasma level of IL-21 (the major cytokine in T_{FH}) is negatively correlated with expression of forkhead box P3 (FOXP3) in Treg cells (228).

3.1.1.7 | The Role of CD28^{null} T Cells in Atherosclerosis

CD4⁺CD28^{null} T cells are characterized by the absence of CD28 expression that is the major costimulatory receptor for CD4⁺ T cells. CD28^{null} T cells have cytotoxic and inflammatory properties. Attractively, CD28^{null} T cells are detected exclusively in the humans and non-human primates, but not in mice (229). Patients with acute coronary syndrome have more CD28^{null} cells in their blood compared with healthy people, and their CD28^{null} T cells are resistant against apoptosis (230, 231). Additionally, unstable plaques may develop by the effects of T cells, such as CD28^{null} T cells. Besides, some of the CD28^{null} T cell subsets react to HSP60 and HSP70, two proteins that are likely to be among atherosclerosis antigens (232, 233). Among the patients with end-stage renal disease, individuals with atherosclerosis have more CD28^{null} cells than those without atherosclerosis (234).

3.1.1.8 | The Role of Tregs in Atherosclerosis

CD4⁺FoxP3⁺ regulatory T cells are a subtype of T cells which can repress immune system responses (235). Indeed, Tregs are a subtype of T cells playing a role in inhibiting immune reactions, maintaining hemostasis, and inducing self-tolerance. Tregs are divided into two main subsets, including natural Tregs (nTregs) and induced Tregs (iTregs). The nTregs detect self-antigens, after being developed in the thymus. These cells are also recognized by expression of CD4, CD25 surficial receptors (a subunit of IL-2 receptor), and FOXP3 transcription factor (236). The iTregs are derived from nTregs, in the presence of IL-10. Those iTregs formed by IL-10 are known as TR1 (237). It has been indicated that transmission of CD4⁺CD25⁺ T or TR1 cells can lead to a decrease in the expansion of atherosclerotic plaque in mouse models (238-240). Those iTregs, stimulated by TGF- β are known as Th3 cells (241). The mechanism related to suppression of immune system are developed through the activity by Treg cells (235). TGF- β and IL-10 are two main cytokines of Treg cells which are responsible for many effects of Treg cells and have strong anti-atherogenic activities. Inactivation of TGF- β or using neutralizing antibodies has been shown to accelerate the development of plaque and vascular inflammation in mouse models of atherosclerosis (242-244). The presence of Tregs in atherosclerosis was shown first by the immunohistochemistry method and following in the aorta of mice using polymerase chain reaction (PCR) technique (245, 246). There is no evidence demonstrating that the presence of Treg cells has a direct role in the development of the disease. Chemokines play a role in recruitment of CD4⁺FoxP3⁺ regulatory T cells. There are documents showing that the size of atherosclerotic plaque is smaller in CXCL10^{-/-}ApoE^{-/-} and CXCL3^{-/-}ApoE^{-/-} mice, where there is a less number of T cells and infiltrated macrophages, and an increased number of CD4⁺FoxP3⁺ regulatory T cells (246, 247). Furthermore, some studies conducted on mouse models revealed that the elimination of Tregs by anti-CD25 antibodies elevates the size of atherosclerotic plaque (239). Tolerance against potential antigens in atherosclerotic plaque has an indispensable role and, according to our knowledge, Tregs play a critical role in creating tolerance. There is evidence showing that mice with a low number of Tregs in spleen suffer from systemic tolerance defect compared with non-atherosclerotic mice (238, 248). Although they can inhibit the activation of T cells and, subsequently, decrease the progression of atherosclerosis, Tregs are likely to form remarkable protection against atherosclerosis by targeting APCs. Tregs can lead to differentiation of macrophages into the anti-inflammatory M2 phenotype as well as inhibit the formation of foam cells *in-vitro* and differentiation of inflammatory M1 phenotype (249, 250).

3.1.2 | The Role of CD8⁺ T Cells in Atherosclerosis

CD8⁺ T cells identify antigen peptides presented by MHC class I. These cells can create cytotoxic T cells that have the ability to kill virus-infected cells and other abnormal cells (e.g., cancerous cells) using various cytotoxic pathways. In patients with CAD, the blood level of cytotoxin-producing CD8⁺ T cells increases compared with healthy individuals, and CD8⁺ T cells in atherosclerotic plaques are plentiful in the humans and mice (251, 252). In advanced stages of human atherosclerosis, the number of CD8⁺ T cells is higher than CD4⁺ T cells, which are principally found in the fibrous cap regions (252, 253). In the majority of studies related to atherosclerosis, similar to CD4⁺ T cells, the specific antigen of CD8⁺ T cells associated with atherosclerosis plaque is unknown. Experimental researches highlighted both atheroprotective and pro-atherogenic roles of CD8⁺ T cells (254). The cytotoxic activity of CD8⁺ T cells, relative to lesion-stabilizing cells (e.g., VSMCs), and production of inflammatory cytokines by CD8⁺ T cells may amplify inflammatory responses in atherosclerotic plaques and intensify the progression and instability of lesion. In contrast, the cytotoxic activity of CD8⁺ T cells, relative to APCs, may limit the progression of atherosclerosis. MHC class I-dependent cytotoxic CD8⁺ T cells contribute to creating inflammation in atherosclerotic plaques and necrotic nuclei (254). Furthermore, a decrease in the number of CD8⁺ T cells by antibodies was found to decrease the severity of atherosclerosis in mice, showing the pro-atherogenic role of CD8⁺ T cells (255, 256). In these cells, the production of granzyme B and IFN- γ is higher than CD8⁺ T cells in non-atherosclerotic mice (256). Results from a study on Ldlr^{-/-} mice fed with high-fat diets revealed that CD8⁺ T cells develop atherosclerosis by regulating the level of peripheral monocytes, through production of IFN- γ (255). However, findings from another study revealed that IFN- γ produced by CD8⁺ T cells has no role in atherosclerosis in Apoe^{-/-} mice fed with high-fat diet (257). A number of experimental researches manifested that CD8⁺ T cells can play an atheroprotective role. Some studies demonstrated regulatory functions of CD8⁺ T cells in atherosclerosis. In the Apoe^{-/-} mice, CD8⁺ T cells were found to exert their atheroprotective effects by ApoB-related peptide (p210) (258). In the Apoe-deficient mice, a subset of CD8⁺ T cells, called CD8⁺ Treg, was reported to decrease the severity of atherosclerosis and potentially atherogenic IgGs by blocking the ICOS-ICOSL signaling among B cells and T_{FH} cells of germinal centers (227).

3.2 | The Role of B Lymphocytes in Atherosclerosis

B cells (B lymphocytes) are a type of white blood cells in the lymphocyte subtype, which play a vital role in the humoral immunity component of the adaptive immune system by secreting antibodies. Moreover, B cells present antigens, which are categorized as professional APCs, and secrete cytokines (259). B cells can be divided into two distinct categories, including B1 and B2. B1 cells account for a minute percentage of B lymphocytes and are derived from the committed precursors of fetus or infant. Actually, B1 and B2 cells are predominant in fetus and adults, respectively. B2 cells incorporate marginal zone B cells and follicular B cells, and B1 cells include B-1b and B1a cells. B1 cells cooperate in innate immunity by producing normal IgM antibodies. B2 cells are the principal population of B cells produced in the bone marrow. After activation by specific antigens, B2 cells produce IgG antibodies and differentiate into plasma cells. Other functions of B2 cells include cytokine production, organization of lymphoid tissues, and antigen presentation (260, 261). B cells are detected in atherosclerotic plaques in humans and mice. B cells, which help in formation of lymphoid follicles, can play a role in atherosclerosis (262, 263). IgG and IgM antibodies are present in atherosclerotic plaques at all stages of disease progression (264). Specific antibodies against ox-LDL were observed in humans and mice (265). In the mice, the anti-oxLDL IgM antibody was demonstrated to provide protection against atherosclerosis, showing that humoral and B cell immunity has anti-atherogenic properties (266). It has been reported that the administration of IgG1 antibodies against ox-LDL epitopes in *Apoe*^{-/-} mice decreases the severity of atherosclerosis (267). Moreover, bone marrow transplantation from B cell-defective mice to *Ldlr*^{-/-} mice (compared to with bone marrow transplantation from healthy mice) was shown to increase the severity of atherosclerosis, which indicates the important role of B cells in the development of atherosclerosis (268). Regulatory B cells are a subset of B lymphocytes, producing IL-10, and are considered as regulatory cells (269). Breg cells seem to play an atheroprotective role. The incidence of atherosclerosis in null mice for IL-10 has been found to be significantly higher than the control group (244). IL-10 decreases the risk of atherosclerosis by exerting apoptosis, modulating lipid metabolism, and inhibiting production of pro-inflammatory mediators (270). Breg cells express Fas ligand, actually inducing apoptosis in target cells, presumably representing an essential mechanism for suppressing pro-inflammatory immune responses in atherosclerotic plaque (270, 271). Surface markers of Breg cells have not been fully identified, so their role in atherosclerosis is unclear yet.

4 | Other T Lymphocytes in Atherosclerosis

4.1 | The Role of Natural Killer T (NKT) Cells in Atherosclerosis

Natural killer T (NKT) cells are classified into two subgroups including invariant NKT (type I or iNKT) cells whose TCRs are not very variable, and type II NKT cells, which have highly variable TCRs. To date, iNKT cells are the only NKTs studied in atherosclerosis. The iNKT cells can be activated by TCR interplay with antigens presented by CD1d molecules, which include antigen glycolipids presented in APCs (272). Some glycolipids have a microbial origin (273). The iNKT cells, similar to CD8⁺ T cells, can express cytotoxic proteins such as granzyme and perforin (274). Most researches performed on mice indicated that iNKT cells play a pro-atherogenic role (275). The iNKT cells can activate immune cells in plaque, by secreting cytokines and promoting atherosclerosis development.

4.2 | The Role of $\gamma\delta$ T Cells in Atherosclerosis

$\gamma\delta$ T cells, unlike $\alpha\beta$ T cells, fail to identify specific antigens (276). There are a few studies conducted on mice about $\gamma\delta$ T cells in atherosclerosis (277-279). Results from the first study conducted in 1993 showed the presence of $\gamma\delta$ T cells in human atherosclerotic plaque (280). Interestingly, the amount of intracellular cholesterol in $\gamma\delta$ T cells regulates their function activation, and proliferation (277). Moreover, a study accomplished on mice revealed that $\gamma\delta$ T cells are a rich source of IL-17 and can regulate atherosclerosis by producing IL-17 (281). According to the documents, progress of early atherosclerosis in Apoe^{-/-} mice, whose $\gamma\delta$ T cells are genetically defective, is similar to Apoe^{-/-} mice $\gamma\delta$ having T cells (278). To date, the definite role of this T cell subset has not been determined in atherosclerosis.

CONCLUSION

The present review was conducted to investigate how immune system cells influence atherosclerosis. Studies demonstrated that LDLs stimulate specific and innate immunity in atherosclerosis. Atherosclerosis is exacerbated by pro-inflammatory responses mediated by

immune cells. However, some cells, such as M2 macrophages, tolerogenic DCs, Bregs, and Tregs reduce inflammatory activity of the immune system in atherosclerotic lesions. Therefore, these regulatory factors can be useful in the treatment and production of therapeutic agents for atherosclerosis. Interestingly, increasing the population of atheroprotective cells such as Tregs can be considered as a treatment for atherosclerosis. For example, inducing IL-2, a key cytokine for Treg differentiation and proliferation, can increase Tregs population and prevent atherosclerosis development. The function of a number of immune cells, including Th2 cells, Th17 cells, CD8⁺ T cells, and some subtypes of DCs, is still unknown in atherosclerosis because both protective and pro-atherogenic functions have been observed. Therefore, there is a need for further studies to determine the function of such cells in human atherosclerosis and to develop therapies stimulating their protective functions. Another suggested treatment includes stimulating the production of antibodies against LDL or oxLDL-related antigens, which can help inhibit inflammation. Various clinical trials demonstrated the importance of inflammation as a goal for future treatments to decrease the severity of atherosclerosis. Nevertheless, the potential risks of anti-inflammatory treatment, such as infection and cancer cannot be ignored.

ACKNOWLEDGMENTS

The authors would like to thank the Cellular and Molecular Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran. This research receive no specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

CONFLICT OF INTEREST

The authors declared that they have no conflicts of interest.

ORCID

Nader Bagheri <https://orcid.org/0000-0001-5196-5513>

REFERENCES

1. Yang Z, Shi L, Xue Y, Zeng T, Shi Y, Lin Y, et al. Interleukin-32 increases in coronary arteries and plasma from patients with coronary artery disease. *Clinica Chimica Acta*. 2019;497:104-9.
2. Hansson GK, Hermansson A. The immune system in atherosclerosis. *Nature immunology*. 2011;12(3):204-12.

3. Aziz S, Ramsdale D. Chronic total occlusions—a stiff challenge requiring a major breakthrough: is there light at the end of the tunnel? *Heart*. 2005;91(suppl 3):iii42-iii8.
4. Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray CJ. Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *The lancet*. 2006;367(9524):1747-57.
5. Mendis S, Puska P, Norrving B, Organization WH. Global atlas on cardiovascular disease prevention and control: World Health Organization; 2011.
6. Libby P, Lichtman AH, Hansson GK. Immune effector mechanisms implicated in atherosclerosis: from mice to humans. *Immunity*. 2013;38(6):1092-104.
7. Tian K, Xu Y, Sahebkar A, Xu S. CD36 in Atherosclerosis: Pathophysiological Mechanisms and Therapeutic Implications. *Current Atherosclerosis Reports*. 2020;22(10):1-10.
8. Weber C, Zernecke A, Libby P. The multifaceted contributions of leukocyte subsets to atherosclerosis: lessons from mouse models. *Nature reviews immunology*. 2008;8(10):802-15.
9. Gebuhrer V, Murphy J, Bordet J, Reck M, McGregor J. Oxidized low-density lipoprotein induces the expression of P-selectin (GMP140/PADGEM/CD62) on human endothelial cells. *Biochemical Journal*. 1995;306(1):293-8.
10. Davies MJ, Gordon J, Gearing A, Pigott R, Woolf N, Katz D, et al. The expression of the adhesion molecules ICAM-1, VCAM-1, PECAM, and E-selectin in human atherosclerosis. *The Journal of pathology*. 1993;171(3):223-9.
11. Hansson GK, Robertson A-KL, Söderberg-Nauclér C. Inflammation and atherosclerosis. *Annu Rev Pathol Mech Dis*. 2006;1:297-329.
12. Ovchinnikov DA. Macrophages in the embryo and beyond: much more than just giant phagocytes. *genesis*. 2008;46(9):447-62.
13. Wynn TA, Chawla A, Pollard JW. Macrophage biology in development, homeostasis and disease. *Nature*. 2013;496(7446):445-55.
14. Xu S, Li L, Yan J, Ye F, Shao C, Sun Z, et al. CML/CD36 accelerates atherosclerotic progression via inhibiting foam cell migration. *Biomedicine & Pharmacotherapy*. 2018;97:1020-31.
15. Ouimet M, Barrett TJ, Fisher EA. HDL and reverse cholesterol transport: basic mechanisms and their roles in vascular health and disease. *Circulation research*. 2019;124(10):1505-18.
16. Westerterp M, Fotakis P, Ouimet M, Bochem AE, Zhang H, Molusky MM, et al. Cholesterol efflux pathways suppress inflammasome activation, NETosis, and atherogenesis. *Circulation*. 2018;138(9):898-912.
17. Peled M, Nishi H, Weinstock A, Barrett TJ, Zhou F, Quezada A, et al. A wild-type mouse-based model for the regression of inflammation in atherosclerosis. *PLoS One*. 2017;12(3):e0173975.
18. Jaitin DA, Adlung L, Thaïss CA, Weiner A, Li B, Descamps H, et al. Lipid-associated macrophages control metabolic homeostasis in a Trem2-dependent manner. *Cell*. 2019;178(3):686-98. e14.
19. Olzmann JA, Carvalho P. Dynamics and functions of lipid droplets. *Nature reviews Molecular cell biology*. 2019;20(3):137-55.
20. Chávez-Sánchez L, Madrid-Miller A, Chávez-Rueda K, Legorreta-Haquet M, Tesoro-Cruz E, Blanco-Favela F. Activation of TLR2 and TLR4 by minimally modified low-density lipoprotein in human macrophages and monocytes triggers the inflammatory response. *Human immunology*. 2010;71(8):737-44.
21. Koelwyn GJ, Corr EM, Erbay E, Moore KJ. Regulation of macrophage immunometabolism in atherosclerosis. *Nature immunology*. 2018;19(6):526-37.
22. Bories GF, Leitinger N. Macrophage metabolism in atherosclerosis. *Febs Letters*. 2017;591(19):3042-60.
23. Williams JW, Giannarelli C, Rahman A, Randolph GJ, Kovacic JC. Macrophage biology, classification, and phenotype in cardiovascular disease: JACC macrophage in CVD series (part 1). *Journal of the American College of Cardiology*. 2018;72(18):2166-80.

24. Wang F, Zhang S, Vuckovic I, Jeon R, Lerman A, Folmes CD, et al. Glycolytic stimulation is not a requirement for M2 macrophage differentiation. *Cell metabolism*. 2018;28(3):463-75. e4.
25. Devaraj S, Jialal I. C-reactive protein polarizes human macrophages to an M1 phenotype and inhibits transformation to the M2 phenotype. *Arteriosclerosis, thrombosis, and vascular biology*. 2011;31(6):1397-402.
26. Hirose K, Iwabuchi K, Shimada K, Kiyanagi T, Iwahara C, Nakayama H, et al. Different responses to oxidized low-density lipoproteins in human polarized macrophages. *Lipids in health and disease*. 2011;10(1):1.
27. Huang W-C, Sala-Newby GB, Susana A, Johnson JL, Newby AC. Classical macrophage activation up-regulates several matrix metalloproteinases through mitogen activated protein kinases and nuclear factor- κ B. *PloS one*. 2012;7(8):e42507.
28. Van Dyken SJ, Locksley RM. Interleukin-4-and interleukin-13-mediated alternatively activated macrophages: roles in homeostasis and disease. *Annual review of immunology*. 2013;31:317-43.
29. Murray PJ, Wynn TA. Protective and pathogenic functions of macrophage subsets. *Nature reviews immunology*. 2011;11(11):723-37.
30. Mantovani A, Sica A, Locati M. Macrophage polarization comes of age. *Immunity*. 2005;23(4):344-6.
31. Biswas SK, Mantovani A. Macrophage plasticity and interaction with lymphocyte subsets: cancer as a paradigm. *Nature immunology*. 2010;11(10):889-96.
32. Khallou-Laschet J, Varthaman A, Fornasa G, Compain C, Gaston A-T, Clement M, et al. Macrophage plasticity in experimental atherosclerosis. *PloS one*. 2010;5(1):e8852.
33. Feng J, Li L, Ou Z, Li Q, Gong B, Zhao Z, et al. IL-25 stimulates M2 macrophage polarization and thereby promotes mitochondrial respiratory capacity and lipolysis in adipose tissues against obesity. *Cellular & molecular immunology*. 2018;15(5):493-505.
34. Isa SA, Ruffino JS, Ahluwalia M, Thomas AW, Morris K, Webb R. M2 macrophages exhibit higher sensitivity to oxLDL-induced lipotoxicity than other monocyte/macrophage subtypes. *Lipids in health and disease*. 2011;10(1):1-12.
35. Bose D, Banerjee S, Chatterjee N, Das S, Saha M, Saha KD. Inhibition of TGF- β induced lipid droplets switches M2 macrophages to M1 phenotype. *Toxicology in Vitro*. 2019;58:207-14.
36. Palma A, Jarrah AS, Tieri P, Cesareni G, Castiglione F. Gene regulatory network modeling of macrophage differentiation corroborates the continuum hypothesis of polarization states. *Frontiers in physiology*. 2018;9:1659.
37. Kwon E-Y, Kim SY, Choi M-S. Luteolin-enriched artichoke leaf extract alleviates the metabolic syndrome in mice with high-fat diet-induced obesity. *Nutrients*. 2018;10(8):979.
38. Martinez F. siCa, a.; ManToVani, a.; and loCaTi, M. Macrophage activation and polarization. *Frontiers in Bioscience*. 2008;13:453-61.
39. Garlanda C, Bottazzi B, Bastone A, Mantovani A. Pentraxins at the crossroads between innate immunity, inflammation, matrix deposition, and female fertility. *Annu Rev Immunol*. 2005;23:337-66.
40. Bottazzi B, Garlanda C, Salvatori G, Jeannin P, Manfredi A, Mantovani A. Pentraxins as a key component of innate immunity. *Current opinion in immunology*. 2006;18(1):10-5.
41. Savchenko A, Imamura M, Ohashi R, Jiang S, Kawasaki T, Hasegawa G, et al. Expression of pentraxin 3 (PTX3) in human atherosclerotic lesions. *The Journal of Pathology: A Journal of the Pathological Society of Great Britain and Ireland*. 2008;215(1):48-55.
42. Zizzo G, Hilliard BA, Monestier M, Cohen PL. Efficient clearance of early apoptotic cells by human macrophages requires M2c polarization and MerTK induction. *The Journal of Immunology*. 2012;189(7):3508-20.
43. Chinetti-Gbaguidi G, Staels B. Macrophage polarization in metabolic disorders: functions and regulation. *Current opinion in lipidology*. 2011;22(5):365.
44. Fleming BD, Mosser DM. Regulatory macrophages: setting the threshold for therapy. *European journal of immunology*. 2011;41(9):2498-502.

45. Pinhal-Enfield G, Ramanathan M, Hasko G, Vogel SN, Salzman AL, Boons G-J, et al. An angiogenic switch in macrophages involving synergy between Toll-like receptors 2, 4, 7, and 9 and adenosine A2A receptors. *The American journal of pathology*. 2003;163(2):711-21.
46. Ferrante CJ, Pinhal-Enfield G, Elson G, Cronstein BN, Hasko G, Outram S, et al. The adenosine-dependent angiogenic switch of macrophages to an M2-like phenotype is independent of interleukin-4 receptor alpha (IL-4R α) signaling. *Inflammation*. 2013;36(4):921-31.
47. Abu El-Asrar AM, Alam K, Siddiquei MM, Van den Eynde K, Mohammad G, De Hertogh G, et al. Myeloid-related protein-14/MRP-14/S100A9/calgranulin B is associated with inflammation in proliferative diabetic retinopathy. *Ocular immunology and inflammation*. 2018;26(4):615-24.
48. Kraakman MJ, Lee MK, Al-Sharea A, Dragoljevic D, Barrett TJ, Montenont E, et al. Neutrophil-derived S100 calcium-binding proteins A8/A9 promote reticulated thrombocytosis and atherogenesis in diabetes. *The Journal of clinical investigation*. 2017;127(6):2133-47.
49. Flynn MC, Pernes G, Lee MKS, Murphy AJ, Nagareddy PR. Monocytes, macrophages and metabolic disease in atherosclerosis. *Frontiers in pharmacology*. 2019;10:666.
50. Kadl A, Meher AK, Sharma PR, Lee MY, Doran AC, Johnstone SR, et al. Identification of a novel macrophage phenotype that develops in response to atherogenic phospholipids via Nrf2. *Circulation research*. 2010;107(6):737-46.
51. Finn AV, Nakano M, Polavarapu R, Karmali V, Saeed O, Zhao X, et al. Hemoglobin directs macrophage differentiation and prevents foam cell formation in human atherosclerotic plaques. *Journal of the American College of Cardiology*. 2012;59(2):166-77.
52. Nielsen MJ, Møller HJ, Moestrup SK. Hemoglobin and heme scavenger receptors. *Antioxidants & redox signaling*. 2010;12(2):261-73.
53. Philippidis P, Mason J, Evans B, Nadra I, Taylor K, Haskard D, et al. Hemoglobin scavenger receptor CD163 mediates interleukin-10 release and heme oxygenase-1 synthesis: antiinflammatory monocyte-macrophage responses in vitro, in resolving skin blisters in vivo, and after cardiopulmonary bypass surgery. *Circulation research*. 2004;94(1):119-26.
54. Gleissner CA, Shaked I, Little KM, Ley K. CXC chemokine ligand 4 induces a unique transcriptome in monocyte-derived macrophages. *The Journal of Immunology*. 2010;184(9):4810-8.
55. Gleissner CA, Ley K. CXCL4 in atherosclerosis: possible roles in monocyte arrest and macrophage foam cell formation. *THROMBOSIS AND HAEMOSTASIS-STUTTGART*. 2007;98(5):917.
56. Gleissner CA, von Hundelshausen P, Ley K. Platelet chemokines in vascular disease. *Arteriosclerosis, thrombosis, and vascular biology*. 2008;28(11):1920-7.
57. Pitsilos S, Hunt J, Mohler ER, Prabhakar AM, Poncz M, Dawicki J, et al. Platelet factor 4 localization in carotid atherosclerotic plaques: correlation with clinical parameters. *Thrombosis and haemostasis*. 2003;90(12):1112-20.
58. Gleissner CA, Shaked I, Erbel C, Böckler D, Katus HA, Ley K. CXCL4 downregulates the atheroprotective hemoglobin receptor CD163 in human macrophages. *Circulation research*. 2010;106(1):203-11.
59. Nichols BA, Bainton DF, Farquhar MG. Differentiation of monocytes: origin, nature, and fate of their azurophil granules. *The Journal of cell biology*. 1971;50(2):498-515.
60. Ziegler-Heitbrock H. Definition of human blood monocytes. *Journal of Leukocyte Biology*. 2000;67(5):603-6.
61. Kratoofil RM, Kubes P, Deniset JF. Monocyte conversion during inflammation and injury. *Arteriosclerosis, thrombosis, and vascular biology*. 2017;37(1):35-42.
62. Aw NH, Canetti E, Suzuki K, Goh J. Monocyte subsets in atherosclerosis and modification with exercise in humans. *Antioxidants*. 2018;7(12):196.
63. Swirski FK, Libby P, Aikawa E, Alcaide P, Luscinskas FW, Weissleder R, et al. Ly-6C hi monocytes dominate hypercholesterolemia-associated monocytosis and give rise to macrophages in atheromata. *The Journal of clinical investigation*. 2007;117(1):195-205.

64. Auffray C, Fogg D, Garfa M, Elain G, Join-Lambert O, Kayal S, et al. Monitoring of blood vessels and tissues by a population of monocytes with patrolling behavior. *Science*. 2007;317(5838):666-70.
65. Sheel M, Engwerda CR. The diverse roles of monocytes in inflammation caused by protozoan parasitic diseases. *Trends in parasitology*. 2012;28(10):408-16.
66. Nahrendorf M, Swirski FK, Aikawa E, Stangenberg L, Wurdinger T, Figueiredo J-L, et al. The healing myocardium sequentially mobilizes two monocyte subsets with divergent and complementary functions. *The Journal of experimental medicine*. 2007;204(12):3037-47.
67. Olzinski AR, Turner GH, Bernard RE, Karr H, Cornejo CA, Aravindhan K, et al. Pharmacological inhibition of CC chemokine receptor 2 decreases macrophage infiltration in the aortic root of the human CC chemokine receptor 2/apolipoprotein E^{-/-} mouse: magnetic resonance imaging assessment. *Arteriosclerosis, thrombosis, and vascular biology*. 2010;30(2):253-9.
68. Potteaux S, Combadiere C, Pezard A, Riou S, Simon T, Rodero M, et al. Combined inhibition of Ccl2, Cx3cr1 and Ccr5 abrogates Ly6chi and Ly6clo monocytoysis and almost abolishes atherosclerosis in hypercholesterolemic mice. *Am Heart Assoc*; 2007.
69. An G, Wang H, Tang R, Yago T, McDaniel JM, McGee S, et al. PSGL-1 is highly expressed on Ly-6Chi monocytes and a major determinant for Ly-6Chi monocyte recruitment to sites of atherosclerosis in mice. *Circulation*. 2008;117(25):3227.
70. Cybulsky MI, Iiyama K, Li H, Zhu S, Chen M, Iiyama M, et al. A major role for VCAM-1, but not ICAM-1, in early atherosclerosis. *The Journal of clinical investigation*. 2001;107(10):1255-62.
71. Rahman K, Vengrenyuk Y, Ramsey SA, Vila NR, Girgis NM, Liu J, et al. Inflammatory Ly6C hi monocytes and their conversion to M2 macrophages drive atherosclerosis regression. *The Journal of clinical investigation*. 2017;127(8):2904-15.
72. Lin J-D, Nishi H, Poles J, Niu X, Mccauley C, Rahman K, et al. Single-cell analysis of fate-mapped macrophages reveals heterogeneity, including stem-like properties, during atherosclerosis progression and regression. *JCI insight*. 2019;4(4).
73. Passlick B, Flieger D, Ziegler-Heitbrock H. Identification and characterization of a novel monocyte subpopulation in human peripheral blood. 1989.
74. Ziegler-Heitbrock L. The CD14⁺ CD16⁺ blood monocytes: their role in infection and inflammation. *Journal of leukocyte biology*. 2007;81(3):584-92.
75. Allen N, Barrett TJ, Guo Y, Nardi M, Ramkhelawon B, Rockman CB, et al. Circulating monocyte-platelet aggregates are a robust marker of platelet activity in cardiovascular disease. *Atherosclerosis*. 2019;282:11-8.
76. Tomas L, Edsfeldt A, Mollet IG, Perisic Matic L, Prehn C, Adamski J, et al. Altered metabolism distinguishes high-risk from stable carotid atherosclerotic plaques. *European heart journal*. 2018;39(24):2301-10.
77. Zhuang J, Han Y, Xu D, Zhu G, Singh S, Chen L, et al. Comparison of circulating dendritic cell and monocyte subsets at different stages of atherosclerosis: insights from optical coherence tomography. *BMC cardiovascular disorders*. 2017;17(1):1-10.
78. Dann R, Hadi T, Montenont E, Boytard L, Alebrahim D, Feinstein J, et al. Platelet-derived MRP-14 induces monocyte activation in patients with symptomatic peripheral artery disease. *Journal of the American College of Cardiology*. 2018;71(1):53-65.
79. Geissmann F, Jung S, Littman DR. Blood monocytes consist of two principal subsets with distinct migratory properties. *Immunity*. 2003;19(1):71-82.
80. Belge K-U, Dayyani F, Horelt A, Siedlar M, Frankenberger M, Frankenberger B, et al. The proinflammatory CD14⁺ CD16⁺ DR⁺⁺ monocytes are a major source of TNF. *The Journal of Immunology*. 2002;168(7):3536-42.
81. Tsujioka H, Imanishi T, Ikejima H, Kuroi A, Takarada S, Tanimoto T, et al. Impact of heterogeneity of human peripheral blood monocyte subsets on myocardial salvage in patients with primary acute myocardial infarction. *Journal of the American College of Cardiology*. 2009;54(2):130-8.

82. Hamers AA, Dinh HQ, Thomas GD, Marcovecchio P, Blatchley A, Nakao CS, et al. Human monocyte heterogeneity as revealed by high-dimensional mass cytometry. *Arteriosclerosis, thrombosis, and vascular biology*. 2019;39(1):25-36.
83. Villani A-C, Satija R, Reynolds G, Sarkizova S, Shekhar K, Fletcher J, et al. Single-cell RNA-seq reveals new types of human blood dendritic cells, monocytes, and progenitors. *Science*. 2017;356(6335).
84. Swirski FK, Nahrendorf M. Cardioimmunology: the immune system in cardiac homeostasis and disease. *Nature Reviews Immunology*. 2018;18(12):733-44.
85. Soehnlein O, Steffens S, Hidalgo A, Weber C. Neutrophils as protagonists and targets in chronic inflammation. *Nature reviews Immunology*. 2017;17(4):248-61.
86. Casanova-Acebes M, Pitaval C, Weiss LA, Nombela-Arrieta C, Chèvre R, N AG, et al. Rhythmic modulation of the hematopoietic niche through neutrophil clearance. *Cell*. 2013;153(5):1025-35.
87. Hayashi F, Means TK, Luster AD. Toll-like receptors stimulate human neutrophil function. *Blood*. 2003;102(7):2660-9.
88. Kolaczowska E, Kubes P. Neutrophil recruitment and function in health and inflammation. *Nature reviews Immunology*. 2013;13(3):159-75.
89. Drechsler M, Megens RT, van Zandvoort M, Weber C, Soehnlein O. Hyperlipidemia-triggered neutrophilia promotes early atherosclerosis. *Circulation*. 2010;122(18):1837-45.
90. Mantovani A, Cassatella MA, Costantini C, Jaillon S. Neutrophils in the activation and regulation of innate and adaptive immunity. *Nature reviews Immunology*. 2011;11(8):519-31.
91. Chen L, Ge B, Casale FP, Vasquez L, Kwan T, Garrido-Martín D, et al. Genetic Drivers of Epigenetic and Transcriptional Variation in Human Immune Cells. *Cell*. 2016;167(5):1398-414.e24.
92. Mawhin MA, Tilly P, Zirka G, Charles AL, Slimani F, Vonesch JL, et al. Neutrophils recruited by leukotriene B4 induce features of plaque destabilization during endotoxaemia. *Cardiovascular research*. 2018;114(12):1656-66.
93. Roumenina LT, Ruseva MM, Zlatarova A, Ghai R, Kolev M, Olova N, et al. Interaction of C1q with IgG1, C-reactive protein and pentraxin 3: mutational studies using recombinant globular head modules of human C1q A, B, and C chains. *Biochemistry*. 2006;45(13):4093-104.
94. Norata GD, Garlanda C, Catapano AL. The long pentraxin PTX3: a modulator of the immunoinflammatory response in atherosclerosis and cardiovascular diseases. *Trends in cardiovascular medicine*. 2010;20(2):35-40.
95. Van Leeuwen M, Gijbels MJ, Duijvestijn A, Smook M, van de Gaar MJ, Heeringa P, et al. Accumulation of myeloperoxidase-positive neutrophils in atherosclerotic lesions in LDLR^{-/-} mice. *Arteriosclerosis, thrombosis, and vascular biology*. 2008;28(1):84-9.
96. Delporte C, Boudjeltia KZ, Noyon C, Furtmüller PG, Nuyens V, Slomianny M-C, et al. Impact of myeloperoxidase-LDL interactions on enzyme activity and subsequent posttranslational oxidative modifications of apoB-100. *Journal of lipid research*. 2014;55(4):747-57.
97. Nicholls SJ, Hazen SL. Myeloperoxidase and cardiovascular disease. *Arteriosclerosis, thrombosis, and vascular biology*. 2005;25(6):1102-11.
98. Klinke A, Nussbaum C, Kubala L, Friedrichs K, Rudolph TK, Rudolph V, et al. Myeloperoxidase attracts neutrophils by physical forces. *Blood*. 2011;117(4):1350-8.
99. Suzuki S, Takeishi Y, Niizeki T, Koyama Y, Kitahara T, Sasaki T, et al. Pentraxin 3, a new marker for vascular inflammation, predicts adverse clinical outcomes in patients with heart failure. *American heart journal*. 2008;155(1):75-81.
100. Wong ND, Gransar H, Narula J, Shaw L, Moon JH, Miranda-Peats R, et al. Myeloperoxidase, subclinical atherosclerosis, and cardiovascular disease events. *JACC: Cardiovascular Imaging*. 2009;2(9):1093-9.
101. Wågsäter D, Zhu C, Björkegren J, Skogsberg J, Eriksson P. MMP-2 and MMP-9 are prominent matrix metalloproteinases during atherosclerosis development in the Ldlr^{-/-}-Apob100/100 mouse. *International journal of molecular medicine*. 2011;28(2):247-53.
102. Warnatsch A, Ioannou M, Wang Q, Papayannopoulos V. Neutrophil extracellular traps license macrophages for cytokine production in atherosclerosis. *Science*. 2015;349(6245):316-20.

103. Fuchs TA, Brill A, Duerschmied D, Schatzberg D, Monestier M, Myers DD, et al. Extracellular DNA traps promote thrombosis. *Proceedings of the National Academy of Sciences*. 2010;107(36):15880-5.
104. Denny MF, Yalavarthi S, Zhao W, Thacker SG, Anderson M, Sandy AR, et al. A distinct subset of proinflammatory neutrophils isolated from patients with systemic lupus erythematosus induces vascular damage and synthesizes type I IFNs. *The journal of immunology*. 2010;184(6):3284-97.
105. Rasmuson J, Kenne E, Wahlgren M, Soehnlein O, Lindbom L. Heparinoid sevuparin inhibits Streptococcus-induced vascular leak through neutralizing neutrophil-derived proteins. *The FASEB Journal*. 2019;33(9):10443-52.
106. TAEKEMA-ROELVINK ME, VAN KOOTEN C, VAN DER KOOIJ S, HEEMSKERK E, DAHA MR. Proteinase 3 enhances endothelial monocyte chemoattractant protein-1 production and induces increased adhesion of neutrophils to endothelial cells by upregulating intercellular cell adhesion molecule-1. *Journal of the American Society of Nephrology*. 2001;12(5):932-40.
107. Chaly YV, Paleolog E, Kolesnikova T, Tikhonov I, Petratchenko E, Voitenok N. Neutrophil α -defensin human neutrophil peptide modulates cytoline production in human monocytes and adhesion molecule expression in endothelial cells. *European cytokine network*. 2000;11(2):257-66.
108. Kougias P, Chai H, Lin PH, Yao Q, Lumsden AB, Chen C. Neutrophil antimicrobial peptide α -defensin causes endothelial dysfunction in porcine coronary arteries. *Journal of vascular surgery*. 2006;43(2):357-63.
109. Kritikou E, van Duijn J, Nahon JE, van der Heijden T, Bouwman M, Groeneveldt C, et al. Disruption of a CD1d-mediated interaction between mast cells and NKT cells aggravates atherosclerosis. *Atherosclerosis*. 2019;280:132-9.
110. Wilcock A, Bahri R, Bulfone-Paus S, Arkwright PD. Mast cell disorders: from infancy to maturity. *Allergy*. 2019;74(1):53-63.
111. JEZIORSKA M, McCOLLUM C, WOOLLEY DE. Mast cell distribution, activation, and phenotype in atherosclerotic lesions of human carotid arteries. *The Journal of Pathology: A Journal of the Pathological Society of Great Britain and Ireland*. 1997;182(1):115-22.
112. Sun J, Sukhova GK, Wolters PJ, Yang M, Kitamoto S, Libby P, et al. Mast cells promote atherosclerosis by releasing proinflammatory cytokines. *Nature medicine*. 2007;13(6):719-24.
113. Meyer N, Zenclussen AC. Mast cells—Good guys with a bad image? *American Journal of Reproductive Immunology*. 2018;80(4):e13002.
114. Wang J, Cheng X, Xiang M-X, Alanne-Kinnunen M, Wang J-A, Chen H, et al. IgE stimulates human and mouse arterial cell apoptosis and cytokine expression and promotes atherogenesis in Apoe^{-/-} mice. *The Journal of clinical investigation*. 2011;121(9).
115. Wernersson S, Pejler G. Mast cell secretory granules: armed for battle. *Nature Reviews Immunology*. 2014;14(7):478-94.
116. Malbec O, Daëron M. The mast cell IgG receptors and their roles in tissue inflammation. *Immunological reviews*. 2007;217(1):206-21.
117. Ylä-Herttuala S, Palinski W, Butler SW, Picard S, Steinberg D, Witztum JL. Rabbit and human atherosclerotic lesions contain IgG that recognizes epitopes of oxidized LDL. *Arteriosclerosis and Thrombosis: A Journal of Vascular Biology*. 1994;14(1):32-40.
118. Lappalainen J, Lindstedt KA, Oksjoki R, Kovanen PT. OxLDL-IgG immune complexes induce expression and secretion of proatherogenic cytokines by cultured human mast cells. *Atherosclerosis*. 2011;214(2):357-63.
119. den Dekker WK, Tempel D, Bot I, Biessen EA, Joosten LA, Netea MG, et al. Mast cells induce vascular smooth muscle cell apoptosis via a toll-like receptor 4 activation pathway. *Arteriosclerosis, thrombosis, and vascular biology*. 2012;32(8):1960-9.
120. Bahrami A, Parsamanesh N, Atkin SL, Banach M, Sahebkar A. Effect of statins on toll-like receptors: a new insight to pleiotropic effects. *Pharmacological research*. 2018;135:230-8.

121. de Vries MR, Wezel A, Schepers A, van Santbrink PJ, Woodruff TM, Niessen HW, et al. Complement factor C5a as mast cell activator mediates vascular remodelling in vein graft disease. *Cardiovascular research*. 2013;97(2):311-20.
122. Xu J-M, Shi G-P. Emerging role of mast cells and macrophages in cardiovascular and metabolic diseases. *Endocrine reviews*. 2012;33(1):71-108.
123. Kovanen PT. Mast Cells as Potential Accelerators of Human Atherosclerosis—From Early to Late Lesions. *International Journal of Molecular Sciences*. 2019;20(18):4479.
124. Clejan S, Japa S, Clemetson C, Hasabnis SS, David O, Talano J. Blood histamine is associated with coronary artery disease, cardiac events and severity of inflammation and atherosclerosis. *Journal of cellular and molecular medicine*. 2002;6(4):583-92.
125. Lätti S, Leskinen M, Shiota N, Wang Y, Kovanen PT, Lindstedt KA. Mast cell-mediated apoptosis of endothelial cells in vitro: A paracrine mechanism involving TNF- α -mediated down-regulation of bcl-2 expression. *Journal of cellular physiology*. 2003;195(1):130-8.
126. Zhang J, Alcaide P, Liu L, Sun J, He A, Lusinskas FW, et al. Regulation of endothelial cell adhesion molecule expression by mast cells, macrophages, and neutrophils. *PloS one*. 2011;6(1):e14525.
127. Zhi X, Xu C, Zhang H, Tian D, Li X, Ning Y, et al. Tryptase promotes atherosclerotic plaque haemorrhage in ApoE^{-/-} mice. *PLoS One*. 2013;8(4):e60960.
128. Yeong P, Ning Y, Xu Y, Li X, Yin L. Tryptase promotes human monocyte-derived macrophage foam cell formation by suppressing LXR α activation. *Biochimica et Biophysica Acta (BBA)-Molecular and Cell Biology of Lipids*. 2010;1801(5):567-76.
129. Mohajeri M, Kovanen PT, Bianconi V, Pirro M, Cicero AF, Sahebkar A. Mast cell tryptase—Marker and maker of cardiovascular diseases. *Pharmacology & therapeutics*. 2019;199:91-110.
130. Lee M, Lindstedt LK, Kovanen PT. Mast cell-mediated inhibition of reverse cholesterol transport. *Arteriosclerosis and thrombosis: a journal of vascular biology*. 1992;12(11):1329-35.
131. Bot I, Bot M, van Heiningen SH, van Santbrink PJ, Lankhuizen IM, Hartman P, et al. Mast cell chymase inhibition reduces atherosclerotic plaque progression and improves plaque stability in ApoE^{-/-} mice. *Cardiovascular research*. 2011;89(1):244-52.
132. Leskinen MJ, Lindstedt KA, Wang Y, Kovanen PT. Mast cell chymase induces smooth muscle cell apoptosis by a mechanism involving fibronectin degradation and disruption of focal adhesions. *Arteriosclerosis, thrombosis, and vascular biology*. 2003;23(2):238-43.
133. Kokkonen J, Kovanen P. Low-density-lipoprotein binding by mast-cell granules. Demonstration of binding of apolipoprotein B to heparin proteoglycan of exocytosed granules. *Biochemical Journal*. 1987;241(2):583-9.
134. Mäyränpää MI, Heikkilä HM, Lindstedt KA, Walls AF, Kovanen PT. Desquamation of human coronary artery endothelium by human mast cell proteases: implications for plaque erosion. *Coronary artery disease*. 2006;17(7):611-21.
135. Moretta L, Moretta A. Unravelling natural killer cell function: triggering and inhibitory human NK receptors. *The EMBO journal*. 2004;23(2):255-9.
136. Farag SS, Fehniger TA, Ruggeri L, Velardi A, Caligiuri MA. Natural killer cell receptors: new biology and insights into the graft-versus-leukemia effect. *Blood*. 2002;100(6):1935-47.
137. Colucci F, Caligiuri MA, Di Santo JP. What does it take to make a natural killer? *Nature Reviews Immunology*. 2003;3(5):413-25.
138. Bobryshev YV, Lord RS. Identification of natural killer cells in human atherosclerotic plaque. *Atherosclerosis*. 2005;180(2):423-7.
139. Dong K, Ge J-H, Gu S-L, Li S, Zhu W-G, Fan F-Y, et al. Ox-LDL can enhance the interaction of mice natural killer cells and dendritic cells via the CD48-2B4 pathway. *Heart and vessels*. 2011;26(6):637-45.
140. Linton MF, Major AS, Fazio S. Proatherogenic role for NK cells revealed. *Am Heart Assoc*; 2004.

141. Allavena P, Bianchi G, Zhou D, Van Damme J, Jílek P, Sozzani S, et al. Induction of natural killer cell migration by monocyte chemotactic protein- 1, - 2 and - 3. *European journal of immunology*. 1994;24(12):3233-6.
142. Greaves DR, Häkkinen T, Lucas AD, Liddiard K, Jones E, Quinn CM, et al. Linked chromosome 16q13 chemokines, macrophage-derived chemokine, fractalkine, and thymus-and activation-regulated chemokine, are expressed in human atherosclerotic lesions. *Arteriosclerosis, thrombosis, and vascular biology*. 2001;21(6):923-9.
143. Li H, Zhu X, Hu L, Li Q, Ma J, Yan J. Loss of exosomal MALAT1 from ox-LDL-treated vascular endothelial cells induces maturation of dendritic cells in atherosclerosis development. *Cell Cycle*. 2019;18(18):2255-67.
144. Shortman K, Naik SH. Steady-state and inflammatory dendritic-cell development. *Nature Reviews Immunology*. 2007;7(1):19-30.
145. Banchereau J, Steinman RM. Dendritic cells and the control of immunity. *Nature*. 1998;392(6673):245-52.
146. Wang L, Li D, Yang K, Hu Y, Zeng Q. Toll-like receptor-4 and mitogen-activated protein kinase signal system are involved in activation of dendritic cells in patients with acute coronary syndrome. *Immunology*. 2008;125(1):122-30.
147. Alderman CJ, Bunyard PR, Chain BM, Foreman JC, Leake DS, Katz DR. Effects of oxidised low density lipoprotein on dendritic cells: a possible immunoregulatory component of the atherogenic micro-environment? *Cardiovascular research*. 2002;55(4):806-19.
148. Shen L-H, Zhou L, Wang B-Y, Pu J, Hu L-H, Chai D-J, et al. Oxidized low-density lipoprotein induces differentiation of RAW264. 7 murine macrophage cell line into dendritic-like cells. *Atherosclerosis*. 2008;199(2):257-64.
149. von Schlieffen E, Oskolkova OV, Schabbauer G, Gruber F, Blüml S, Genest M, et al. Multi-hit inhibition of circulating and cell-associated components of the toll-like receptor 4 pathway by oxidized phospholipids. *Arteriosclerosis, thrombosis, and vascular biology*. 2009;29(3):356-62.
150. Naiki Y, Sorrentino R, Wong MH, Michelsen KS, Shimada K, Chen S, et al. TLR/MyD88 and liver X receptor α signaling pathways reciprocally control Chlamydia pneumoniae-induced acceleration of atherosclerosis. *The Journal of immunology*. 2008;181(10):7176-85.
151. Deng J, Ma-Krupa W, Gewirtz AT, Younge BR, Goronzy JrJ, Weyand CM. Toll-like receptors 4 and 5 induce distinct types of vasculitis. *Circulation research*. 2009;104(4):488-95.
152. Yilmaz A, Arditi M. Giant cell arteritis: dendritic cells take two T's to tango. *Am Heart Assoc*; 2009.
153. Han JW, Shimada K, Ma-Krupa W, Johnson TL, Nerem RM, Goronzy JrJ, et al. Vessel wall-embedded dendritic cells induce T-cell autoreactivity and initiate vascular inflammation. *Circulation research*. 2008;102(5):546-53.
154. Merad M, Sathe P, Helft J, Miller J, Mortha A. The dendritic cell lineage: ontogeny and function of dendritic cells and their subsets in the steady state and the inflamed setting. *Annual review of immunology*. 2013;31:563-604.
155. Charo IF, Taubman MB. Chemokines in the pathogenesis of vascular disease. *Circulation research*. 2004;95(9):858-66.
156. Weis M, Schlichting CL, Engleman EG, Cooke JP. Endothelial determinants of dendritic cell adhesion and migration: new implications for vascular diseases. *Arteriosclerosis, thrombosis, and vascular biology*. 2002;22(11):1817-23.
157. Liu P, Yu Y-RA, Spencer JA, Johnson AE, Vallanat CT, Fong AM, et al. CX3CR1 deficiency impairs dendritic cell accumulation in arterial intima and reduces atherosclerotic burden. *Arteriosclerosis, thrombosis, and vascular biology*. 2008;28(2):243-50.
158. Paulson KE, Zhu S-N, Chen M, Nurmohamed S, Jongstra-Bilen J, Cybulsky MI. Resident intimal dendritic cells accumulate lipid and contribute to the initiation of atherosclerosis. *Circulation research*. 2010;106(2):383-90.

159. Zaguri R, Verbovetski I, Atallah M, Trahtemberg U, Krispin A, Nahari E, et al. 'Danger' effect of low-density lipoprotein (LDL) and oxidized LDL on human immature dendritic cells. *Clinical & Experimental Immunology*. 2007;149(3):543-52.
160. Nickel T, Schmauss D, Hanssen H, Sicic Z, Krebs B, Jankl S, et al. oxLDL uptake by dendritic cells induces upregulation of scavenger-receptors, maturation and differentiation. *Atherosclerosis*. 2009;205(2):442-50.
161. Nickel T, Pfeiler S, Summo C, Kopp R, Meimarakis G, Sicic Z, et al. oxLDL downregulates the dendritic cell homing factors CCR7 and CCL21. *Mediators of inflammation*. 2012;2012.
162. Niessner A, Sato K, Chaikof EL, Colmegna I, Goronzy JrJ, Weyand CM. CLINICAL PERSPECTIVE. *Circulation*. 2006;114(23):2482-9.
163. Yilmaz A, Weber J, Cicha I, Stumpf C, Klein M, Raithel D, et al. Decrease in circulating myeloid dendritic cell precursors in coronary artery disease. *Journal of the American College of Cardiology*. 2006;48(1):70-80.
164. Van Brussel I, Van Vré EA, De Meyer GR, Vrints CJ, Bosmans JM, Bult H. Decreased numbers of peripheral blood dendritic cells in patients with coronary artery disease are associated with diminished plasma Flt3 ligand levels and impaired plasmacytoid dendritic cell function. *Clinical science (London, England : 1979)*. 2011;120(9):415-26.
165. Liu J, Cao X. Regulatory dendritic cells in autoimmunity: a comprehensive review. *Journal of autoimmunity*. 2015;63:1-12.
166. Morelli AE, Thomson AW. Tolerogenic dendritic cells and the quest for transplant tolerance. *Nature Reviews Immunology*. 2007;7(8):610-21.
167. Selenko-Gebauer N, Majdic O, Szekeres A, Höfler G, Guthann E, Korthäuer U, et al. B7-H1 (programmed death-1 ligand) on dendritic cells is involved in the induction and maintenance of T cell anergy. *The Journal of Immunology*. 2003;170(7):3637-44.
168. Darrasse-Jèze G, Deroubaix S, Mouquet H, Victora GD, Eisenreich T, Yao K-h, et al. Feedback control of regulatory T cell homeostasis by dendritic cells in vivo. *Journal of Experimental Medicine*. 2009;206(9):1853-62.
169. Hermansson A, Johansson DK, Ketelhuth DF, Andersson J, Zhou X, Hansson GK. Immunotherapy with tolerogenic apolipoprotein B-100-loaded dendritic cells attenuates atherosclerosis in hypercholesterolemic mice. *Circulation*. 2011;123(10):1083-91.
170. Lan YY, Wang Z, Raimondi G, Wu W, Colvin BL, De Creus A, et al. "Alternatively activated" dendritic cells preferentially secrete IL-10, expand Foxp3⁺ CD4⁺ T cells, and induce long-term organ allograft survival in combination with CTLA4-Ig. *The Journal of Immunology*. 2006;177(9):5868-77.
171. van Duijn J, Kritikou E, Benne N, van der Heijden T, van Puijvelde GH, Kröner MJ, et al. CD8⁺ T-cells contribute to lesion stabilization in advanced atherosclerosis by limiting macrophage content and CD4⁺ T-cell responses. *Cardiovascular research*. 2019;115(4):729-38.
172. Bobryshev YV. Dendritic cells in atherosclerosis: current status of the problem and clinical relevance. *European heart journal*. 2005;26(17):1700-4.
173. Stemme S, Faber B, Holm J, Wiklund O, Witztum JL, Hansson GK. T lymphocytes from human atherosclerotic plaques recognize oxidized low density lipoprotein. *Proceedings of the National Academy of Sciences*. 1995;92(9):3893-7.
174. Palinski W, Tangirala RK, Miller E, Young SG, Witztum JL. Increased autoantibody titers against epitopes of oxidized LDL in LDL receptor-deficient mice with increased atherosclerosis. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 1995;15(10):1569-76.
175. Salonen JT, Korpela H, Salonen R, Nyyssonen K, Yla-Herttuala S, Yamamoto R, et al. Autoantibody against oxidised LDL and progression of carotid atherosclerosis. *The Lancet*. 1992;339(8798):883-7.
176. Zhou X, Robertson A-KL, Hjerpe C, Hansson GK. Adoptive transfer of CD4⁺ T cells reactive to modified low-density lipoprotein aggravates atherosclerosis. *Arteriosclerosis, thrombosis, and vascular biology*. 2006;26(4):864-70.

177. George J, Harats D, Gilburd B, Afek A, Levy Y, Schneiderman J, et al. Immunolocalization of β 2-glycoprotein I (apolipoprotein H) to human atherosclerotic plaques: potential implications for lesion progression. *Circulation*. 1999;99(17):2227-30.
178. George J, Afek A, Gilburd B, Blank M, Levy Y, Aron-Maor A, et al. Induction of early atherosclerosis in LDL-receptor-deficient mice immunized with β 2-glycoprotein I. *Circulation*. 1998;98(11):1108-15.
179. Hansson GK, Holm J, Jonasson L. Detection of activated T lymphocytes in the human atherosclerotic plaque. *The American journal of pathology*. 1989;135(1):169.
180. Whitman SC, Ravisankar P, Daugherty A. IFN- γ deficiency exerts gender-specific effects on atherogenesis in apolipoprotein E-/-mice. *Journal of interferon & cytokine research*. 2002;22(6):661-70.
181. Gupta S, Pablo AM, c Jiang X, Wang N, Tall AR, Schindler C. IFN-gamma potentiates atherosclerosis in ApoE knock-out mice. *The Journal of clinical investigation*. 1997;99(11):2752-61.
182. Buono C, Come CE, Stavrakis G, Maguire GF, Connelly PW, Lichtman AH. Influence of interferon- γ on the extent and phenotype of diet-induced atherosclerosis in the LDLR-deficient mouse. *Arteriosclerosis, thrombosis, and vascular biology*. 2003;23(3):454-60.
183. Whitman SC, Ravisankar P, Elam H, Daugherty A. Exogenous interferon- γ enhances atherosclerosis in apolipoprotein E-/- mice. *The American journal of pathology*. 2000;157(6):1819-24.
184. Niwa T, Wada H, Ohashi H, Iwamoto N, Ohta H, Kirii H, et al. Interferon- γ produced by bone marrow-derived cells attenuates atherosclerotic lesion formation in LDLR-deficient mice. *Journal of atherosclerosis and thrombosis*. 2004;11(2):79-87.
185. Gerdes N, Sukhova GK, Libby P, Reynolds RS, Young JL, Schönbeck U. Expression of interleukin (IL)-18 and functional IL-18 receptor on human vascular endothelial cells, smooth muscle cells, and macrophages: implications for atherogenesis. *The Journal of experimental medicine*. 2002;195(2):245-57.
186. Leon MA, Zuckerman S. Gamma interferon: a central mediator in atherosclerosis. *Inflammation research*. 2005;54(10):395-411.
187. Harvey EJ, Ramji DP. Interferon- γ and atherosclerosis: pro-or anti-atherogenic? *Cardiovascular research*. 2005;67(1):11-20.
188. Hansson GK, Hellstrand M, Rymo L, Rubbia L, Gabbiani G. Interferon gamma inhibits both proliferation and expression of differentiation-specific alpha-smooth muscle actin in arterial smooth muscle cells. *The Journal of experimental medicine*. 1989;170(5):1595-608.
189. Amento EP, Ehsani N, Palmer H, Libby P. Cytokines and growth factors positively and negatively regulate interstitial collagen gene expression in human vascular smooth muscle cells. *Arteriosclerosis and thrombosis: a journal of vascular biology*. 1991;11(5):1223-30.
190. McLaren JE, Ramji DP. Interferon gamma: a master regulator of atherosclerosis. *Cytokine & growth factor reviews*. 2009;20(2):125-35.
191. Buono C, Binder CJ, Stavrakis G, Witztum JL, Glimcher LH, Lichtman AH. T-bet deficiency reduces atherosclerosis and alters plaque antigen-specific immune responses. *Proceedings of the National Academy of Sciences*. 2005;102(5):1596-601.
192. Sun J, Hartvigsen K, Chou M-Y, Zhang Y, Sukhova GK, Zhang J, et al. Deficiency of antigen presenting cell invariant chain reduces atherosclerosis in mice. *Circulation*. 2010;122(8):808.
193. Lutgens E, Lievens D, Beckers L, Donners M, Daemen M. CD40 and its ligand in atherosclerosis. *Trends in cardiovascular medicine*. 2007;17(4):118-23.
194. Mach F, Schönbeck U, Sukhova GK, Atkinson E, Libby P. Reduction of atherosclerosis in mice by inhibition of CD40 signalling. *Nature*. 1998;394(6689):200-3.
195. Lutgens E, Cleutjens KB, Heeneman S, Kotliansky VE, Burkly LC, Daemen MJ. Both early and delayed anti-CD40L antibody treatment induces a stable plaque phenotype. *Proceedings of the National Academy of Sciences*. 2000;97(13):7464-9.
196. Lutgens E, Gorelik L, Daemen MJ, de Muinck ED, Grewal IS, Kotliansky VE, et al. Requirement for CD154 in the progression of atherosclerosis. *Nature medicine*. 1999;5(11):1313-6.

197. Elhage R, Jawien J, Rudling M, Ljunggren H-G, Takeda K, Akira S, et al. Reduced atherosclerosis in interleukin-18 deficient apolipoprotein E-knockout mice. *Cardiovascular research*. 2003;59(1):234-40.
198. Huber SA, Sakkinen P, David C, Newell M, Tracy R. T helper–cell phenotype regulates atherosclerosis in mice under conditions of mild hypercholesterolemia. *Circulation*. 2001;103(21):2610-6.
199. Davenport P, Tipping PG. The role of interleukin-4 and interleukin-12 in the progression of atherosclerosis in apolipoprotein E-deficient mice. *The American journal of pathology*. 2003;163(3):1117-25.
200. King VL, Szilvassy SJ, Daugherty A. Interleukin-4 deficiency decreases atherosclerotic lesion formation in a site-specific manner in female LDL receptor–/– mice. *Arteriosclerosis, thrombosis, and vascular biology*. 2002;22(3):456-61.
201. Leskinen MJ, Kovanen PT, Lindstedt KA. Regulation of smooth muscle cell growth, function and death in vitro by activated mast cells—a potential mechanism for the weakening and rupture of atherosclerotic plaques. *Biochemical pharmacology*. 2003;66(8):1493-8.
202. Shimizu K, Shichiri M, Libby P, Lee RT, Mitchell RN. Th2-predominant inflammation and blockade of IFN- γ signaling induce aneurysms in allografted aortas. *The Journal of clinical investigation*. 2004;114(2):300-8.
203. Binder CJ, Hartvigsen K, Chang M-K, Miller M, Broide D, Palinski W, et al. IL-5 links adaptive and natural immunity specific for epitopes of oxidized LDL and protects from atherosclerosis. *The Journal of clinical investigation*. 2004;114(3):427-37.
204. Hörkkö S, Bird DA, Miller E, Itabe H, Leitinger N, Subbanagounder G, et al. Monoclonal autoantibodies specific for oxidized phospholipids or oxidized phospholipid–protein adducts inhibit macrophage uptake of oxidized low-density lipoproteins. *The Journal of clinical investigation*. 1999;103(1):117-28.
205. Schmitz J, Owyang A, Oldham E, Song Y, Murphy E, McClanahan TK, et al. IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines. *Immunity*. 2005;23(5):479-90.
206. Miller AM, Xu D, Asquith DL, Denby L, Li Y, Sattar N, et al. IL-33 reduces the development of atherosclerosis. *Journal of Experimental Medicine*. 2012;209(13):2515.
207. Park H, Li Z, Yang XO, Chang SH, Nurieva R, Wang Y-H, et al. A distinct lineage of CD4 T cells regulates tissue inflammation by producing interleukin 17. *Nature immunology*. 2005;6(11):1133-41.
208. Harrington LE, Hatton RD, Mangan PR, Turner H, Murphy TL, Murphy KM, et al. Interleukin 17–producing CD4+ effector T cells develop via a lineage distinct from the T helper type 1 and 2 lineages. *Nature immunology*. 2005;6(11):1123-32.
209. McGeachy MJ, Bak-Jensen KS, Chen Y, Tato CM, Blumenschein W, McClanahan T, et al. TGF- β and IL-6 drive the production of IL-17 and IL-10 by T cells and restrain TH-17 cell–mediated pathology. *Nature immunology*. 2007;8(12):1390-7.
210. Yoshida H, Yoshiyuki M. Regulation of immune responses by interleukin-27. *Immunological reviews*. 2008;226(1):234-47.
211. Van Es T, Van Puijvelde G, Ramos O, Segers F, Joosten L, van Den Berg W, et al. Attenuated atherosclerosis upon IL-17R signaling disruption in LDLr deficient mice. *Biochemical and biophysical research communications*. 2009;388(2):261-5.
212. Erbel C, Dengler TJ, Wangler S, Lasitschka F, Bea F, Wambsganss N, et al. Expression of IL-17A in human atherosclerotic lesions is associated with increased inflammation and plaque vulnerability. *Basic research in cardiology*. 2011;106(1):125-34.
213. Zhou L, Chong MM, Littman DR. Plasticity of CD4+ T cell lineage differentiation. *Immunity*. 2009;30(5):646-55.
214. Quintana FJ, Basso AS, Iglesias AH, Korn T, Farez MF, Bettelli E, et al. Control of T reg and TH 17 cell differentiation by the aryl hydrocarbon receptor. *Nature*. 2008;453(7191):65-71.

215. Zhang F, Meng G, Strober W. Interactions among the transcription factors Runx1, ROR γ t and Foxp3 regulate the differentiation of interleukin 17–producing T cells. *Nature immunology*. 2008;9(11):1297-306.
216. Kaplan MH. Th9 cells: differentiation and disease. *Immunological reviews*. 2013;252(1):104-15.
217. Gregersen I, Skjelland M, Holm S, Holven KB, Krogh-Sørensen K, Russell D, et al. Increased systemic and local interleukin 9 levels in patients with carotid and coronary atherosclerosis. *PLoS One*. 2013;8(8):e72769.
218. Zhou Q, Fu Y, Hu L, Li Q, Jin M, Jiang E. Relationship of circulating chemerin and omentin levels with Th17 and Th9 cell immune responses in patients with asthma. *Journal of Asthma*. 2018;55(6):579-87.
219. Shohan M, Dehghani R, Khodadadi A, Dehnavi S, Ahmadi R, Joudaki N, et al. Interleukin-22 and intestinal homeostasis: Protective or destructive? *IUBMB Life*. 2020;72(8):1585-602.
220. Azizi G, Yazdani R, Mirshafiey A. Th22 cells in autoimmunity: a review of current knowledge. *European annals of allergy and clinical immunology*. 2015;47(4):108-17.
221. Zhang L, Wang T, Wang X-q, Du R-z, Zhang K-n, Liu X-g, et al. Elevated frequencies of circulating Th22 cell in addition to Th17 cell and Th17/Th1 cell in patients with acute coronary syndrome. *PloS one*. 2013;8(12):e71466.
222. Xia Q, Xiang X, Patel S, Puranik R, Xie Q, Bao S. Characterisation of IL-22 and interferon-gamma-inducible chemokines in human carotid plaque. *International journal of cardiology*. 2012;154(2):187-9.
223. Fatkhullina AR, Peshkova IO, Dzutsev A, Aghayev T, McCulloch JA, Thovarai V, et al. An interleukin-23-interleukin-22 axis regulates intestinal microbial homeostasis to protect from diet-induced atherosclerosis. *Immunity*. 2018;49(5):943-57. e9.
224. Crotty S. T follicular helper cell biology: a decade of discovery and diseases. *Immunity*. 2019;50(5):1132-48.
225. Ryu H, Lim H, Choi G, Park Y-J, Cho M, Na H, et al. Atherogenic dyslipidemia promotes autoimmune follicular helper T cell responses via IL-27. *Nature Immunology*. 2018;19(6):583-93.
226. Nus M, Sage AP, Lu Y, Masters L, Lam BY, Newland S, et al. Marginal zone B cells control the response of follicular helper T cells to a high-cholesterol diet. *Nature medicine*. 2017;23(5):601-10.
227. Clement M, Guedj K, Andreatta F, Morvan M, Bey L, Khallou-Laschet J, et al. Control of the T follicular helper–germinal center B-cell axis by CD8+ regulatory T cells limits atherosclerosis and tertiary lymphoid organ development. *Circulation*. 2015;131(6):560-70.
228. Gaddis DE, Padgett LE, Wu R, McSkimming C, Romines V, Taylor AM, et al. Apolipoprotein AI prevents regulatory to follicular helper T cell switching during atherosclerosis. *Nature communications*. 2018;9(1):1-15.
229. Weng N-p, Akbar AN, Goronzy J. CD28– T cells: their role in the age-associated decline of immune function. *Trends in immunology*. 2009;30(7):306-12.
230. Dumitriu IE, Baruah P, Finlayson CJ, Loftus IM, Antunes RF, Lim P, et al. High levels of costimulatory receptors OX40 and 4-1BB characterize CD4+ CD28null T cells in patients with acute coronary syndrome. *Circulation research*. 2012;110(6):857-69.
231. Kovalcsik E, Antunes RF, Baruah P, Kaski JC, Dumitriu IE. Proteasome-mediated reduction in proapoptotic molecule Bim renders CD4+ CD28null T cells resistant to apoptosis in acute coronary syndrome. *Circulation*. 2015;131(8):709-20.
232. Yadav AK, Kumar V, Jha V. Heat shock proteins 60 and 70 specific proinflammatory and cytotoxic response of CD4+ CD28null cells in chronic kidney disease. *Mediators of inflammation*. 2013;2013.
233. Roy P, Ali AJ, Kobiyama K, Ghosheh Y, Ley K. Opportunities for an atherosclerosis vaccine: from mice to humans. *Vaccine*. 2020.
234. Okba AM, Raafat MAER, Farres MN, Melek NAEN, Amin MM, Gendy NN. Expanded peripheral CD4+ CD28null T cells and its association with atherosclerotic changes in patients with end stage renal disease on hemodialysis. *Human immunology*. 2019;80(9):748-54.

235. Shevach EM, DiPaolo RA, Andersson J, Zhao DM, Stephens GL, Thornton AM. The lifestyle of naturally occurring CD4⁺ CD25⁺ Foxp3⁺ regulatory T cells. *Immunological reviews*. 2006;212(1):60-73.
236. Stephens GL, Shevach EM. Foxp3⁺ regulatory T cells: selfishness under scrutiny. *Immunity*. 2007;27(3):417-9.
237. Wakkach A, Fournier N, Brun V, Breittmayer J-P, Cottrez F, Groux H. Characterization of dendritic cells that induce tolerance and T regulatory 1 cell differentiation in vivo. *Immunity*. 2003;18(5):605-17.
238. Mor A, Planer D, Luboshits G, Afek A, Metzger S, Chajek-Shaul T, et al. Role of naturally occurring CD4⁺ CD25⁺ regulatory T cells in experimental atherosclerosis. *Arteriosclerosis, thrombosis, and vascular biology*. 2007;27(4):893-900.
239. Ait-Oufella H, Salomon BL, Potteaux S, Robertson A-KL, Gourdy P, Zoll J, et al. Natural regulatory T cells control the development of atherosclerosis in mice. *Nature medicine*. 2006;12(2):178-80.
240. Mallat Z, Gojova A, Brun V, Esposito B, Fournier N, Cottrez F, et al. Induction of a regulatory T cell type 1 response reduces the development of atherosclerosis in apolipoprotein E-knockout mice. *Circulation*. 2003;108(10):1232-7.
241. Zhao C, Davies JD. A peripheral CD4⁺ T cell precursor for naive, memory, and regulatory T cells. *Journal of Experimental Medicine*. 2010;207(13):2883-94.
242. Mallat Z, Besnard S, Duriez M, Deleuze V, Emmanuel F, Bureau MF, et al. Protective role of interleukin-10 in atherosclerosis. *Circulation research*. 1999;85(8):e17-e24.
243. Mallat Z, Gojova A, Marchiol-Fournigault C, Esposito B, Kamaté C, Merval R, et al. Inhibition of transforming growth factor- β signaling accelerates atherosclerosis and induces an unstable plaque phenotype in mice. *Circulation research*. 2001;89(10):930-4.
244. Pinderski Oslund LJ, Hedrick CC, Olvera T, Hagenbaugh A, Territo M, Berliner JA, et al. Interleukin-10 blocks atherosclerotic events in vitro and in vivo. *Arteriosclerosis, thrombosis, and vascular biology*. 1999;19(12):2847-53.
245. Veillard NR, Steffens S, Burger F, Pelli G, Mach F. Differential expression patterns of proinflammatory and antiinflammatory mediators during atherogenesis in mice. *Arteriosclerosis, thrombosis, and vascular biology*. 2004;24(12):2339-44.
246. Heller EA, Liu E, Tager AM, Yuan Q, Lin AY, Ahluwalia N, et al. Chemokine CXCL10 promotes atherogenesis by modulating the local balance of effector and regulatory T cells. *Circulation*. 2006;113(19):2301-12.
247. Veillard NR, Steffens S, Pelli G, Lu B, Kwak BR, Gerard C, et al. Differential influence of chemokine receptors CCR2 and CXCR3 in development of atherosclerosis in vivo. *Circulation*. 2005;112(6):870-8.
248. Wigren M, Bengtsson D, Dunér P, Olofsson K, Björkbacka H, Bengtsson E, et al. Atheroprotective effects of Alum are associated with capture of oxidized LDL antigens and activation of regulatory T cells. *Circulation Research*. 2009;104(12):e62-e70.
249. Lin J, Li M, Wang Z, He S, Ma X, Li D. The role of CD4⁺ CD25⁺ regulatory T cells in macrophage-derived foam-cell formation. *Journal of lipid research*. 2010;51(5):1208-17.
250. Liu G, Ma H, Qiu L, Li L, Cao Y, Ma J, et al. Phenotypic and functional switch of macrophages induced by regulatory CD4⁺ CD25⁺ T cells in mice. *Immunology and cell biology*. 2011;89(1):130-42.
251. Hwang Y, Yu HT, Kim D-H, Jang J, Kim HY, Kang I, et al. Expansion of CD8⁺ T cells lacking the IL-6 receptor α chain in patients with coronary artery diseases (CAD). *Atherosclerosis*. 2016;249:44-51.
252. Gewaltig J, Kummer M, Koella C, Cathomas G, Biedermann BC. Requirements for CD8 T-cell migration into the human arterial wall. *Human pathology*. 2008;39(12):1756-62.
253. Paul VSV, Paul CMP, Kuruvilla S. Quantification of various inflammatory cells in advanced atherosclerotic plaques. *Journal of Clinical and Diagnostic Research: JCDR*. 2016;10(5):EC35.

254. Dimayuga PC, Zhao X, Yano J, Lio WM, Zhou J, Mihailovic PM, et al. Identification of apoB-100 Peptide-Specific CD 8+ T Cells in Atherosclerosis. *Journal of the American Heart Association*. 2017;6(7):e005318.
255. Cochain C, Koch M, Chaudhari SM, Busch M, Pelisek J, Boon L, et al. CD8+ T cells regulate monopoiesis and circulating Ly6Chigh monocyte levels in atherosclerosis in mice. *Circulation research*. 2015;117(3):244-53.
256. Seijkens TT, Poels K, Meiler S, Van Tiel CM, Kusters PJ, Reiche M, et al. Deficiency of the T cell regulator Casitas B-cell lymphoma-B aggravates atherosclerosis by inducing CD8+ T cell-mediated macrophage death. *European heart journal*. 2019;40(4):372-82.
257. Kyaw T, Winship A, Tay C, Kanellakis P, Hosseini H, Cao A, et al. Cytotoxic and proinflammatory CD8+ T lymphocytes promote development of vulnerable atherosclerotic plaques in apoE-deficient mice. *Circulation*. 2013;127(9):1028-39.
258. Chyu K-Y, Zhao X, Dimayuga PC, Zhou J, Li X, Yano J, et al. CD8+ T cells mediate the athero-protective effect of immunization with an ApoB-100 peptide. *PLoS One*. 2012;7(2):e30780.
259. Bagchi-Chakraborty J, Francis A, Bray T, Masters L, Tsiantoulas D, Nus M, et al. B cell Fcγ receptor IIb modulates atherosclerosis in male and female mice by controlling adaptive germinal center and innate B-1-cell responses. *Arteriosclerosis, thrombosis, and vascular biology*. 2019;39(7):1379-89.
260. Aubry M-C, Riehle DL, Edwards WD, Maradit-Kremers H, Roger VL, Sebo TJ, et al. B-Lymphocytes in plaque and adventitia of coronary arteries in two patients with rheumatoid arthritis and coronary atherosclerosis: preliminary observations. *Cardiovascular pathology*. 2004;13(4):233-6.
261. Zhou X, Hansson G. Detection of B cells and proinflammatory cytokines in atherosclerotic plaques of hypercholesterolaemic apolipoprotein E knockout mice. *Scandinavian journal of immunology*. 1999;50(1):25-30.
262. Watanabe M, Sangawa A, Sasaki Y, Yamashita M, Tanaka-Shintani M, Shintaku M, et al. Distribution of inflammatory cells in adventitia changed with advancing atherosclerosis of human coronary artery. *Journal of atherosclerosis and thrombosis*. 2007;14(6):325-31.
263. Gräbner R, Lötzer K, Döpping S, Hildner M, Radke D, Beer M, et al. Lymphotoxin β receptor signaling promotes tertiary lymphoid organogenesis in the aorta adventitia of aged ApoE^{-/-} mice. *Journal of Experimental Medicine*. 2009;206(1):233-48.
264. Van Leeuwen M, Damoiseaux J, Duijvestijn A, Tervaert JC. The therapeutic potential of targeting B cells and anti-oxLDL antibodies in atherosclerosis. *Autoimmunity reviews*. 2009;9(1):53-7.
265. Tsimikas S. Oxidized low-density lipoprotein biomarkers in atherosclerosis. *Current atherosclerosis reports*. 2006;8(1):55-61.
266. Witztum JL, Binder CJ, Chou M-Y, Fogelstrand L, Hartvigsen K, Shaw PX, et al. Natural antibodies in murine atherosclerosis. *Current drug targets*. 2008;9(3):190-5.
267. Schiopu A, Freundéus B, Jansson B, Söderberg I, Ljungcrantz I, Araya Z, et al. Recombinant antibodies to an oxidized low-density lipoprotein epitope induce rapid regression of atherosclerosis in Apobec-1^{-/-}/low-density lipoprotein receptor^{-/-} mice. *Journal of the American College of Cardiology*. 2007;50(24):2313-8.
268. Major AS, Fazio S, Linton MF. B-lymphocyte deficiency increases atherosclerosis in ldl receptor-null mice. *Arteriosclerosis, thrombosis, and vascular biology*. 2002;22(11):1892-8.
269. Sage AP, Nus M, Baker LL, Finigan AJ, Masters LM, Mallat Z. Regulatory B cell-specific interleukin-10 is dispensable for atherosclerosis development in mice. *Arteriosclerosis, thrombosis, and vascular biology*. 2015;35(8):1770-3.
270. von der Thüsen JH, van Berkel TJ, Biessen EA. Induction of rapid atherogenesis by perivascular carotid collar placement in apolipoprotein E-deficient and low-density lipoprotein receptor-deficient mice. *Circulation*. 2001;103(8):1164-70.
271. Mauri C, Bosma A. Immune regulatory function of B cells. *Annual review of immunology*. 2012;30:221-41.
272. Getz GS, Reardon CA. Natural killer T cells in atherosclerosis. *Nature Reviews Cardiology*. 2017;14(5):304-14.

273. Bennstein SB. Unraveling natural killer T-cells development. *Frontiers in Immunology*. 2018;8:1950.
274. Bondarenko S, Catapano AL, Norata GD. The CD1d-natural killer T cell axis in atherosclerosis. *Journal of innate immunity*. 2014;6(1):3-12.
275. Li Y, To K, Kanellakis P, Hosseini H, Deswaerte V, Tipping P, et al. CD4+ natural killer T cells potentially augment aortic root atherosclerosis by perforin-and granzyme B-dependent cytotoxicity. *Circulation research*. 2015;116(2):245-54.
276. Nielsen MM, Witherden DA, Havran WL. $\gamma\delta$ T cells in homeostasis and host defence of epithelial barrier tissues. *Nature Reviews Immunology*. 2017;17(12):733-45.
277. Cheng HY, Wu R, Gebre AK, Hanna RN, Smith DJ, Parks JS, et al. Increased cholesterol content in gammadelta ($\gamma\delta$) T lymphocytes differentially regulates their activation. *PLoS One*. 2013;8(5):e63746.
278. Cheng HY, Wu R, Hedrick CC. Gammadelta ($\gamma\delta$) T lymphocytes do not impact the development of early atherosclerosis. *Atherosclerosis*. 2014;234(2):265-9.
279. Vu DM, Tai A, Tatro JB, Karas RH, Huber BT, Beasley D. $\gamma\delta$ T cells are prevalent in the proximal aorta and drive nascent atherosclerotic lesion progression and neutrophilia in hypercholesterolemic mice. *PLoS One*. 2014;9(10):e109416.
280. Kleindienst R, Xu Q, Willeit J, Waldenberger FR, Weimann S, Wick G. Immunology of atherosclerosis. Demonstration of heat shock protein 60 expression and T lymphocytes bearing alpha/beta or gamma/delta receptor in human atherosclerotic lesions. *Am J Pathol*. 1993;142(6):1927-37.
281. Stark MA, Huo Y, Burcin TL, Morris MA, Olson TS, Ley K. Phagocytosis of apoptotic neutrophils regulates granulopoiesis via IL-23 and IL-17. *Immunity*. 2005;22(3):285-94.
282. Drechsler M, Megens RT, van Zandvoort M, Weber C, Soehnlein O. Hyperlipidemia-triggered neutrophilia promotes early atherosclerosis. *Circulation*. 2010;122(18):1837-45.
283. Chen L, Ge B, Casale FP, Vasquez L, Kwan T, Garrido-Martín D, et al. Genetic drivers of epigenetic and transcriptional variation in human immune cells. *Cell*. 2016;167(5):1398-414. e24.
284. Stark MA, Huo Y, Burcin TL, Morris MA, Olson TS, Ley K. Phagocytosis of apoptotic neutrophils regulates granulopoiesis via IL-23 and IL-17. *Immunity*. 2005;22(3):285-94.
285. Tardif J-C, McMurray JJ, Klug E, Small R, Schumi J, Choi J, et al. Effects of succinobucol (AGI-1067) after an acute coronary syndrome: a randomised, double-blind, placebo-controlled trial. *The Lancet*. 2008;371(9626):1761-8.
286. Lehrer-Graiwer J, Singh P, Abdelbaky A, Vucic E, Korsgren M, Baruch A, et al. FDG-PET imaging for oxidized LDL in stable atherosclerotic disease: a phase II study of safety, tolerability, and anti-inflammatory activity. *JACC: Cardiovascular Imaging*. 2015;8(4):493-4.
287. Tardif J-C, Tanguay J-F, Wright SR, Duchatelle V, Petroni T, Grégoire JC, et al. Effects of the P-selectin antagonist inclacumab on myocardial damage after percutaneous coronary intervention for non-ST-segment elevation myocardial infarction: results of the SELECT-ACS trial. *Journal of the American College of Cardiology*. 2013;61(20):2048-55.
288. Stähli BE, Gebhard C, Duchatelle V, Cournoyer D, Petroni T, Tanguay JF, et al. Effects of the P-selectin antagonist inclacumab on myocardial damage after percutaneous Coronary intervention according to timing of infusion: insights from the SELECT-ACS trial. *Journal of the American Heart Association*. 2016;5(11):e004255.
289. Nidorf SM, Eikelboom JW, Budgeon CA, Thompson PL. Low-dose colchicine for secondary prevention of cardiovascular disease. *Journal of the American College of Cardiology*. 2013;61(4):404-10.
290. Abbate A, Kontos MC, Abouzaki NA, Melchior RD, Thomas C, Van Tassell BW, et al. Comparative safety of interleukin-1 blockade with anakinra in patients with ST-segment elevation acute myocardial infarction (from the VCU-ART and VCU-ART2 pilot studies). *The American journal of cardiology*. 2015;115(3):288-92.
291. Ridker PM, Everett BM, Thuren T, MacFadyen JG, Chang WH, Ballantyne C, et al. Antiinflammatory therapy with canakinumab for atherosclerotic disease. *New England journal of medicine*. 2017;377(12):1119-31.

292. Nicholls SJ, Kastelein JJ, Schwartz GG, Bash D, Rosenson RS, Cavender MA, et al. Varespladib and cardiovascular events in patients with an acute coronary syndrome: the VISTA-16 randomized clinical trial. *Jama*. 2014;311(3):252-62.
293. Investigators S. Darapladib for preventing ischemic events in stable coronary heart disease. *New England Journal of Medicine*. 2014;370(18):1702-11.
294. O'Donoghue ML, Braunwald E, White HD, Steen DL, Lukas MA, Tarka E, et al. Effect of darapladib on major coronary events after an acute coronary syndrome: the SOLID-TIMI 52 randomized clinical trial. *Jama*. 2014;312(10):1006-15.
295. Tardif J-C, L'Allier PL, Ibrahim R, Grégoire JC, Nozza A, Cossette M, et al. Treatment with 5-lipoxygenase inhibitor VIA-2291 (Atreleuton) in patients with recent acute coronary syndrome. *Circulation: Cardiovascular Imaging*. 2010;3(3):298-307.
296. Gaztanaga J, Farkouh M, Rudd JH, Brotz TM, Rosenbaum D, Mani V, et al. A phase 2 randomized, double-blind, placebo-controlled study of the effect of VIA-2291, a 5-lipoxygenase inhibitor, on vascular inflammation in patients after an acute coronary syndrome. *Atherosclerosis*. 2015;240(1):53-60.
297. Hakonarson H, Thorvaldsson S, Helgadóttir A, Gudbjartsson D, Zink F, Andresdóttir M, et al. Effects of a 5-lipoxygenase-activating protein inhibitor on biomarkers associated with risk of myocardial infarction: a randomized trial. *Jama*. 2005;293(18):2245-56.
298. Elkhawad M, Rudd JH, Sarov-Blat L, Cai G, Wells R, Davies LC, et al. Effects of p38 mitogen-activated protein kinase inhibition on vascular and systemic inflammation in patients with atherosclerosis. *JACC: Cardiovascular Imaging*. 2012;5(9):911-22.
299. Newby LK, Marber MS, Melloni C, Sarov-Blat L, Aberle LH, Aylward PE, et al. Losmapimod, a novel p38 mitogen-activated protein kinase inhibitor, in non-ST-segment elevation myocardial infarction: a randomised phase 2 trial. *The Lancet*. 2014;384(9949):1187-95.
300. O'Donoghue ML, Glaser R, Cavender MA, Aylward PE, Bonaca MP, Budaj A, et al. Effect of losmapimod on cardiovascular outcomes in patients hospitalized with acute myocardial infarction: a randomized clinical trial. *Jama*. 2016;315(15):1591-9.

Table 1. The role of innate immune cells in atherosclerosis

Innate Immune Cells	Function during Atherosclerosis	Atheroprotective or Pro-atherogenic	Ref
Macrophage	<ul style="list-style-type: none"> Transformed into foam cells and form atherosclerotic plaque Imbalance between macrophage: atherosclerosis progress <p>❖ Classic or M1 Macrophages</p> <ul style="list-style-type: none"> Acute-phase proteins (APPs) leads to formation of M1 macrophages and progression of atherosclerotic plaque Destruction of fibrous layer, plaque rupture, and eventually myocardial infarction through production of collagenase (MMP1) and stromelysin (MMP-10) <p>❖ Alternative or M2 Macrophages</p> <ul style="list-style-type: none"> M2 macrophages predominate in the early stages of atherosclerosis May play an atheroprotective role M2 macrophages are categorized into four subclasses, including M2a, M2b, M2c, and M2d <p>❖ Other types of macrophages in atherosclerosis plaque include M4, Mox, M(Hb), and Mhem</p>	<p>Pro-atherogenic</p> <p>Atheroprotective and Pro-atherogenic</p>	<p>(14-19)</p> <p>(25, 27)</p> <p>(21, 32, 36, 46)</p> <p>(47-58)</p>
Monocyte	<p>❖ Two subgroups of mouse monocytes have been identified</p> <ul style="list-style-type: none"> Gr1⁺Ly6C^{high}CCR2⁺CX3CR^{low} (inflammatory) Gr1⁻Ly6C^{low}CCR2⁻CX3CR^{high} <ul style="list-style-type: none"> Ly-6C^{high} monocytes is raised in the first days following acute myocardial infarction Ly-6C^{low} monocytes is raised following acute myocardial infarction and can boost wound healing and regeneration of blood vessels Ly6C^{low} and Ly6C^{high} monocytes become macrophages M2 and M1, respectively Monocyte subtypes may play a vital role in inhibition and progression of atherosclerosis <p>❖ Monocytes in the humans are classified into three subgroups including</p> <ul style="list-style-type: none"> non-classical (CD14⁺ CD16⁺⁺) intermediate (CD14⁺⁺ CD16⁺) classical (CD14⁺⁺ CD16⁻) CD14⁺CD16^{high} monocytes promote atherosclerosis by the increased production of 	<p>Pro-atherogenic</p>	<p>(63-66)</p> <p>(71, 72)</p> <p>(73-79)</p>

	TNF- α		
Neutrophil	<ul style="list-style-type: none"> Elevated cholesterol levels: increase release of neutrophils and size of atherosclerotic plaque Direct relationship between the number of peripheral neutrophils and size of the atherosclerotic lesion Inflammatory activity in atherosclerotic plaques via secretion of various mediators ❖ Pentraxin 3, ROS, neutrophil elastases, MPO, leukotriene B4, neutrophil extracellular traps (NET), connexin, IFN-γ, and MMPs Neutrophils are also involved in development of atherosclerosis by secreting azurocidin, proteinase 3, and defensin. 	Pro-atherogenic	(105, 282, 283)
Mast Cell	<ul style="list-style-type: none"> Inactivation of mast cells by FcϵR1: decrease the number of inflammatory cells like macrophages and T cells in atherosclerotic plaque Activation of mast cells by TLR4 induces apoptosis in vascular smooth muscle cells of atherosclerotic plaques Mast cells cause plaque instability and exposing of necrotic nucleus through secretion of various mediators: ❖ Histamine, heparin, proteases (tryptase and chymase), and multiple cytokines (such as IL-6, IL-8, MCP1, TNF-α, and IFN-γ) 	Pro-atherogenic	(114, 115, 119, 120, 134)
Natural Killer Cell	<ul style="list-style-type: none"> NK cells are leading sources for production of IFN-γ: decrease in the IFN-γ levels significantly reduces atherosclerotic plaques in mice Overall, the function of NK cells in atherosclerosis has not been identified 	Has not been identified	(139, 140)
Dendritic cells	<ul style="list-style-type: none"> DCs have a pivotal role in both innate and adaptive immune systems Decreased number of DCs reduces lipid accumulation and number of foam cells and ultimately reduces size of atherosclerotic plaques Interactions of DCs with NK cells enhance production of IL-12 and IFN-γ: may have a role in formation and progression of atherosclerosis plaques The interaction between mDC, pDC, and other immune cells activates other immune cells, rupture plaques, and eventually leads to atherosclerosis Regulatory DCs interfere with anergy and depletion of inflammatory T cells and production and proliferation of atheroprotective Tregs 	Atheroprotective and Pro-atherogenic	(139, 145, 158, 165-168)

Table 2. The role of specific immune cells in atherosclerosis

Specific Immune Cells	Function during Atherosclerosis	Atheroprotective or Pro-atherogenic	Ref
T Lymphocytes	<ul style="list-style-type: none"> ❖ CD4⁺ T Cell <ul style="list-style-type: none"> ✓ Th1 Cells <ul style="list-style-type: none"> • The most abundant population of T cells present in atherosclerotic plaques • Activate other cells of the immune system by secretion IFN-γ • IFN-γ: prompts the formation of atherosclerotic plaques, formation of foam cells, and rupture of atherosclerotic plaques • IFN-γ: decreasing concentrations of serum cholesterol, scavenger receptor A, suppression of LDLR-related protein, MMP9, CD36, and inhibition of lipoprotein lipases ✓ Th2 Cells <ul style="list-style-type: none"> • Produce IL-25, IL-4, IL-5, IL-9, and IL-13 • Protective role against atherosclerosis because they suppress Th1 cells • IL-4: activation of mast cells, apoptosis of SMCs, plaque rupture, increased protease production, and decreased collagen production • IL-5: protective role in atherosclerosis ✓ Th17 Cells <ul style="list-style-type: none"> • Failure in signaling of IL-17 receptor (IL-17R: decrease in the size of atherosclerosis plaque and serum concentration of anti-oxLDL IgG antibodies) • IL-17 cytokine is related to lipid-rich, complicated, and unstable plaques ✓ Th9 Cells <ul style="list-style-type: none"> • Main source of IL-9 • Plasma level of IL-9 is higher in 	<p>Pro-atherogenic</p> <p>Atheroprotective</p> <p>Has not been identified</p> <p>Has not been</p>	<p>(173, 179, 186-190)</p> <p>(201, 203)</p> <p>(211, 212)</p> <p>(216, 217)</p> <p>(218, 221)</p> <p>(217)</p> <p>(230, 231)</p> <p>(249, 250)</p> <p>(251, 252,</p>

	<p>patients with atherosclerosis than healthy individuals</p> <p>✓ Th22 Cells</p> <ul style="list-style-type: none"> • The level of Th22 cells in the blood from patients with acute coronary syndrome was reported to increase compared with the healthy control group • Plasma level of IL-22 is higher in patients with symptomatic atherosclerosis than those with asymptomatic atherosclerosis <p>✓ T_{FH} Cells</p> <ul style="list-style-type: none"> • The pathway ICOS and its ligand (ICOSL) is vital for differentiation and sustaining of TFH cells • It has been reported that blocking the ICOS-ICOSL signaling pathway: decreases the progression of atherosclerosis and leads to a reduction in the number of T_{FH} cells <p>✓ CD28^{null} T Cells</p> <ul style="list-style-type: none"> • Patients with acute coronary syndrome have more TCD28null cells in their blood compared with healthy people, and their CD28null T cells are resistant against apoptosis <p>✓ Tregs</p> <ul style="list-style-type: none"> • TGF-β and IL-10 are two main cytokines of Treg cells which are responsible for many effects of Treg cells and have strong anti-atherogenic activities • Inhibit the activation of T cells and, subsequently, decrease the progression of atherosclerosis, can lead to differentiation of macrophages into the anti-inflammatory M2 	<p>identified</p> <p>Has not been identified</p> <p>Pro-atherogenic</p> <p>Pro-atherogenic</p> <p>Atheroprotective</p>	<p>254)</p>
--	---	--	-------------

	<p>phenotype, inhibit the formation of foam cells and differentiation of inflammatory M1 phenotype</p> <p>❖ CD8⁺ T Cells</p> <ul style="list-style-type: none"> • In patients with CAD, the blood level of cytotoxin-producing CD8⁺ T cells increases compared with healthy individuals • The cytotoxic activity of CD8⁺ T cells, relative to lesion-stabilizing cells (e.g., VSMCs), and production of inflammatory cytokines by CD8⁺ T cells may amplify inflammatory responses in atherosclerotic plaques • In contrast, the cytotoxic activity of CD8⁺ T cells, relative to APCs, may limit the progression of atherosclerosis 	Atheroprotective and Pro-atherogenic	
B Lymphocytes	<ul style="list-style-type: none"> • B cells can be divided into two distinct categories, including B1 and B2 • B2 cells incorporate marginal zone B cells and follicular B cells, and B1 cells include B-1b and B1a cells • B cells, which help in formation of lymphoid follicles, can play a role in atherosclerosis • Regulatory B cells are a subset of B lymphocytes, producing IL-10 • IL-10: decreases the risk of atherosclerosis by exerting apoptosis, modulating lipid metabolism, and inhibiting production of pro-inflammatory mediators 	Has not been identified	(262, 263, 270)

Table 3. Clinical trials of inflammation modulation in atherosclerotic cardiovascular disease.

Agent	Pathway/target	Clinical Trial (Ref. #)
Succinobulol	Oxidized LDL	ARISE (285)
BI-204 (MLDL1278A)	oxLDL antibody	GLACIER (286)
Inclacumab	P-selectin	SELECT-ACS (287)
Inclacumab	P-selectin	SELECT-CABG (288)
Colchicine	Neutrophil mobility	LoDoCo (289)
Colchicine	Neutrophil mobility	ColCot
Anakinra	IL-1RI	IL-HEART(290)
Canakinumab	IL-1 β	CANTOS (291)
Etanercept	TNF- α	ENTRACTE
Tocilizumab	IL-6	ENTRACTE
Methotrexate	IL-6, TNF	CIRT
Verespladib	sPLA2	VISTA-16 (292)
Darapladib	LpPLA2	STABILITY(293)
Darapladib	LpPLA2	SOLID-TIMI 52(294)
Atreleuton	5-LO inhibitor	1. FRANCIS (295) 2. VISTA-16 (296)
Veliflapon	FLAP inhibitor	Hakonarson et al. (297)
Losmapimod	p38 MAPK inhibitor	1. Elkhawad et al (298) 2. SOLSTICE (299) 3. LATITUDE-TIMI (300)

LDL: low-density lipoprotein; oxLD: oxidized low-density lipoprotein; sPLA2: secretory phospholipase A2; LpPLA2: lipoprotein-associated phospholipase A2; TNF: tumor necrosis factor; IL: interleukin; 5-LO: 5-lipoxygenase; FLAP: 5-lipoxygenase activating protein; p38 MAPK: p38 mitogen-activated protein kinase.

Figure 1: The role of macrophages in atherosclerosis. Atherosclerosis begins with the deposition of lipids. Endothelial activation increases the expression of leukocyte adhesion molecules such as E and P-selectins, VCAM-1 and ICAM-1 glycoproteins, and MCP-1 chemokine, which stimulates inflammatory monocytes by migration and infiltration. Circulating monocytes are activated, by adhering to endothelial cells (ECs), and are attracted to the atherosclerotic lesion site, by entering the subendothelial cell space. Inside the atherosclerosis plaque, macrophages absorb lipid particles and convert into foam cells, and finally form the early lesions of atherosclerosis. The reactive oxygen species (ROS) produced by macrophages lead to the oxidation of low-density lipoprotein (LDL) in the vascular wall and the formation of oxidized low-density lipoprotein (ox-LDL), which consequently lead to monocyte differentiation into macrophages and the formation of foam cells. Ox-LDL induces necrosis of foam cells, which forms a necrotic nucleus, leading to instability and rupture of atherosclerotic plaques. Pre-inflammatory macrophages M1 and anti-inflammatory M2 are polarized by the cytokines Th1 and Th2, respectively. Haem-induced phenotypes, such as M (Hb) and Mhem like M2, display anti-inflammatory effects such as resistance to lipid uptake and suppression of oxidative stress. Mox and M4 phenotypes have little potential for phagocytosis and, potentially have pre-inflammatory properties, by expressing proatherogenic markers.

Figure 2: Role of neutrophils in atherosclerosis. Neutrophils are of the most important components of innate immune responses and contain numerous inflammatory mediators that are freely activated. Activated oxygen species, myeloperoxidase (MPO), and matrix metalloproteinases (MMPs) are involved in extracellular matrix degradation and atherosclerotic plaque erosion. By binding to C1q, pentraxin induces complement activation and allows leukocytes to be recruited. On the luminal side, activated neutrophils secrete granular proteins, example.g., cathelicidin and cathepsin G, which directly or indirectly attract myeloid cells. Neutrophils activate and dysregulate endothelial layer cells (ECs) and extracellular matrix (ECM) by secretion of reactive oxygen species (ROS) and proteases on the luminal and intimal sides of atherosclerotic plaques. MPO attracts neutrophils and Ox-LDL. Ox-LDL is swallowed by macrophages and differentiates them into foam cells. Neutrophils can additionally be exposed to NETosis, in which neutrophils release extracellular traps that can play a role in thrombus formation, antigen-presenting cell activation, or endothelial cell cytotoxicity.

Figure 3: Role of T cells in atherosclerosis. Naive CD4 + T helper (Th) cells have several subsets, and in fact, stimulatory molecules stimulate T cells to express transcription factors that lead to differentiation into Th phenotypes. CD4 + T cells can have both pro-atherogenic and atheroprotective characteristics. Atherosclerotic lesions contain Th1, Th2, Th9, Th17, Th22, Treg, and follicular helper T (TFH) cells. Actually, these cells exert their atheroprotective effects by cytotoxicity activity against antigen-presenting cells (APCs) and the inhibition of production of inflammatory subsets of CD4 + T cells. Invariant natural killer T (iNKT) cells are activated by interplay of T cell receptor (TCR) receptors with CD1d molecules, comprising antigenic glycolipids presented in APCs. Activation of iNKT cells gives rise to the rapid discharge of cytokines associated with T helper 1 (Th1), Th2 and, Th17 cells and activate other immune system cells in atherosclerotic lesions. iNKT cells can additionally develop atherosclerosis through apoptosis induction of atherosclerosis plaque cells by releasing cytotoxic proteins such as perforin and granzyme B. $\gamma\delta$ T cells are a subset of T cells detected in atherosclerotic lesions of mice. $\gamma\delta$ T cells are a rich source of IL-17, thus, these cells can regulate atherosclerosis through the production of IL-17. Nevertheless, the definite role of $\gamma\delta$ T cells in atherosclerosis is unclear.