

Could changing the DNA methylation landscape promote the destruction of Epstein-Barr virus-associated cancers?

Article (Published Version)

Sinclair, Alison J (2021) Could changing the DNA methylation landscape promote the destruction of Epstein-Barr virus-associated cancers? *Frontiers in Cellular and Infection Microbiology*, 11. a695093 1-6. ISSN 2235-2988

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/99100/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.



Could Changing the DNA Methylation Landscape Promote the Destruction of Epstein-Barr Virus-Associated Cancers?

Alison J. Sinclair*

School of Life Sciences, University of Sussex, Brighton, United Kingdom

OPEN ACCESS

Edited by:

Kostas Paschos,
Imperial College London,
United Kingdom

Reviewed by:

Takayuki Murata,
Fujita Health University, Japan
Renfeng Li,
Virginia Commonwealth University,
United States

*Correspondence:

Alison J. Sinclair
a.j.sinclair@sussex.ac.uk

Specialty section:

This article was submitted to
Virus and Host,
a section of the journal
Frontiers in Cellular and
Infection Microbiology

Received: 14 April 2021

Accepted: 10 May 2021

Published: 28 May 2021

Citation:

Sinclair AJ (2021) Could Changing the DNA Methylation Landscape Promote the Destruction of Epstein-Barr Virus-Associated Cancers? *Front. Cell. Infect. Microbiol.* 11:695093. doi: 10.3389/fcimb.2021.695093

DNA methylation at CpG motifs provides an epigenetic route to regulate gene expression. In general, an inverse correlation between DNA hypermethylation at CpG motifs and gene expression is observed. Epstein Barr-virus (EBV) infects people and the EBV genome resides in the nucleus where either its replication cycle initiates or it enters a long-term latency state where the viral genome becomes hypermethylated at CpG motifs. Viral gene expression shows a largely inverse correlation with DNA hypermethylation. DNA methylation occurs through the action of DNA methyl transferase enzymes: writer DNA methyl transferases add methyl groups to specific regions of unmethylated DNA; maintenance DNA methyl transferases reproduce the pattern of DNA methylation during genome replication. The impact of DNA methylation is achieved through the association of various proteins specifically with methylated DNA and their influence on gene regulation. DNA methylation can be changed through altering DNA methyl transferase activity or through the action of enzymes that further modify methylated CpG motifs. Azacytidine prodrugs that are incorporated into CpG motifs during DNA replication are recognized by DNA methyl transferases and block their function resulting in hypomethylation of DNA. EBV-associated cancers have hypermethylated viral genomes and many carcinomas also have highly hypermethylated cellular genomes. Decitabine, a member of the azacytidine prodrug family, reactivates viral gene expression and promotes the recognition of lymphoma cells by virus-specific cytotoxic T-cells. For EBV-associated cancers, the impact of decitabine on the cellular genome and the prospect of combining decitabine with other therapeutic approaches is currently unknown but exciting.

Keywords: Epstein-Barr virus, DNA methylation, CpG motif, decitabine, demethylation, epigenetics

INTRODUCTION

DNA methylation at the 5-position of the cytosine ring of CpG motifs in DNA provides an epigenetic route to regulate gene expression (Klose and Bird, 2006). Extensive DNA methylation is generally associated with mammalian genes that are not expressed (Suzuki and Bird, 2008), with some cell-type specific genes showing a consistent pattern of DNA methylation found in key regions of the genome (Schmidl et al., 2009).

The methylation of DNA at CpG motifs is achieved and maintained by the interplay of several activities within the cell. 1. *De novo* DNA methyl transferases enzymes (also termed ‘writers’) are responsible for the initial addition of methyl groups to previously unmethylated CpG motifs. 2. Maintenance DNA methyl transferases enzymes are responsible for maintaining the DNA methylation pattern in daughter cells by adding methyl groups to the newly replicated strand of the hemi-methylated CpG motifs that are synthesized during genome replication (Klose and Bird, 2006). 3. A set of ‘eraser’ enzymes work in effective opposition to the DNA methyl transferases, by modifying the methylation mark (Tsiouplis et al., 2020). 4. A set of proteins termed ‘readers’ specifically recognize and bind to methylated CpG motifs and recruit other proteins to that region of the genome. Together, these enzymes act to mark the expression of linked genes (Mahmood and Rabbani, 2019). The mechanism by which the methylated state of the DNA is able to alter gene expression mediate through the action of methyl-DNA binding proteins that recruit chromatin modifiers (Du et al., 2015).

Epstein-Barr virus (EBV) is a common virus that infects people in a generally asymptomatic manner. The virus resides for life within the pool of B-lymphocytes (Thorley-Lawson and Babcock, 1999). For most people, EBV infection is asymptomatic, but it can cause infectious mononucleosis (IM, glandular fever) and a range of cancers both carcinomas (Gastric, Nasopharyngeal) and lymphomas (Burkitt’s lymphoma, Hodgkin’s lymphoma, NK/T-cell lymphoma, diffuse large B-cell lymphoma and post-transplant-like lymphoma) (Vetsika and Callan, 2004; Shannon-Lowe and Rickinson, 2019). EBV lytic replication and production of infectious virus is rare in cancer cells; by this stage the virus has entered into a latent state. The viral genome is maintained in the cancer cells by replicating once per cell cycle in time the host genome, but few viral genes are expressed. The EBV genome consists of double strand DNA and following infection it resides as a circular form in the nucleus of infected cells (Tsurumi et al., 2005; Hammerschmidt and Sugden, 2013). As with the cellular genome, the EBV genome is subject to DNA methylation at CpG motifs and, analogous to the cellular genome, DNA methylation of the viral genome is strongly linked to gene expression (Minarovits, 2006). The DNA methylation pattern of the EBV genome changes during its life-cycle, being largely unmethylated in the infectious virion, then gaining methylation following the infection of cells (Bergbauer et al., 2010). By the time that EBV-associated carcinomas and lymphomas develop, the viral genome is heavily methylated at CpG motifs and few of the approximately 90 viral genes are expressed (Shannon-Lowe and Rickinson, 2019).

The cellular genomes of cancers can also be impacted by DNA methylation. Importantly, some EBV-associated cancers exhibit very high levels of DNA methylation of the cellular genome that correlates with reduced expression of specific tumour suppressor genes (Bass et al., 2014; Stanland and Luftig, 2020).

Deliberately attempting to reduce the DNA methylation in EBV-associated cancer with the aim of activating the expression of cellular and viral genes has the potential to achieve two potentially therapeutic events. First, it may result in

reactivation of the expression of viral genes that expose the cells to attack by the immune system. Secondly, it may result in the expression important cellular genes that cause the cancer cells to stop proliferating or to die through apoptosis. Deliberately attempting to alter DNA methylation therefore presents an attractive research direction EBV which may lead to new therapeutic approaches to treat EBV-associated cancer in the future.

Enzymes That Regulate DNA Methylation

The process of establishing this epigenetic change to gene expression is undertaken through the deposition of DNA methylation onto the cytosine residues of CpG motifs in genomes by *de novo* DNA methyltransferase (DNMT) enzymes that add a methyl group to the cytosine of specific CpG motifs (Klose and Bird, 2006). The individual roles of *de novo* DNA methyl transferases DNMT3A and DNMT3B for methylating regulatory regions of developmental genes and the X-chromosome respectively, have been established and they are also candidates to accomplish *de novo* DNA methylation in somatic cells (Law and Jacobsen, 2010). Once established, the DNA methylation pattern is copied during each cycle of DNA replication by the action of the maintenance DNA methyl transferase, DNMT1 (Law and Jacobsen, 2010); thereby maintaining the set pattern of DNA methylation in daughter cells. As methylated DNA replicates, hemi-methylated CpG motifs consisting of the methylated template DNA strand and the newly replicated and so unmethylated strand are produced. DNMT1 recognizes the hemi-methylated CpG motif and adds a methyl group to the cytosine (**Figure 1**).

The effective removal of CpG DNA methylation on the host genome occurs through one of two routes. If the maintenance DNMT enzyme is compromised during DNA replication then hemi-methylated CpG motifs will be present after one round of replication with the potential for fully unmethylated CpG motifs after a second round of replication. The second route is achieved through the actions of a set of proteins (Ten-Eleven-Translocation (TET) family) that further modify the methyl-group on CpG motifs. (Tsiouplis et al., 2020).

DNA Methylation of the Host and Viral Genomes in EBV-Associated Cancer

Global methylome analysis of over 30 different cancer types by The Cancer Genome Atlas Network (TCGA) revealed that the highest level of hypermethylation of cellular DNA occurred in Epstein-Barr virus associated Gastric Carcinoma (Bass et al., 2014). Affected genes include a set of tumour suppressor genes (Okada et al., 2013; He et al., 2015; Nishikawa et al., 2018). Hypermethylation of cellular DNA is also seen in Epstein-Barr virus associated Burkitt’s lymphoma and Nasopharyngeal carcinoma, but to a lesser degree (Stanland and Luftig, 2020).

In EBV-associated carcinomas and lymphomas, the EBV genome is also subject to widespread DNA methylation and highly restricted patterns of viral gene expression (Minarovits, 2006). Immediately following infection of cells, transient low-level expression of many of the 90-viral genes occurs from the




FIGURE 1 | Impact of decitabine and 5' azacytidine on the maintenance methylation of DNA during genome replication. The presence of an isolated C and two CpG motifs in genome are shown prior to replication. The methylation group on each methyl cytosine of the central CpG motif is shown (blue triangle). Following one round of semi-conservative DNA replication hemi-methylated CpG motifs occur. These are recognized by DNMT1 and a methyl group added to the unmethylated cytosine. Thus the end product has the same methylation state as the genome prior to replication. In the presence of decitabine or 5'azacytidine, these modified bases are incorporated into the newly synthesized strand (shown in red). The hemi-methylated CpG motifs are recognized by DNMT1, and are covalently bound to the modified cytosine bases. The genome remains in a hemi-methylated state.

unmethylated EBV genome. However, the viral genome is rapidly subject to DNA methylation at CpG motifs and settles into one of several highly restricted sets of viral latency gene expression (Tierney et al., 2000; Bergbauer et al., 2010). Three major patterns are common in cancer (latency I, II (also termed IIa) and III), but other patterns of viral gene expression exist including Wp-restricted (Kelly et al., 2009) and IIb latency (Price and Luftig, 2015). Around 90% of the EBV genes are not expressed in any lymphomas and carcinomas, and for the majority of the EBV-associated cancers, the promoters driving expression of the genes generating the most immunogenic T-cell responses (EBNA 2 and 3 family) are not expressed (Minarovits, 2006). In addition, the lytic replication cycle genes (especially BZLF1, BRLF1, BMRF1 and BMLF1) which are highly immunogenic are also silent (Taylor et al., 2015). So, the EBV genome itself provides a wealth of immunogenic genes that are often silenced by DNA hypermethylation.

Clues about how the hypermethylation of cellular and viral DNA occurs in EBV-associated cancer comes from cell culture experiments where it was demonstrated that *de novo* methylation of cellular DNA occurs within days of EBV infection (Queen et al., 2013). Two EBV genes have been implicated: LMP1 and LMP2A. LMP1 activates the expression of both *de novo* and maintenance DNA methyl transferases (Tsai et al., 2006; Peng et al., 2016) and so could both aid the initiation and ensure the maintenance of the DNA methylation changes. LMP1 is expressed in some but not all EBV-associated cancer, however, it has been shown to be expressed transiently after EBV infection of gastric epithelial cells (Matsusaka et al., 2017) in addition to during the immortalization of B-lymphocytes by EBV. It is therefore conceivable that LMP1 might play a transient early role in reprogramming DNA methylation in EBV-associated cancers. In addition, LMP2A which is expressed in some but not all, EBV-associated cancers, drives the upregulation of the

maintenance DNA methyl transferase (Hino et al., 2009) and the down regulation of TET gene expression (Namba-Fukuyo et al., 2016). These changes indicate that LMP2A is also a candidate to contribute to global increases in DNA methylation in EBV-associated cancers.

A further piece in this puzzle is the enduring impact of EBV on hypermethylation of the cellular genome even after the viral genome has been lost from the cells (Birdwell et al., 2014). So, clues about the mechanisms by which the hypermethylation of the cellular and viral genomes occurs have been identified and it is possible that LMP1 and LMP2A play transient roles in reprogramming DNA methylation early after entry into cells.

The correlation of EBV-associated gastric cancer with DNA hypermethylation of the cellular genome, the potential contribution of EBV genes to changing the expression of components of the DNA methylation machinery and the relevance of the genes that are potentially silenced by hypermethylation in EBV-associated gastric cancer all support the hypothesis that actively demethylating the genome of EBV-associated cancer cells may be beneficial.

Impact of Changing the DNA Methylation in EBV-Infected Cells

Attempts to alter the DNA methylation within EBV-infected cells focus on the maintenance DNA methyltransferase. The impact of totally inhibiting the action of the maintenance DNA methyl transferase would be the hypomethylation of the cellular and viral genomes within 1-2 rounds of cell replication (Mani and Herceg, 2010). The reduction in the DNA methylation landscape resulting from this in EBV infected cells is predicted to have impacts on the expression of both viral and cellular genes with the potential for immunogenic viral genes rendering the cancer cells susceptible to attack by the immune system and of tumour suppressor genes reducing growth and survival of the

cancer cell. However, global demethylation is not a targeted approach and it may also drive unfavorable changes in cellular gene expression.

Cytidine analogues that irreversibly inhibit the maintenance DNA methyl transferases and so promote hypomethylation of DNA were identified over 50-years ago (Sorm et al., 1964; Sorm and Vesely, 1968) - 5'Azacytidine (4-Amino-1-(β -D-ribofuranosyl)-1,3,5-triazin-2(1H)-one; azacytidine, 5-aza-CR; Vidaza[®]) and its deoxyribose version, Decitabine (2'-Deoxy-5-azacytidine, 4-Amino-1-(2-deoxy- β -D-ribofuranosyl)-1,3,5-triazin-2(1H)-one, 5-aza-CdR; Dacogen[®]). Both cytidine analogues are processed within cells to the triphosphate derivatives and can be incorporated into newly synthesized DNA during the S-phase of the cell replication cycle (**Figure 1**) (Mani and Herceg, 2010). Once embedded within the genome those modified bases form hemi-methylated CpG motifs that are recognized by the maintenance DNA methyl transferase and they irreversibly trap and inhibit the DNMT1, leading to rapid hypomethylation of CpG motifs throughout the genome (**Figure 1**) (Howell et al., 2010; Mani and Herceg, 2010).

Preclinical studies showed that both 5'azacytidine and decitabine are highly active cytotoxic agents against many types of cancer cells (Howell et al., 2010). Furthermore, clinical studies showed that both 5'azacytidine and decitabine are beneficial for people with some forms of haematological cancer (Silverman et al., 2002; Kantarjian et al., 2006; Fenaux et al., 2009; Benton et al., 2014; Mayer et al., 2014). A 5'Azacytidine small

scale trial in patients with EBV-associated cancers revealed that demethylation of the viral genome was successfully achieved (Chan et al., 2004).

Impact of Decitabine as a Demethylation Agent of the EBV Genome in EBV-Infected Cells

An exciting new development in the fight to reprogram the epigenetics of EBV-infected cells to make them sensitized to immunotherapy was reported recently (Dalton et al., 2020). This builds on previous successes of infusing virus-antigen-specific autologous or allogeneic CTLs to treat EBV cancers (Heslop et al., 2010; Bollard et al., 2014; Bollard and Barrett, 2014; Chia et al., 2014; Kazi et al., 2019; Prockop et al., 2020). Dalton hypothesized that if they could reactivate expression of viral antigens using sub-cytotoxic doses of drugs, the cancer cells may become primed for immune attack. In a search to re-purpose drugs that could promote the expression of antigenic viral genes in the aggressive B-lymphomas that express the most restricted pattern of EBV gene expression (latency I), they identified decitabine as a strong candidate to reprogram the expression of immunogenic viral latency genes. Decitabine induced the expression of the two key viral promoters, with the resulting proteins identified in 27-58% of cells in a cell culture model system (**Figure 2**) (Dalton et al., 2020). Excitingly, the reprogramming of viral gene expression was reproduced in a more




FIGURE 2 | Reversing epigenetic silencing of the viral genome allows recognition by T-cells. EBV-associated cancer cells are shown in blue. In the top panel the viral genome (red oval) is hyper methylated at CpG motifs and largely silent. MHC is shown at the cell surface (blue shape). In the bottom panel, following decitabine treatment, the genome is de-methylated (red dashed-line oval), reactivated and expresses immunogenic viral proteins (red) that are displayed with MHC as peptides at the surface of the cell. This allows the EBV-specific T-cells (yellow) to recognize the cancer cells.

physiologically challenging mouse xenograft tumour model and the change in gene expression was shown to have a durable impact. Importantly, EBV-specific CTLs were able to recognise and kill the decitabine-treated tumour cells *in vitro* and critically they were shown to infiltrate the xenograft tumours *in vivo* (Figure 1) (Dalton et al., 2020), providing proof-of-principle.

DISCUSSION

The pressing question is whether decitabine treatment could provide a relevant clinical approach to treat EBV-associated lymphomas and carcinomas? Decitabine has a proven potential to drive hypomethylation of the viral genome and to prime EBV-associated lymphoma cells to re-express immunogenic viral genes (Dalton et al., 2020). This provides the first part of a potential therapeutic strategy. In combination with the action of the patient's existing immune response, this may be sufficient to boost an immune attack, or this could be supplemented by infusion of EBV-specific CTLs or T-cells expressing EBV-specific chimeric antigen receptors (CAR-T therapy) (Chicaybam et al., 2019; Dragon et al., 2020; Prockop et al., 2020; Heslop et al., 2021). However, a potential limitation to this approach is that the EBV genome was not reactivated in every lymphoma cell following decitabine treatment (Dalton et al., 2020), and so not all tumor cells would be rendered susceptible to immune attack. Whether reactivation could be stimulated to generate a homogeneous

response remains an open question. Interestingly, the sub-cytopathic dose of decitabine used appeared sufficient to stop the lymphoma growing (Dalton et al., 2020). Alterations to host gene expression might also occur as a result of decitabine and may negatively impact on the growth or survival of tumor cells. The recent discovery that LMP1 directs the expression of a set of tumor-associated antigens in a B-lymphoma that are recognized by T-cells (Choi et al., 2021) supports this avenue, although it is not known yet whether the changes in gene expression involves a change in DNA methylation. Changes in expression of cellular genes may be especially relevant for EBV-associated carcinomas, where a high proportion of the cellular genome is methylated. In addition, in non-EBV infected cells it has recently been shown that decitabine can act in combination with chemotherapy to improve its efficacy (Wu et al., 2019). This suggests another avenue of research for combined EBV-associated cancer treatments.

The recent advances that allow for the reversal of DNA methylation and epigenetic silencing of viral gene expression and those that develop specific viral immunotherapies provide promising avenues of research for EBV-associated cancers in the future.

AUTHOR CONTRIBUTIONS

The author confirms being the sole contributor of this work and has approved it for publication.

REFERENCES

- Bass, A. J., Thorsson, V., Shmulevich, I., Reynolds, S. M., Miller, M., Bernard, B., et al. (2014). Comprehensive Molecular Characterization of Gastric Adenocarcinoma. *Nature* 513 (7517), 202–209. doi: 10.1038/nature13480
- Benton, C. B., Thomas, D. A., Yang, H., Ravandi, F., Rytting, M., O'Brien, S., et al. (2014). Safety and Clinical Activity of 5-Aza-2'-Deoxycytidine (Decitabine) With or Without Hyper-CVAD in Relapsed/Refractory Acute Lymphocytic Leukaemia. *Br. J. Haematol* 167 (3), 356–365. doi: 10.1111/bjh.13050
- Bergbauer, M., Kalla, M., Schmeinek, A., Gobel, C., Rothbauer, U., Eck, S., et al. (2010). CpG-methylation Regulates a Class of Epstein-Barr Virus Promoters. *PLoS Pathog.* 6 (9), e1001114. doi: 10.1371/journal.ppat.1001114
- Birdwell, C. E., Queen, K. J., Kilgore, P. C., Rollyson, P., Trutschl, M., Cvek, U., et al. (2014). Genome-Wide DNA Methylation as an Epigenetic Consequence of Epstein-Barr Virus Infection of Immortalized Keratinocytes. *J. Virol.* 88 (19), 11442–11458. doi: 10.1128/JVI.00972-14
- Bollard, C. M., and Barrett, A. J. (2014). Cytotoxic T Lymphocytes for Leukemia and Lymphoma. *Hematol. Am. Soc. Hematol. Educ. Program* 2014 (1), 565–569. doi: 10.1182/asheducation-2014.1.565
- Bollard, C. M., Gottschalk, S., Torrano, V., Diouf, O., Ku, S., Hazrat, Y., et al. (2014). Sustained Complete Responses in Patients With Lymphoma Receiving Autologous Cytotoxic T Lymphocytes Targeting Epstein-Barr Virus Latent Membrane Proteins. *J. Clin. Oncol.* 32 (8), 798–808. doi: 10.1200/JCO.2013.51.5304
- Chan, A. T., Tao, Q., Robertson, K. D., Flinn, I. W., Mann, R. B., Klencke, B., et al. (2004). Azacitidine Induces Demethylation of the Epstein-Barr Virus Genome in Tumors. *J. Clin. Oncol.* 22 (8), 1373–1381. doi: 10.1200/JCO.2004.04.185
- Chia, W. K., Teo, M., Wang, W. W., Lee, B., Ang, S. F., Tai, W. M., et al. (2014). Adoptive T-cell Transfer and Chemotherapy in the First-Line Treatment of Metastatic and/or Locally Recurrent Nasopharyngeal Carcinoma. *Mol. Ther.* 22 (1), 132–139. doi: 10.1038/mt.2013.242
- Chicaybam, L., Abdo, L., Carneiro, M., Peixoto, B., Viegas, M., de Sousa, P., et al. (2019). Car T Cells Generated Using Sleeping Beauty Transposon Vectors and Expanded With an EBV-Transformed Lymphoblastoid Cell Line Display Antitumor Activity in Vitro and In Vivo. *Hum. Gene Ther.* 30 (4), 511–522. doi: 10.1089/hum.2018.218
- Choi, I. K., Wang, Z., Ke, Q., Hong, M., Paul, D. W. Jr., Fernandes, S. M., et al. (2021). Mechanism of EBV Inducing Anti-Tumour Immunity and its Therapeutic Use. *Nature* 590 (7844), 157–162. doi: 10.1038/s41586-020-03075-w
- Dalton, T., Doubrovina, E., Pankov, D., Reynolds, R., Scholze, H., Selvakumar, A., et al. (2020). Epigenetic Reprogramming Sensitizes Immunologically Silent EBV+ Lymphomas to Virus-Directed Immunotherapy. *Blood* 135 (21), 1870–1881. doi: 10.1182/blood.2019004126
- Dragon, A. C., Zimmermann, K., Nerretre, T., Sandfort, D., Lahrberg, J., Kloss, S., et al. (2020). CAR-T Cells and TRUCKs That Recognize an EBNA-3C-derived Epitope Presented on HLA-B*35 Control Epstein-Barr Virus-Associated Lymphoproliferation. *J. Immunother. Cancer* 8 (2), 1–16. doi: 10.1136/jitc-2020-000736
- Du, Q., Luu, P. L., Stirzaker, C., and Clark, S. J. (2015). Methyl-CpG-binding Domain Proteins: Readers of the Epigenome. *Epigenomics* 7 (6), 1051–1073. doi: 10.2217/epi.15.39
- Fenaux, P., Mufti, G. J., Hellstrom-Lindberg, E., Santini, V., Finelli, C., Giagounidis, A., et al. (2009). Efficacy of Azacitidine Compared With That of Conventional Care Regimens in the Treatment of Higher-Risk Myelodysplastic Syndromes: A Randomised, Open-Label, Phase III Study. *Lancet Oncol.* 10 (3), 223–232. doi: 10.1016/S1470-2045(09)70003-8
- Hammerschmidt, W., and Sugden, B. (2013). Replication of Epstein-Barr Viral DNA. *Cold Spring Harb. Perspect. Biol.* 5 (1), a013029. doi: 10.1101/cshperspect.a013029
- He, D., Zhang, Y. W., Zhang, N. N., Zhou, L., Chen, J. N., Jiang, Y., et al. (2015). Aberrant Gene Promoter Methylation of P16, FHIT, Crbp1, WWOX, and DLC-1 in Epstein-Barr Virus-Associated Gastric Carcinomas. *Med. Oncol.* 32 (4), 92. doi: 10.1007/s12032-015-0525-y
- Heslop, H. E., Sharma, S., and Rooney, C. M. (2021). Adoptive T-Cell Therapy for Epstein-Barr Virus-Related Lymphomas. *J. Clin. Oncol.* 39 (5), 514–524. doi: 10.1200/JCO.20.01709

- Heslop, H. E., Slobod, K. S., Pule, M. A., Hale, G. A., Rousseau, A., Smith, C. A., et al. (2010). Long-Term Outcome of EBV-Specific T-Cell Infusions to Prevent or Treat EBV-related Lymphoproliferative Disease in Transplant Recipients. *Blood* 115 (5), 925–935. doi: 10.1182/blood-2009-08-239186
- Hino, R., Uozaki, H., Murakami, N., Ushiku, T., Shinozaki, A., Ishikawa, S., et al. (2009). Activation of DNA Methyltransferase 1 by EBV Latent Membrane Protein 2A Leads to Promoter Hypermethylation of PTEN Gene in Gastric Carcinoma. *Cancer Res.* 69 (7), 2766–2774. doi: 10.1158/0008-5472.CAN-08-3070
- Howell, P. M., Liu, Z., and Khong, H. T. (2010). Demethylating Agents in the Treatment of Cancer. *Pharmaceut (Basel)* 3 (7), 2022–2044. doi: 10.3390/ph3072022
- Kantarjian, H., Issa, J. P., Rosenfeld, C. S., Bennett, J. M., Albitar, M., DiPersio, J., et al. (2006). Decitabine Improves Patient Outcomes in Myelodysplastic Syndromes: Results of a Phase III Randomized Study. *Cancer* 106 (8), 1794–1803. doi: 10.1002/cncr.21792
- Kazi, S., Mathur, A., Wilkie, G., Cheal, K., Battle, R., McGowan, N., et al. (2019). Long-Term Follow Up After Third-Party Viral-Specific Cytotoxic Lymphocytes for Immunosuppression- and Epstein-Barr Virus-Associated Lymphoproliferative Disease. *Haematologica* 104 (8), e356–e359. doi: 10.3324/haematol.2018.207548
- Kelly, G. L., Long, H. M., Stylianou, J., Thomas, W. A., Leese, A., Bell, A. I., et al. (2009). An Epstein-Barr Virus Anti-Apoptotic Protein Constitutively Expressed in Transformed Cells and Implicated in Burkitt Lymphomagenesis: The Wp/BHRF1 Link. *PLoS Pathog.* 5 (3), e1000341. doi: 10.1371/journal.ppat.1000341
- Klose, R. J., and Bird, A. P. (2006). Genomic DNA Methylation: The Mark and its Mediators. *Trends Biochem. Sci.* 31 (2), 89–97. doi: 10.1016/j.tibs.2005.12.008
- Law, J. A., and Jacobsen, S. E. (2010). Establishing, Maintaining and Modifying DNA Methylation Patterns in Plants and Animals. *Nat. Rev. Genet.* 11 (3), 204–220. doi: 10.1038/nrg2719
- Mahmood, N., and Rabbani, S. A. (2019). DNA Methylation Readers and Cancer: Mechanistic and Therapeutic Applications. *Front. Oncol.* 9, 489. doi: 10.3389/fonc.2019.00489
- Mani, S., and Herceg, Z. (2010). DNA Demethylating Agents and Epigenetic Therapy of Cancer. *Adv. Genet.* 70, 327–340. doi: 10.1016/B978-0-12-380866-0.60012-5
- Matsusaka, K., Funata, S., Fukuyo, M., Seto, Y., Aburatani, H., Fukayama, M., et al. (2017). Epstein-Barr Virus Infection Induces Genome-Wide *De Novo* DNA Methylation in Non-Neoplastic Gastric Epithelial Cells. *J. Pathol.* 242 (4), 391–399. doi: 10.1002/path.4909
- Mayer, J., Arthur, C., Delaunay, J., Mazur, G., Thomas, X. G., Wierzbowska, A., et al. (2014). Multivariate and Subgroup Analyses of a Randomized, Multinational, Phase 3 Trial of Decitabine vs Treatment Choice of Supportive Care or Cytarabine in Older Patients With Newly Diagnosed Acute Myeloid Leukemia and Poor- or Intermediate-Risk Cytogenetics. *BMC Cancer* 14, 69. doi: 10.1186/1471-2407-14-69
- Minarovits, J. (2006). Epigenotypes of Latent Herpesvirus Genomes. *Curr. Top. Microbiol. Immunol.* 310, 61–80. doi: 10.1007/3-540-31181-5_5
- Namba-Fukuyo, H., Funata, S., Matsusaka, K., Fukuyo, M., Rahmutulla, B., Mano, Y., et al. (2016). TET2 Functions as a Resistance Factor Against DNA Methylation Acquisition During Epstein-Barr Virus Infection. *Oncotarget* 7 (49), 81512–81526. doi: 10.18632/oncotarget.13130
- Nishikawa, J., Iizasa, H., Yoshiyama, H., Shimokuri, K., Kobayashi, Y., Sasaki, S., et al. (2018). Clinical Importance of Epstein(-)Barr Virus-Associated Gastric Cancer. *Cancers (Basel)* 10 (6), 1–13. doi: 10.3390/cancers10060167
- Okada, T., Nakamura, M., Nishikawa, J., Sakai, K., Zhang, Y., Saito, M., et al. (2013). Identification of Genes Specifically Methylated in Epstein-Barr Virus-Associated Gastric Carcinomas. *Cancer Sci.* 104 (10), 1309–1314. doi: 10.1111/cas.12228
- Peng, H., Chen, Y., Gong, P., Cai, L., Lyu, X., Jiang, Q., et al. (2016). Higher Methylation Intensity Induced by EBV LMP1 Via NF-kappaB/DNMT3b Signaling Contributes to Silencing of PTEN Gene. *Oncotarget* 7 (26), 40025–40037. doi: 10.18632/oncotarget.9474
- Price, A. M., and Luftig, M. A. (2015). To be or Not IIb: A Multi-Step Process for Epstein-Barr Virus Latency Establishment and Consequences for B Cell Tumorigenesis. *PLoS Pathog.* 11 (3), e1004656. doi: 10.1371/journal.ppat.1004656
- Prockop, S., Doubrovina, E., Suser, S., Heller, G., Barker, J., Dahi, P., et al. (2020). Off-the-Shelf EBV-Specific T Cell Immunotherapy for Rituximab-Refractory EBV-associated Lymphoma Following Transplantation. *J. Clin. Invest.* 130 (2), 733–747. doi: 10.1172/JCI121127
- Queen, K. J., Shi, M., Zhang, F., Cvek, U., and Scott, R. S. (2013). Epstein-Barr Virus-Induced Epigenetic Alterations Following Transient Infection. *Int. J. Cancer* 132 (9), 2076–2086. doi: 10.1002/ijc.27893
- Schmidl, C., Klug, M., Boeld, T. J., Andreesen, R., Hoffmann, P., Edinger, M., et al. (2009). Lineage-Specific DNA Methylation in T Cells Correlates With Histone Methylation and Enhancer Activity. *Genome Res.* 19 (7), 1165–1174. doi: 10.1101/gr.091470.109
- Shannon-Lowe, C., and Rickinson, A. (2019). The Global Landscape of EBV-Associated Tumors. *Front. Oncol.* 9, 713. doi: 10.3389/fonc.2019.00713
- Silverman, L. R., Demakos, E. P., Peterson, B. L., Kornblith, A. B., Holland, J. C., Odchimar-Reissig, R., et al. (2002). Randomized Controlled Trial of Azacitidine in Patients With the Myelodysplastic Syndrome: A Study of the Cancer and Leukemia Group B. *J. Clin. Oncol.* 20 (10), 2429–2440. doi: 10.1200/JCO.2002.04.117
- Sorm, F., Piskala, A., Cihak, A., and Vesely, J. (1964). 5-Azacytidine, a New, Highly Effective Cancerostatic. *Experientia* 20 (4), 202–203. doi: 10.1007/BF02135399
- Sorm, F., and Vesely, J. (1968). Effect of 5-Aza-2'-Deoxycytidine Against Leukemic and Hemopoietic Tissues in AKR Mice. *Neoplasma* 15 (4), 339–343.
- Stanland, L. J., and Luftig, M. A. (2020). The Role of EBV-Induced Hypermethylation in Gastric Cancer Tumorigenesis. *Viruses* 12 (11), 1–16. doi: 10.3390/v12111222
- Suzuki, M. M., and Bird, A. (2008). DNA Methylation Landscapes: Provocative Insights From Epigenomics. *Nat. Rev. Genet.* 9 (6), 465–476. doi: 10.1038/nrg2341
- Taylor, G. S., Long, H. M., Brooks, J. M., Rickinson, A. B., and Hislop, A. D. (2015). The Immunology of Epstein-Barr Virus-Induced Disease. *Annu. Rev. Immunol.* 33, 787–821. doi: 10.1146/annurev-immunol-032414-112326
- Thorley-Lawson, D. A., and Babcock, G. J. (1999). A Model for Persistent Infection With Epstein-Barr Virus: The Stealth Virus of Human B Cells. *Life Sci.* 65 (14), 1433–1453. doi: 10.1016/S0024-3205(99)00214-3
- Tierney, R. J., Kirby, H. E., Nagra, J. K., Desmond, J., Bell, A. I., and Rickinson, A. B. (2000). Methylation of Transcription Factor Binding Sites in the Epstein-Barr Virus Latent Cycle Promoter Wp Coincides With Promoter Down-Regulation During Virus-Induced B-cell Transformation. *J. Virol.* 74 (22), 10468–10479. doi: 10.1128/JVI.74.22.10468-10479.2000
- Tsai, C. L., Li, H. P., Lu, Y. J., Hsueh, C., Liang, Y., Chen, C. L., et al. (2006). Activation of DNA Methyltransferase 1 by EBV Lmp1 Involves C-Jun NH(2)-terminal Kinase Signaling. *Cancer Res.* 66 (24), 11668–11676. doi: 10.1158/0008-5472.CAN-06-2194
- Tsiouplis, N. J., Bailey, D. W., Chiou, L. F., Wissink, F. J., and Tsagaratou, A. (2020). Tet-Mediated Epigenetic Regulation in Immune Cell Development and Disease. *Front. Cell Dev. Biol.* 8, 623948. doi: 10.3389/fcell.2020.623948
- Tsurumi, T., Fujita, M., and Kudoh, A. (2005). Latent and Lytic Epstein-Barr Virus Replication Strategies. *Rev. Med. Virol.* 15 (1), 3–15. doi: 10.1002/rmv.441
- Vetsika, E.-K., and Callan, M. (2004). Infectious Mononucleosis and Epstein-Barr Virus. *Expert Rev. Mol. Med.* 6 (23), 1–16. doi: 10.1017/S1462399404008440
- Wu, M., Sheng, L., Cheng, M., Zhang, H., Jiang, Y., Lin, S., et al. (2019). Low Doses of Decitabine Improve the Chemotherapy Efficacy Against Basal-Like Bladder Cancer by Targeting Cancer Stem Cells. *Oncogene* 38 (27), 5425–5439. doi: 10.1038/s41388-019-0799-1

Conflict of Interest: The author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2021 Sinclair. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.