

The trafficking and targeting of P2X receptors

Article (Published Version)

Robinson, Lucy E and Murrell-Lagnado, Ruth D (2013) The trafficking and targeting of P2X receptors. *Frontiers in Cellular Neuroscience*, 7. p. 233. ISSN 1662-5102

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/52585/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.



The trafficking and targeting of P2X receptors

Lucy E. Robinson and Ruth D. Murrell-Lagnado*

Department of Pharmacology, University of Cambridge, Cambridge, UK

Edited by:

Baljit S. Khakh, University of California
Los Angeles, USA

Reviewed by:

Daryl L. Davies, University of Southern
California School of Pharmacy, USA
Richard Evans, University of Leicester,
UK

*Correspondence:

Ruth D. Murrell-Lagnado, Department
of Pharmacology, University of
Cambridge, Tennis Court Road,
Cambridge, CB2 1PD, UK
e-mail: rdm1003@cam.ac.uk

The functional expression of P2X receptors at the plasma membrane is dependent on their trafficking along secretory and endocytic pathways. There are seven P2X receptor subunits, and these differ in their subcellular distributions because they have very different trafficking properties. Some are retained within the endoplasmic reticulum (ER), while others are predominantly at the cell surface or within endosomes and lysosomes. Changes in recruitment of receptors to and from the plasma membrane provides a way of rapidly up- or down-regulating the cellular response to adenosine triphosphate (ATP). An additional layer of regulation is the targeting of these receptors within the membranes of each compartment, which affects their stability, function and the nature of the effector proteins with which they form signaling complexes. The trafficking and targeting of P2X receptors is regulated by their interactions with other proteins and with lipids and we can expect this to vary in a cell-type specific manner and in response to changes in the environment giving rise to differences in receptor activity and function.

Keywords: P2X receptor trafficking, lipid rafts, receptor endocytosis, receptor regulation, P2X receptor targeting

INTRODUCTION

P2X receptors open an integral ion channel at the plasma membrane in response to binding extracellular adenosine triphosphate (ATP). Some subtypes of P2X receptor are predominantly within intracellular membranes, but there is no compelling evidence that the receptors on intracellular membranes open in response to ATP, at least in mammals. Nonetheless, this remains an intriguing possibility, given that the distantly related P2X-like receptors in *Dictyostelium* are located and function within contractile vacuoles (Fountain et al., 2007; Ludlow et al., 2009; Sivaramakrishnan and Fountain, 2012a,b; Baines et al., 2013). The best-established role of the internal mammalian P2X receptors is, therefore, to regulate the expression and activity of receptors at the cell surface. Here we consider three related issues concerning the targeting and trafficking of P2X receptors: first, primary location, and the amino acid motifs which determine it; second, regulation of mobility both within the plasma membrane and between the plasma membrane and intracellular membranes; third, targeting to lipid rafts and the effects of the lipid environment on receptor signaling.

SUBCELLULAR LOCALIZATION OF P2X RECEPTORS

Trimeric P2X receptor complexes assemble and are core glycosylated within the endoplasmic reticulum (ER) and then traffic via the trans-Golgi network (TGN) to the plasma membrane.

They are subsequently internalized and either recycled back to the surface or targeted to late endosomes and lysosomes. The kinetics of these processes determines receptor distribution.

ER RESIDENT P2X RECEPTORS

P2X receptors are predominantly found within the ER, at the plasma membrane or within late endosomes and lysosomes, dependent upon the subtype (Figure 1). The only full-length

P2X receptor that is retained within the ER and is therefore non-functional is P2X6 (Ormond et al., 2006). Imaging of P2X6 receptors by atomic force microscopy indicates that the subunits do not assemble to form stable homotrimeric complexes, but they do form stable heterotrimers with either P2X2 or P2X4 (Bobanovic et al., 2002; Barrera et al., 2005, 2007; Ormond et al., 2006). The P2X2/6 and P2X4/6 receptors are expressed as functional receptors at the plasma membrane and have trafficking properties that resemble the P2X2 and P2X4 homomeric receptors respectively. In the category of ER resident P2X receptors there is also the human P2X5 receptor. Although the full-length P2X5 receptor traffics to the cell surface, the predominant allele expressed in most humans gives rise to an exon 10-deleted variant which is retained in the ER (Bo et al., 2003; Kotnis et al., 2010; Compan et al., 2012).

PLASMA MEMBRANE P2X RECEPTORS

Two subtypes that traffic relatively slowly through the secretory pathway and hence often appear to have a predominantly ER distribution are P2X2 and P2X7 receptors. P2X2 receptors are stably expressed at the plasma membrane, but, when heterologously expressed, they accumulate slowly at the cell surface. This slow traffic might be important for facilitating interactions with other proteins along the way (Bobanovic et al., 2002). For example, in spinal cord neurons, intracellular P2X2 receptors interact with GABA_A receptors and co-traffic to the surface (Shrivastava et al., 2011). Another protein that interacts with P2X2 receptors to regulate its targeting to synapses is the beta-amyloid precursor protein-binding protein Fe65 (Masin et al., 2006). There is also the neuronal calcium sensor, visinin-like protein-1 (VILIP-1) that interacts with P2X2 in a calcium-dependent manner (Chau-mont et al., 2008). These interacting proteins affect the stability, targeting and function of the receptors at the plasma membrane.




FIGURE 1 | The subcellular distribution of P2X receptors. P2X receptor subtypes differ in their trafficking properties and hence are localized to different subcellular compartments. P2X6 receptors are retained within the ER but can assemble with P2X4 and P2X6 subunits to form heterotrimer that traffic to the cell surface. The predominant human allele of P2X5 lacks exon 10 and is also retained in the ER. P2X2 and P2X7 receptors traffic

relatively slowly through the secretory pathway but are stably expressed at the surface. P2X1 receptors are expressed at the cell surface but rapidly cycle to and from recycling endosomes. P2X3 and P2X4 receptors are constitutively internalized and delivered to late endosomes and lysosomes. Within the lysosomes, P2X3 receptors are rapidly degraded but P2X4 receptors resist degradation and can recycle back to the surface.

P2X7 receptor trafficking is dependent upon cell-type and species. For example, in human monocytes and lymphocytes, P2X7 receptors are predominantly intracellular, but upon differentiation of monocytes to macrophages receptors locate to the plasma membrane (Hickman et al., 1994; Gu et al., 2000; Gudipaty et al., 2001). Native P2X7 receptors in rodent microglia and macrophages are also predominantly at the plasma membrane (Boumechache et al., 2009). What regulates the rate at which P2X7 receptors traffic from the ER to the cell surface is unknown, although mutagenesis analysis suggests that it involves the cytoplasmic C-terminal domain of the receptor (Denlinger et al., 2003; Smart et al., 2003).

P2X1 receptors are predominantly at the cell surface, and at least one interacting protein has been identified, namely heat shock protein 90, which further promotes their trafficking and plasma membrane expression (Lalo et al., 2010, 2012). At the plasma membrane, P2X1 receptors differ in trafficking behavior from P2X2 receptors. P2X2 receptors are relatively stable and show little constitutive internalization over a period of an hour (Bobanovic et al., 2002). In contrast, measurements of P2X1 receptor mobility by fluorescence recovery after photobleaching (FRAP) shows that receptors undergo considerable internalization and recycling over a similar time period (Lalo et al., 2010). Their surface localization indicates that recycling is rapid compared to their rate of endocytosis and targeting to late endosomes.

ENDO-LYSOSOMAL P2X RECEPTORS

P2X3 and P2X4 receptors also undergo rapid constitutive internalization from the plasma membrane, but unlike P2X1 they are predominantly localized to late endosomes and lysosomes (Bobanovic et al., 2002; Qureshi et al., 2007; Vacca et al., 2009). For P2X3 receptors this has been shown for heterologously expressed receptors in HEK293 cells and also for native receptors in dorsal root ganglion (DRG) neurons (Vacca et al., 2009). For P2X4 receptors, their localization to endolysosomes has been shown in immune and endothelial cells as well as for the heterologously expressed receptors in neurons (Bobanovic et al., 2002; Royle et al., 2002; Qureshi et al., 2007; Toulme et al., 2010). Constitutive endocytosis of P2X4 receptors is mediated via a dynamin/clathrin-dependent process and can be inhibited using dynasore (Bobanovic et al., 2002). Treatment with dynasore rapidly up-regulates P2X4 receptors at the surface of some but not all cells that express this receptor (Boumechache et al., 2009). In cultured microglial cells, native P2X4 receptors are rapidly up-regulated, indicating that they are continually cycling to and from the surface. In contrast, in bone marrow derived macrophages, total P2X4 receptor expression is high, but expression at the surface is very low and not increased by 1 h incubation with dynasore, suggesting that under basal conditions, receptors remain within endolysosomal compartments.

P2X3 receptors have a high rate of turnover, caused by rapid endocytosis and targeting to lysosomes, where the receptor is degraded (Vacca et al., 2009). In contrast, P2X4 receptors are surprisingly stable: they are resistant to degradation in the lysosome and they show little turnover over a period of 24 h (Qureshi et al., 2007). Their resistance to degradation is dependent upon multiple N-linked glycans which decorate the loop between the first and second transmembrane domains (TMD), which is predicted to face towards the lumen of lysosomes. Glycosylation is thought to play a similar role in protecting other lysosome-targeted proteins, such as Lamp-1, from degradation (Kundra and Kornfeld, 1999).

P2XR MOTIFS THAT DETERMINE RECEPTOR TRAFFICKING

P2X receptors share a YXXXK in the C-terminus which regulates surface expression (Chaumont et al., 2004). This motif is situated eight residues downstream of TMD2, and in P2X7 after an additional 18 amino acid cysteine-rich region. For P2X2 receptors, mutations around this motif reduce the stability of the receptor at the plasma membrane and increase internalization. Mutations within the motif similarly reduce the plasma membrane expression of other P2X receptor subtypes (Chaumont et al., 2004).

The P2X4 receptor has two tyrosine-based endocytic motifs within the C-terminus and one di-leucine-like motif within the N-terminus (Royle et al., 2002, 2005; Qureshi et al., 2007). Mutations of Y378, which forms part of a non-canonical YXXGΦ motif, substantially slow receptor endocytosis, suggesting that this is the more accessible of the two tyrosine-based motifs (Royle et al., 2002, 2005). These mutants still, however, show targeting to lysosomes, but this is further inhibited by mutating the leucine and isoleucine pair within the N-terminus.

Though the P2X3 receptor does not share any endocytic motifs with P2X4, there is a di-leucine motif in its C-terminal tail and also a consensus sequence for ubiquitination (DSGΦXS) that is suggested to be involved in the endocytosis and rapid degradation of the receptor (Vacca et al., 2009).

For P2X7 receptors, several mutations, deletions and naturally occurring single nucleotide polymorphisms (SNPs) within the long C-terminal tail have been shown to disrupt its normal trafficking to the plasma membrane. Attention has mostly focused on the distal end of the C-terminus of P2X7 (573–590), where there is a region with strong amino acid identity to the lipopolysaccharide (LPS) binding region of the LPS binding protein (Denlinger et al., 2001). Truncations and mutations within a region overlapping this site (551–581) disrupt normal receptor trafficking in HEK293 cells (Smart et al., 2003). For example, the I568N SNP in human P2X7 receptors disrupts normal trafficking and function (Wiley et al., 2003), as does substitution of acidic residues for the dibasic R578 and K579 (Denlinger et al., 2003). More recently it was shown that mutations within this region also disrupt the normal targeting of rat P2X7 receptors in polarized epithelial cells (Bradley et al., 2010). Alanine substitutions at P582–Q587 switched receptor targeting from the basolateral to the apical membrane but without disrupting plasma membrane expression or receptor function. Although site-directed mutagenesis has revealed critical residues in P2X7 receptor trafficking, the mechanism(s) underlying their involvement remains unknown.

REGULATION OF RECEPTOR TRAFFICKING AND MOBILITY

The activation of P2X receptors regulates their trafficking to and from the plasma membrane and their mobility within the plasma membrane in a calcium-dependent manner. Agonist-stimulated P2X receptor internalization and recycling back to the plasma membrane was first shown for P2X1 in rat vas deferens (Ennion and Evans, 2001). A later study of P2X1 mobility in HEK293 cells showed an increase in the rate of FRAP at the cell surface in the presence of agonist, which was dependent upon a rise in calcium, upon clathrin-mediated endocytosis and upon trafficking of vesicles back to the surface (Lalo et al., 2010). P2X1 receptors rapidly desensitize in the presence of agonist and inhibiting the internalization and recycling of receptors reduces the rate of recovery from desensitization. The P2X3 receptor also rapidly desensitizes and shows agonist-stimulated internalization (Vacca et al., 2009). P2X4 receptors desensitize more slowly but inhibiting dynamin-mediated endocytosis similarly slows the resensitization process (Murrell-Lagnado, unpublished). Thus, receptor retrieval and recycling appears to be important for maintaining the activity of the surface receptors.

Enhanced translocation of intracellular receptors to the plasma membrane has been shown to be a mechanism for up-regulating receptor function, particularly for those receptors that are predominantly intracellular. P2X4 receptors translocate from endolysosomes back to the surface, whereas for P2X3 receptors it is unclear whether up-regulation involves increased delivery from the secretory or endocytic pathway. ATP produces a transient increase in the number of P2X3 receptors at the surface causing sensitization of the current to repetitive doses (Vacca et al., 2009). Another example of increased trafficking of P2X3 receptors to the plasma membrane is in trigeminal sensory neurons in response to calcitonin gene-related peptide (CGRP; Fabbretti et al., 2006). A 1 h incubation with CGRP increased both the amplitude of P2X3 receptor currents and their rate of recovery from desensitization. For P2X4 receptors in macrophages, surface expression is increased in response to stimuli that promote lysosome exocytosis either by increasing cytosolic calcium or by alkalinization of the lysosomes (Qureshi et al., 2007). In the cerebellar microglial cell line, C8-B4, P2X4 receptor currents are negligible in resting cells, but after activating cells with either LPS or fibronectin, receptors translocate from lysosomes to the surface to enhance receptor-mediated currents (Toulme et al., 2010). Anti-depressants, which inhibit these currents, act by blocking this translocation process rather than by directly inhibiting the opening of the channel pore (Toulme et al., 2010). This mode of action could prove to be a useful way of selectively targeting the different subtypes with new therapeutics.

It is not only the retrieval and recycling of P2X receptors that is sensitive to agonist stimulation: their mobility within the lateral plane of the plasma membrane is promoted by the binding of ATP triggering a local influx of calcium. P2X2 receptor mobility was measured in hippocampal neurons by imaging single molecules using a quantum dot-based approach, and a similar approach was used with P2X4 receptors in microglia (Richler et al., 2011; Toulme and Khakh, 2012). In both cases two populations of receptor were observed, characterized as the

mobile and the slowly mobile pool. Neither population correlated with clusters of receptors or receptors localized in lipid rafts, so the molecular basis for the different mobility is unclear. Both populations showed increased mobility in response to ATP. For P2X2 receptors, mobility was also increased by the co-expression of VILIP-1 (Richler et al., 2011). The implications of this increased lateral mobility for receptor signaling remains to be established.

TARGETING OF P2X RECEPTORS TO LIPID RAFTS

The plasma membrane is an extremely heterogeneous environment and the trafficking and function of P2X receptors are regulated by the proteins and lipids within their immediate environment, with which they interact. Lipid rafts are commonly defined as microdomains rich in cholesterol, sphingolipids and saturated phospholipids, but there is heterogeneity amongst these domains in terms of their protein and lipid composition (Pike, 2004). While some proteins are targeted to rafts, others are excluded, affecting the nature of the signaling complexes formed and their stability within the membrane.

Lipid rafts are often identified biochemically by their low buoyant density in a sucrose density gradient, their resistance to solubilization in Triton-X 100 and the presence of protein markers such as caveolin-1 (Pike, 2004). Several of the P2X receptors have been shown to associate with lipid rafts, but the degree to which they distribute between the raft and non-raft fractions depends upon the cells in which they are expressed and the method used to prepare the rafts (Allsopp et al., 2010). P2X1-4 receptors expressed in HEK293 cells associate with rafts prepared using a detergent-free method (Allsopp et al., 2010). When rafts are instead prepared using Triton-X 100, the receptors shift to non-raft fractions. P2X1 and P2X2 receptors are more resistant to extraction from rafts by Triton-X 100 than are P2X3 and P2X4 receptors, suggesting that P2X1 and P2X2 interact more strongly with the cholesterol enriched domains (Allsopp et al., 2010).

Native P2X receptors have also been shown to target to lipid rafts. For P2X1 receptors, the distribution between raft and non-raft fractions is dependent upon the cell type. For example, P2X1 receptors in smooth muscle preparations from artery, vas deferens and bladder are almost exclusively in rafts, whereas only 20% of P2X1 receptors in platelets are in rafts (Vial and Evans, 2005; Vial et al., 2006). Native P2X3 receptors in trigeminal sensory neurons target to lipid rafts and, in a transgenic migraine mouse model, up-regulation of these receptors correlates with an increase in the abundance of lipid rafts and an increase in the fraction of P2X3 receptors within rafts (Gnanasekaran et al., 2011). Thus regulation of lipid rafts provides a mechanism for changing the functional expression of these receptors. P2X7 receptors are also found within rafts, both for heterologously-expressed receptors in HEK293 cells and for native receptors in rat submandibular glands, peritoneal macrophages and mouse lung alveolar cells (Garcia-Marcos et al., 2006a,b; Barth et al., 2007, 2008; Gonnord et al., 2009). Similar to P2X1-4, the association of P2X7 receptors with rafts is dependent upon the method used for isolating them. The receptors target to rafts prepared in detergent-free conditions, but this is reduced by low concentrations of Triton-X-100. This might reflect a weak association of P2X7 receptors with rafts or a difference in the nature of the rafts isolated by these methods.

Rafts prepared in the absence of detergent retain a greater fraction of inner leaflet-membrane lipids, in particular phosphatidylserine (Pike et al., 2002) and this might be involved in stabilizing the association with P2X7 receptors.

How P2X receptors target to rafts is unclear. For P2X7 receptors there is evidence to support the involvement of both caveolin-1 and palmitoylation of the receptor within its cytoplasmic C-terminal domain (Barth et al., 2007, 2008; Gonnord et al., 2009). P2X7 receptors palmitoylated with a radiolabeled palmitate were detected exclusively in lipid rafts, while inhibiting palmitoylation reduced receptor targeting to rafts. In mouse lung alveolar epithelial cells, P2X7R and caveolin-1 were detected in the same native complexes and caveolin-1 co-immunoprecipitated with P2X7 receptors (Weinhold et al., 2010). Also, epithelial cells from the caveolin-1 knock-out mice showed reduced levels of P2X7 immunostaining at the plasma membrane (Barth et al., 2007, 2008). A role for caveolin-1 could explain cell-type dependent differences in P2X7 receptor trafficking and function, such as between fibroblasts, rich in caveolin-1, and some immune cells, deficient in caveolin-1.

Depleting plasma membrane cholesterol with methyl- β -cyclodextrin disrupts lipid rafts and alters the function of some but not all of the P2X receptors. P2X1 receptor currents are strongly inhibited by cholesterol depletion, whereas P2X2 receptor currents are unchanged (Vial and Evans, 2005; Allsopp et al., 2010). A region within the N-terminus of P2X1 proximal to TMD1 was identified as an important determinant of cholesterol sensitivity (Allsopp et al., 2010). Cholesterol sensitivity of P2X1 receptor currents was, however, abolished by a cytoskeletal stabilizing agent, suggesting that lipid rafts regulate P2X1 by affecting its interaction with the cytoskeleton (Lalo et al., 2011). P2X3 receptor currents in trigeminal sensory neurons were inhibited by methyl- β -cyclodextrin treatment and desensitization was accelerated (Gnanasekaran et al., 2011). P2X4 receptor currents in Thp-1 monocytes were similarly inhibited by depleting cholesterol (Li and Fountain, 2012). The role of lipid rafts as regulators of P2X7 receptor signaling is of particular interest because receptor stimulation activates lipid-metabolizing enzymes, including phospholipases and sphingomyelinases, that reside in lipid rafts and whose substrates are also enriched in rafts. In cells from rat submandibular glands, cholesterol depletion reduced ATP-stimulated ceramide generation and phospholipase A2 activation, consistent with the idea that targeting to rafts controls signaling between the P2X7 receptor and its downstream effectors (Garcia-Marcos et al., 2006a).

SUMMARY AND OUTLOOK

For many of the P2X receptors we now have a basic understanding of their trafficking properties and subcellular distributions, and in some cases have identified regulators that can alter their trafficking to and from the plasma membrane. For the P2X4 receptor, a key unanswered question is why it stably resides within endolysosomes. Only for the type II alveolar cells has a role for P2X4 receptors in the lamellar bodies, which resemble secretory lysosomes, been demonstrated, and only upon their fusion with the plasma membrane (Miklavc et al., 2011). It remains to be established whether P2X4 receptors have additional roles within conventional

or secretory lysosomes of other cells. P2X7 receptors are known to be up-regulated in many different cell types under inflammatory conditions, contributing to pathology (Lister et al., 2007). We would like to understand what regulates the trafficking of these receptors to and from the cell surface. For all of the P2X receptors we would like to understand how the lipid environment controls their function and stability. For P2X7 receptors this interaction might also operate in the reverse direction: events downstream of P2X7 receptor activation, including sphingomyelinase activation and ceramide generation directly modify raft structure, thereby providing the potential for cross-talk with other receptors that are modulated by lipid rafts.

REFERENCES

- Allsopp, R. C., Lalo, U., and Evans, R. J. (2010). Lipid raft association and cholesterol sensitivity of P2X1-4 receptors for ATP: chimeras and point mutants identify intracellular amino-terminal residues involved in lipid regulation of P2X1 receptors. *J. Biol. Chem.* 285, 32770–32777. doi: 10.1074/jbc.m110.148940
- Baines, A., Parkinson, K., Sim, J. A., Bragg, L., Thompson, C. R., and North, R. A. (2013). Functional properties of five Dictyostelium discoideum P2X receptors. *J. Biol. Chem.* 288, 20992–21000. doi: 10.1074/jbc.m112.445346
- Barrera, N. P., Henderson, R. M., Murrell-Lagnado, R. D., and Edwardson, J. M. (2007). The stoichiometry of P2X2/6 receptor heteromers depends on relative subunit expression levels. *Biophys. J.* 93, 505–512. doi: 10.1529/biophysj.106.101048
- Barrera, N. P., Ormond, S. J., Henderson, R. M., Murrell-Lagnado, R. D., and Edwardson, J. M. (2005). Atomic force microscopy imaging demonstrates that P2X2 receptors are trimers but that P2X6 receptor subunits do not oligomerize. *J. Biol. Chem.* 280, 10759–10765. doi: 10.1074/jbc.m412265200
- Barth, K., Weinhold, K., Guenther, A., Linge, A., Gereke, M., and Kasper, M. (2008). Characterization of the molecular interaction between caveolin-1 and the P2X receptors 4 and 7 in E10 mouse lung alveolar epithelial cells. *Int. J. Biochem. Cell Biol. Cell Biol.* 40, 2230–2239. doi: 10.1016/j.biocel.2008.03.001
- Barth, K., Weinhold, K., Guenther, A., Young, M. T., Schnittler, H., and Kasper, M. (2007). Caveolin-1 influences P2X7 receptor expression and localization in mouse lung alveolar epithelial cells. *FEBS J.* 274, 3021–3033. doi: 10.1111/j.1742-4658.2007.05830.x
- Bo, X., Jiang, L. H., Wilson, H. L., Kim, M., Burnstock, G., Surprenant, A., et al. (2003). Pharmacological and biophysical properties of the human P2X5 receptor. *Mol. Pharmacol.* 63, 1407–1416. doi: 10.1124/mol.63.6.1407
- Bobanovic, L. K., Royle, S. J., and Murrell-Lagnado, R. D. (2002). P2X receptor trafficking in neurons is subunit specific. *J. Neurosci.* 22, 4814–4824.
- Boumehache, M., Masin, M., Edwardson, J. M., Gorecki, D. C., and Murrell-Lagnado, R. (2009). Analysis of assembly and trafficking of native p2x4 and p2x7 receptor complexes in rodent immune cells. *J. Biol. Chem.* 284, 13446–13454. doi: 10.1074/jbc.m901255200
- Bradley, H. J., Liu, X., Collins, V., Owid, G., Goli, G. R., Smith, M., et al. (2010). Identification of an intracellular microdomain of the P2X7 receptor that is crucial in basolateral membrane targeting in epithelial cells. *FEBS Lett.* 584, 4740–4744. doi: 10.1016/j.febslet.2010.11.007
- Chaumont, S., Compan, V., Toulme, E., Richler, E., Housley, G. D., Rassendren, F., et al. (2008). Regulation of P2X2 receptors by the neuronal calcium sensor VILIP1. *Sci. Signal.* 1, ra8. doi: 10.1126/scisignal.1162329
- Chaumont, S., Jiang, L. H., Penna, A., North, R. A., and Rassendren, F. (2004). Identification of a trafficking motif involved in the stabilization and polarization of P2X receptors. *J. Biol. Chem.* 279, 29628–29638. doi: 10.1074/jbc.m403940200
- Compan, V., Ullmann, L., Stelmashenko, O., Chemin, J., Chaumont, S., and Rassendren, F. (2012). P2X2 and P2X5 subunits define a new heteromeric receptor with P2X7-like properties. *J. Neurosci.* 32, 4284–4296. doi: 10.1523/jneurosci.6332-11.2012
- Denlinger, L. C., Fiset, P. L., Sommer, J. A., Watters, J. J., Prabhu, U., Dubyak, G. R., et al. (2001). Cutting edge: the nucleotide receptor P2X7 contains multiple protein- and lipid-interaction motifs including a potential binding site for bacterial lipopolysaccharide. *J. Immunol.* 167, 1871–1876.
- Denlinger, L. C., Sommer, J. A., Parker, K., Gudipaty, L., Fiset, P. L., Watters, J. W., et al. (2003). Mutation of a dibasic amino acid motif within the C terminus of the P2X7 nucleotide receptor results in trafficking defects and impaired function. *J. Immunol.* 171, 1304–1311.
- Ennion, S. J., and Evans, R. J. (2001). Agonist-stimulated internalisation of the ligand-gated ion channel P2X(1) in rat vas deferens. *FEBS Lett.* 489, 154–158. doi: 10.1016/s0014-5793(01)02102-0
- Fabbretti, E., D'Arco, M., Fabbro, A., Simonetti, M., Nistri, A., and Giniatullin, R. (2006). Delayed upregulation of ATP P2X3 receptors of trigeminal sensory neurons by calcitonin gene-related peptide. *J. Neurosci.* 26, 6163–6171. doi: 10.1523/jneurosci.0647-06.2006
- Fountain, S. J., Parkinson, K., Young, M. T., Cao, L., Thompson, C. R., and North, R. A. (2007). An intracellular P2X receptor required for osmoregulation in Dictyostelium discoideum. *Nature* 448, 200–203. doi: 10.1038/nature05926
- Garcia-Marcos, M., Perez-Andres, E., Tandel, S., Fontanils, U., Kumps, A., Kabre, E., et al. (2006a). Coupling of two pools of P2X7 receptors to distinct intracellular signaling pathways in rat submandibular gland. *J. Lipid Res.* 47, 705–714. doi: 10.1194/jlr.m500408-jlr200
- Garcia-Marcos, M., Pochet, S., Tandel, S., Fontanils, U., Astigarraga, E., Fernandez-Gonzalez, J. A., et al. (2006b). Characterization and comparison of raft-like membranes isolated by two different methods from rat submandibular gland cells. *Biochim. Biophys. Acta* 1758, 796–806. doi: 10.1016/j.bbammem.2006.05.008
- Gnanasekaran, A., Sundukova, M., van den Maagdenberg, A. M., Fabbretti, E., and Nistri, A. (2011). Lipid rafts control P2X3 receptor distribution and function in trigeminal sensory neurons of a transgenic migraine mouse model. *Mol. Pain* 7:77. doi: 10.1186/1744-8069-7-77
- Gonnord, P., Delarasse, C., Auger, R., Benihoud, K., Prigent, M., Cuif, M. H., et al. (2009). Palmitoylation of the P2X7 receptor, an ATP-gated channel, controls its expression and association with lipid rafts. *FASEB J.* 23, 795–805. doi: 10.1096/fj.08-114637
- Gu, B. J., Zhang, W. Y., Bendall, L. J., Chessell, I. P., Buell, G. N., and Wiley, J. S. (2000). Expression of P2X(7) purinoceptors on human lymphocytes and monocytes: evidence for nonfunctional P2X(7) receptors. *Am. J. Physiol. Cell Physiol.* 279, C1189–C1197.
- Gudipaty, L., Humphreys, B. D., Buell, G., and Dubyak, G. R. (2001). Regulation of P2X(7) nucleotide receptor function in human monocytes by extracellular ions and receptor density. *Am. J. Physiol. Cell Physiol.* 280, C943–C953.
- Hickman, S. E., el Khoury, J., Greenberg, S., Schieren, I., and Silverstein, S. C. (1994). P2Z adenosine triphosphate receptor activity in cultured human monocyte-derived macrophages. *Blood* 84, 2452–2456.
- Kotnis, S., Bingham, B., Vasilyev, D. V., Miller, S. W., Bai, Y., Yeola, S., et al. (2010). Genetic and functional analysis of human P2X5 reveals a distinct pattern of exon 10 polymorphism with predominant expression of the nonfunctional receptor isoform. *Mol. Pharmacol.* 77, 953–960. doi: 10.1124/mol.110.063636
- Kundra, R., and Kornfeld, S. (1999). Asparagine-linked oligosaccharides protect Lamp-1 and Lamp-2 from intracellular proteolysis. *J. Biol. Chem.* 274, 31039–31046. doi: 10.1074/jbc.274.43.31039
- Lalo, U., Allsopp, R. C., Mahaut-Smith, M. P., and Evans, R. J. (2010). P2X1 receptor mobility and trafficking: regulation by receptor insertion and activation. *J. Neurochem.* 113, 1177–1187. doi: 10.1111/j.1471-4159.2010.06730.x
- Lalo, U., Jones, S., Roberts, J. A., Mahaut-Smith, M. P., and Evans, R. J. (2012). Heat shock protein 90 inhibitors reduce trafficking of ATP-gated P2X1 receptors and human platelet responsiveness. *J. Biol. Chem.* 287, 32747–32754. doi: 10.1074/jbc.m112.376566
- Lalo, U., Roberts, J. A., and Evans, R. J. (2011). Identification of human P2X1 receptor-interacting proteins reveals a role of the cytoskeleton in receptor regulation. *J. Biol. Chem.* 286, 30591–30599. doi: 10.1074/jbc.m111.253153
- Li, J., and Fountain, S. J. (2012). Fluvastatin suppresses native and recombinant human P2X4 receptor function. *Purinergic Signal.* 8, 311–316. doi: 10.1007/s11302-011-9289-9
- Lister, M. F., Sharkey, J., Sawatzky, D. A., Hodgkiss, J. P., Davidson, D. J., Rossi, A. G., et al. (2007). The role of the purinergic P2X7 receptor in inflammation. *J. Inflamm. (Lond.)* 4:5. doi: 10.1186/1476-9255-4-5
- Ludlow, M. J., Durai, L., and Ennion, S. J. (2009). Functional characterization of intracellular Dictyostelium discoideum P2X receptors. *J. Biol. Chem.* 284, 35227–35239. doi: 10.1074/jbc.M109.045674
- Masin, M., Kerschensteiner, D., Dumke, K., Rubio, M. E., and Soto, F. (2006). Fe65 interacts with P2X2 subunits at excitatory synapses and modulates receptor function. *J. Biol. Chem.* 281, 4100–4108. doi: 10.1074/jbc.m507735200

- Miklavc, P., Mair, N., Wittekindt, O. H., Haller, T., Dietl, P., Felder, E., et al. (2011). Fusion-activated Ca²⁺ entry via vesicular P2X₄ receptors promotes fusion pore opening and exocytotic content release in pneumocytes. *Proc. Natl. Acad. Sci. U S A* 108, 14503–14508. doi: 10.1073/pnas.1101039108
- Ormond, S. J., Barrera, N. P., Qureshi, O. S., Henderson, R. M., Edwardson, J. M., and Murrell-Lagnado, R. D. (2006). An uncharged region within the N terminus of the P2X₆ receptor inhibits its assembly and exit from the endoplasmic reticulum. *Mol. Pharmacol.* 69, 1692–1700. doi: 10.1124/mol.105.020404
- Pike, L. J. (2004). Lipid rafts: heterogeneity on the high seas. *Biochem. J.* 378(Pt 2), 281–292. doi: 10.1042/bj20031672
- Pike, L. J., Han, X., Chung, K. N., and Gross, R. W. (2002). Lipid rafts are enriched in arachidonic acid and plasmenylethanolamine and their composition is independent of caveolin-1 expression: a quantitative electrospray ionization/mass spectrometric analysis. *Biochemistry* 41, 2075–2088. doi: 10.1021/bi0156557
- Qureshi, O. S., Paramasivam, A., Yu, J. C., and Murrell-Lagnado, R. D. (2007). Regulation of P2X₄ receptors by lysosomal targeting, glycan protection and exocytosis. *J. Cell Sci.* 120(Pt 21), 3838–3349. doi: 10.1242/jcs.010348
- Richler, E., Shigetomi, E., and Khakh, B. S. (2011). Neuronal P2X₂ receptors are mobile ATP sensors that explore the plasma membrane when activated. *J. Neurosci.* 31, 16716–16730. doi: 10.1523/jneurosci.3362-11.2011
- Royle, S. J., Bobanovic, L. K., and Murrell-Lagnado, R. D. (2002). Identification of a non-canonical tyrosine-based endocytic motif in an ionotropic receptor. *J. Biol. Chem.* 277, 35378–35385. doi: 10.1074/jbc.m204844200
- Royle, S. J., Qureshi, O. S., Bobanovic, L. K., Evans, P. R., Owen, D. J., and Murrell-Lagnado, R. D. (2005). Non-canonical YXXGPhi endocytic motifs: recognition by AP2 and preferential utilization in P2X₄ receptors. *J. Cell Sci.* 118(Pt 14), 3073–3080. doi: 10.1242/jcs.02451
- Shrivastava, A. N., Triller, A., Sieghart, W., and Sarto-Jackson, I. (2011). Regulation of GABA(A) receptor dynamics by interaction with purinergic P2X₂ receptors. *J. Biol. Chem.* 286, 14455–14468. doi: 10.1074/jbc.m110.165282
- Sivaramakrishnan, V., and Fountain, S. J. (2012a). A mechanism of intracellular P2X receptor activation. *J. Biol. Chem.* 287, 28315–28326. doi: 10.1074/jbc.m112.372565
- Sivaramakrishnan, V., and Fountain, S. J. (2012b). Intracellular P2X receptors as novel calcium release channels and modulators of osmoregulation in Dictyostelium: a comparison of two common laboratory strains. *Channels (Austin)* 7, 43–46. doi: 10.4161/chan.22737
- Smart, M. L., Gu, B., Panchal, R. G., Wiley, J., Cromer, B., Williams, D. A., et al. (2003). P2X₇ receptor cell surface expression and cytolytic pore formation are regulated by a distal C-terminal region. *J. Biol. Chem.* 278, 8853–8860. doi: 10.1074/jbc.m211094200
- Toulme, E., Garcia, A., Samways, D., Egan, T. M., Carson, M. J., and Khakh, B. S. (2010). P2X₄ receptors in activated C8-B4 cells of cerebellar microglial origin. *J. Gen. Physiol.* 135, 333–353. doi: 10.1085/jgp.200910336
- Toulme, E., and Khakh, B. S. (2012). Imaging P2X₄ receptor lateral mobility in microglia: regulation by calcium and p38 MAPK. *J. Biol. Chem.* 287, 14734–14748. doi: 10.1074/jbc.m111.329334
- Vacca, F., Giustizieri, M., Ciotti, M. T., Mercuri, N. B., and Volonte, C. (2009). Rapid constitutive and ligand-activated endocytic trafficking of P2X receptor. *J. Neurochem.* 109, 1031–1041. doi: 10.1111/j.1471-4159.2009.06029.x
- Vial, C., and Evans, R. J. (2005). Disruption of lipid rafts inhibits P2X₁ receptor-mediated currents and arterial vasoconstriction. *J. Biol. Chem.* 280, 30705–30711. doi: 10.1074/jbc.m504256200
- Vial, C., Fung, C. Y., Goodall, A. H., Mahaut-Smith, M. P., and Evans, R. J. (2006). Differential sensitivity of human platelet P2X₁ and P2Y₁ receptors to disruption of lipid rafts. *Biochem. Biophys. Res. Commun.* 343, 415–419. doi: 10.1016/j.bbrc.2006.02.174
- Weinhold, K., Krause-Buchholz, U., Rodel, G., Kasper, M., and Barth, K. (2010). Interaction and interrelation of P2X₇ and P2X₄ receptor complexes in mouse lung epithelial cells. *Cell. Mol. Life Sci.* 67, 2631–2642. doi: 10.1007/s00018-010-0355-1
- Wiley, J. S., Dao-Ung, L. P., Li, C., Shemon, A. N., Gu, B. J., Smart, M. L., et al. (2003). An Ile-568 to Asn polymorphism prevents normal trafficking and function of the human P2X₇ receptor. *J. Biol. Chem.* 278, 17108–17113. doi: 10.1074/jbc.m212759200

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 01 October 2013; accepted: 07 November 2013; published online: 22 November 2013.

Citation: Robinson LE and Murrell-Lagnado RD (2013) The trafficking and targeting of P2X receptors. *Front. Cell. Neurosci.* 7:233. doi: 10.3389/fncel.2013.00233
This article was submitted to the journal *Frontiers in Cellular Neuroscience*.

Copyright © 2013 Robinson and Murrell-Lagnado. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.